

Non-innocent radical ion intermediates in photoredox catalysis: parallel reduction modes enable coupling of diverse aryl chlorides

Alyah F. Chmiel,¹ Oliver P. Williams,¹ Colleen P. Chernowsky,¹ Charles S. Yeung,^{*,2} and Zachary K. Wickens^{*,1}

¹Department of Chemistry, University of Wisconsin-Madison; Madison, Wisconsin, 53706, United States.

²Discovery Chemistry, Merck & Co., Inc., 33 Avenue Louis Pasteur, Boston, MA 02115, United States

ABSTRACT: We describe a photocatalytic system that elicits potent photoreductant activity from conventional photocatalysts by leveraging radical anion intermediates generated *in situ*. The combination of isophthalonitrile and sodium formate promotes diverse aryl radical coupling reactions from abundant but difficult to reduce aryl chloride substrates. Mechanistic studies reveal two parallel pathways for substrate reduction both enabled by a key terminal reductant byproduct, carbon dioxide radical anion.

Reductive activation of organic molecules via single electron transfer (SET) is a fundamental elementary step that underpins diverse and powerful synthetic transformations.¹⁻⁴ Photoredox catalysis promotes SET through conversion of energy from visible light into chemical redox potential and has enabled a suite of carbon-carbon and carbon-heteroatom bond-forming reactions.⁵⁻⁹ When considering whether a substrate will be suitable for photoredox reduction, two primary catalyst parameters are initially considered: (1) $E_{1/2}(\text{PC}^+/\text{PC}^*)$ and (2) $E_{1/2}(\text{PC}/\text{PC}^{\cdot-})$.^{7,10,11} These values reflect redox potentials bounded by the energy of photons in the visible region, a limitation compounded by energy losses to intersystem crossing.¹² As a result, many abundant but challenging to reduce substrates are excluded from photoredox activation based on these guidelines (Figure 1A).¹³⁻¹⁵

Aryl radicals are reactive intermediates that engage in a myriad of synthetically valuable transformations.¹⁶⁻¹⁸ Classically, aryl radical intermediates are generated from aryl diazonium salts, iodides, or bromides.¹⁹⁻²⁹ Aryl chlorides are rarely used as radical precursors despite the fact they comprise over two thirds³⁰ of commercially available aryl halides (Figure 1B).³¹⁻³⁵ This is a consequence of their resistance to reductive activation,¹³ and high $\text{C}(\text{sp}^2)\text{-Cl}$ BDE.^{36,37} König recently pioneered an elegant strategy, termed consecutive photoinduced electron transfer (conPET), wherein a photochemically generated radical anion is subsequently excited.^{38,39} This approach primes the photocatalyst with an electron prior to excitation and, in principle, can generate much deeper reduction potentials through $E_{1/2}(\text{PC}/\text{PC}^{\cdot-})$. Indeed, later implementations of this conPET strategy unlocked exceptionally challenging reductions.^{40,41} However, all recent advances in visible light photoredox methods that reduce electronically diverse chloroarenes have been limited to proteodefunctionalization and borylation reactions.^{40,42-45} Recent electrophotocatalytic⁴⁶⁻⁴⁹ approaches have directly generated these electron-primed photocatalysts cathodically.^{50,51} While this strategy has begun to expand the range of radical coupling reactions that engage aryl chlorides, a general approach to leverage the expansive pool of aryl chloride substrates in radical couplings has remained elusive and the need for electrochemical equipment remains a barrier in some




Figure 1. (A) Energy limitations for photoredox catalysis and electron-primed photoredox catalysis. (B) Aryl chlorides abundance and reactivity as aryl radical precursors. (C) Strategy employing chemical reductants to exploit electron-primed photoredox catalysis. All V vs. SCE.

settings.^{52,53} In particular, net-reductive radical coupling processes, such as alkene hydroarylation⁵⁴⁻⁵⁶, have remained elusive for aryl chloride substrates for all modern methods. We suspect that the paucity of net-reductive processes is a consequence of the intrinsic challenges of circumventing premature reduction of the aryl radical intermediate ($E_{\text{red}}(\text{Ph}^{\cdot}/\text{Ph}^-) = +0.05 \text{ V vs. SCE}$)⁵⁷ in the presence of a stoichiometric reductant.

Our group recently used electrochemistry to examine a diverse set of organic radical anions for photocatalytic activity in the reductive cleavage of strong $\text{C}(\text{sp}^2)\text{-O}$ and $\text{C}(\text{sp}^2)\text{-N}$ bonds.⁵⁸ These experiments revealed that numerous radical

anions, including those derived from commonly employed photoredox catalysts, can serve as potent photocatalytic reductants upon cathodic reduction. These data fit into a growing body of literature from our group⁵⁹ and others^{40,60} that suggest photocatalyst-based redox events can engender more potent activity from conventional photocatalysts. Taken together, these data led us to consider whether we could redesign a photocatalytic system to favor formation of photoactive radical anion intermediates to elicit deeply reducing potentials and expand the repertoire of coupling reactions available from aryl chlorides under operationally simple conditions (Figure 1C). Herein, we disclose that selection of an appropriate reductant to generate and maintain an active electron-primed photoredox catalyst *in situ* enables reduction potentials far beyond those expected from conventional catalyst selection criteria. These new reduction conditions promote a diverse array of intermolecular coupling reactions, including net-reductive coupling processes, from readily available aryl chloride substrates.

We first evaluated a suite of organic compounds recently found to possess photocatalytically active radical anion congeners¹ for activity in the dehalogenation of PhCl ($E_{\text{red}} = -2.8$ V vs. SCE). Considering only conventional photoredox catalyst selection parameters (PC/PC⁻ and PC^{*}/PC⁺), this reduction would be exceedingly endothermic (>1 V) for the photocatalysts under investigation. Therefore, activity in this assay would implicate *in situ* formation of a distinct and potent reductant. Initially, we examined a range of trialkylamine reductants because these are common reductants in photoredox catalysis, including in conPET strategies (Figure 2A).^{38,40,62} We found that each catalyst modestly promoted this energetically demanding dehalogenation reaction. The isophthalonitrile catalysts, which are both excellent neutral chromophores⁶³ as well as electron-primed photoredox catalysts,⁵⁸ promoted the reaction most efficiently albeit still in poor yield. To exclude hydrogen atom transfer (XAT) aryl radical generation,⁶⁴⁻⁶⁶ we examined the reductive defunctionalization of anilinium and aryl phosphate salts (Table 1). These are each challenging reductive cleavage reactions^{67,68} of non-polarizable leaving groups unlikely to undergo XAT processes. We found both substrates underwent productive defunctionalization, albeit in diminished yield (Figure S3).

A broader survey of reductants less commonly employed in photoredox catalysis (Figure S4) revealed that sodium formate

substantially enhanced the photoreductant activity of **4-DPAIPN** (Figure 2A). We suspect this improvement occurs because formate salts undergo a second-order hydrogen atom transfer (HAT) process upon oxidation⁶⁹ that results in formic acid and carbon dioxide radical anion.⁷⁰⁻⁷³ As a consequence, a second reducing equivalent is liberated from formate after initial oxidation. We suspect that the carbon dioxide radical anion can either reduce another equivalent of photocatalyst or promote the reaction by direct reduction of substrate ($E_{1/2}(\text{CO}_2/\text{CO}_2^{\cdot-}) = -2.2$ V vs SCE).⁷⁴ In each mechanistic manifold, the SET is rendered irreversible by the release of CO₂ gas. This scenario contrasts starkly with the trialkylamine reductants, which result in oxidizing amine radical cation intermediates ($E_{1/2}(\text{NR}_3/\text{NR}_3^{\cdot+}) < 1$ V vs SCE) that could deactivate the radical anion photocatalyst via back electron transfer (Figure 2B).^{62,75,76}

We next evaluated the potency of this new catalytic system. Having established that chlorobenzene could be reduced (-2.8 V vs SCE), we probed dehalogenation of increasingly electron-rich aryl chlorides. These experiments revealed that substrates with reduction potentials as low as -3.4 V vs. SCE are efficiently reduced. Additionally, these conditions promoted the challenging reductive cleavage of both an anilinium and aryl phosphate substrate. Taken together, these data clearly implicate processes beyond a conventional photoredox manifold. For example, the reduction of **6** would be predicted to be endothermic by nearly 2 V (>40 kcal/mol at room temperature) based on the most reducing conventional redox couple of **4-DPAIPN** ($E_{1/2}(\text{PC}/\text{PC}^{\cdot-}) = -1.5$ V vs SCE).⁷⁷ We next subjected radical clock **9** to this formate-driven system to validate the intermediacy of an aryl radical. As anticipated, these conditions furnished the five-membered ring product in high selectivity for radical cyclization. Despite its exceptionally reducing potentials, we suspected that this operationally simple procedure would be amenable to high-throughput techniques widely employed in medicinal chemistry. To this end, we rapidly evaluated the tolerance of complex drug-like scaffolds using a commercially available informer plate designed to challenge modern cross-coupling technology. We found that not only was photocatalytic activity retained in the well-plate format but that several of these functional group rich molecules were effectively transformed (Figure S6).







Figure 2. (A) Unlocking radical anion photocatalyst reactivity by evaluation of reductant for catalyst activation. Reactions were conducted on 0.1 mmol scale with 10 mol % **4-DPAIPN** and 3 equiv NaCHO₂. Reactions were analyzed via gas chromatography. (B) Overview of key considerations for chemical reductants as catalyst activators. All V vs. SCE.

Table 1. Evaluation of catalytic system.^{a,b,c}

Although, in principle, aryl radicals are highly versatile synthetic intermediates, premature reduction precludes radical coupling reactions in many cases. This is particularly problematic when potent reductants are required. To evaluate selectivity for radical coupling, we targeted redox-neutral photo-Arbusov and radical borylation processes. These established aryl radical transformations produce biologically relevant aryl phosphonates⁷⁸ and versatile organoboron products.⁷⁹ In both cases, we found that chloroarene substrates readily underwent the desired radical coupling process.⁸⁰ We found that both difficult to reduce electron-rich aryl chlorides and substrates bearing potentially reducible functional groups such as esters and amides were well-tolerated (Table 2). Furthermore, the catalytic system tolerated medically relevant heterocycles.

Next, we evaluated the reductive hydroarylation of alkenes. This challenging aryl radical reaction requires precise control over the relative rates of radical coupling versus protodehalogenation. HAT is mechanistically required to furnish product and cannot be simply suppressed. Initially, we targeted the synthesis of aryethylamines via hydroarylation. Recently, Jui and coworkers reported that aryl radical intermediates productively couple with vinyl carbamates to produce the aryethylamine pharmacophore.⁵⁶ Although one of the most reducing conventional photocatalysts was employed, the majority of the reaction scope was comprised of aryl iodide substrates and only aryl chloride substrates bearing withdrawing groups were viable. Intriguingly, we found that although the vinylcarbamate substrate

Table 2. Intermolecular couplings from challenging aryl chloride precursors.^a**Table 3.** Reductant Impact on Selectivity^h

^aReactions were run on 0.4 mmol scale. Isolated yield unless otherwise noted. ^bNMR yield. ^cGC yield. ^dReactions were run with 4-DPAIPN (12-15 mol %), NaCHO₂ (3 equiv), P(OEt)₃ (5 equiv). ^eReactions were run with 4-DPAIPN (5 mol %), NaCHO₂ (3 equiv), B₂pin₂ (3 equiv) and Cs₂CO₃ (3 equiv). ^fReactions were run with 4-DPAIPN (6 mol %), NaCHO₂ (3 equiv), N-vinyl carbamate (2.5 equiv) and CySH (5 mol %). ^gReactions were run with 4-DPAIPN (6 mol %), NaCHO₂ (3 equiv), alkene (5 equiv) and CySH (5 mol %). ^hReactions were run with conditions d-g, using either EtN(iPr)₂ or NaCHO₂ as the reductant. See SI for details.




Figure 3. (A) UV-vis spectrum demonstrating 4-DPAIPN⁻ generation in the presence of sodium formate and light. (B) UV-vis spectrum demonstrating that 4-DPAIPN⁻ reverts to 4-DPAIPN upon exposure to PhCl and subsequent irradiation.


is thermodynamically easier to reduce than most chloroarenes ($E_{\text{red}} = -2.2$ V vs. SCE), these potent reductive conditions selectively transformed chlorobenzene into *N*-Boc phenethylamine in high yield. Even as the gap between the chloroarene and vinyl carbamate coupling partner widens, synthetically useful arylethylamine yields are still observed. Similar to the other radical couplings, we found aryl chlorides bearing reductively sensitive functional groups were tolerated. We also found these conditions promoted the coupling of aryl chlorides and unactivated alkenes despite the fact that such a hydroarylation remains challenging with any aryl radical precursor.⁸¹ Finally, we questioned whether formate was uniquely effective for each of these radical coupling reactions or whether alkylamines were suitable terminal reductants. While EtN(*i*-Pr)₂ and 4-DPAIPN promote photoreduction of chlorobenzene, both reactivity and radical selectivity were diminished in each coupling reaction. Of note, competitive proteodehalogenation nearly precluded net-reductive hydroarylation when alkylamines were used.⁸²

Having established a generally applicable catalytic system to engage aryl chloride substrates in radical coupling reactions, we next conducted a preliminary mechanistic investigation into the process. First, we probed whether an electron-primed photoredox mechanism—wherein the 4-DPAIPN radical anion is generated and subsequently excited—was feasible under these conditions. We irradiated a mixture of 4-DPAIPN and sodium formate while monitoring speciation by UV-Vis (Figure 3A). This resulted in a decrease in 4-DPAIPN features and growth of new features consistent with electrochemically reduced 4-DPAIPN (Figure S10). Next, we probed the photoreduction of aryl chlorides. Chlorobenzene was added to the reaction mixture and, upon irradiation, the UV-Vis features of 4-DPAIPN were restored (Figure 3B). As expected based on the >1 V underpotential, no return of 4-DPAIPN was observed upon addition of chlorobenzene to 4-DPAIPN radical anion in the absence of light. Consistent with this mechanistic picture, Stern-Volmer analysis resulted in no measurable quenching of excited 4-DPAIPN by chlorobenzene. In contrast, formate salts did quench the excited state. Cyclohexane thiol, which was added to the net-reductive transformations as an HAT co-catalyst, also quenches the excited state and likely mediates the electron-

transfer events in these systems by an analogous mechanism (see Figure S8). Taken together, these experiments are consistent with our working hypothesis that photooxidation of formate results in the 4-DPAIPN radical anion which can be subsequently excited to photoreduce chlorobenzene and return 4-DPAIPN.

While the UV-Vis experiments indicate that an electron-primed photoredox mechanism is feasible, we recognized that carbon dioxide radical anion is sufficiently reducing ($E_{\text{red}}(\text{CO}_2/\text{CO}_2^{\bullet-}) = -2.2$ V vs SCE)⁷⁴ to promote reductive fragmentation of some of the aryl chloride substrates studied without the intervention of an electron-primed photoredox manifold. To evaluate the relative contribution of direct substrate reduction by $\text{CO}_2^{\bullet-}$, we attempted to generate this radical anion directly by homolysis of (PhS)₂ under 370 nm light in the absence of 4-DPAIPN.⁸³ We envisioned thiyl radical could abstract a hydrogen atom from formate to directly generate $\text{CO}_2^{\bullet-}$ *in situ* (Figure S11).^{72,84,85} These conditions resulted in quantitative conversion of 4-chlorobenzonitrile ($E_{\text{red}} = -2.1$ V vs SCE). However, only 9% conversion of chlorobenzene ($E_{\text{red}} = -2.8$ V vs SCE) and <5% conversion of chloroanisole ($E_{\text{red}} = -2.9$ V vs SCE) were observed under these photocatalyst-free

Scheme 1. Plausible mechanism for aryl chloride reduction and the roles of formate and its by-products.



conditions. In contrast, all three of substrates are dehalogenated in comparable efficiency through use of the **4-DPAIPN** conditions. We suspect both mechanisms operate in parallel for substrates within the bound of the potency of $\text{CO}_2^{\cdot-}$ (-2.2 V vs SCE) but that an electron-primed photoredox manifold supports more thermodynamically demanding reductions. In both cases, the $\text{CO}_2^{\cdot-}$ reductant byproduct plays an active role either (a) reducing the substrate directly or (b) reducing the photocatalyst to reactivate it without requiring persistent multiphoton excitation (Scheme 1).

Overall, we have illustrated that use of a formate-based terminal reductant in combination with an isophthalonitrile photocatalyst can engage aryl chlorides in diverse synthetically useful coupling reactions. We anticipate that these operationally simple reaction conditions comprise a broadly useful approach to photochemically induce difficult reductive processes. Beyond the immediate synthetic utility, these results are important because they challenge the notion that the terminal reductant can be viewed as merely an electron-source to turn over the photocatalyst. These data fit within a growing body of literature that suggests terminal reductant byproducts can play a non-innocent role in photoredox catalysis.^{65,66,75} We suspect that these results could also offer an alternative explanation for recent examples wherein isophthalonitrile catalysts have appeared to reduce substrates beyond their expected redox potentials^{66,86} and, more broadly, illustrate the importance of radical ion intermediates in photoredox catalysis.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures, characterization data, and spectra (PDF)

AUTHOR INFORMATION

Corresponding Authors

* Zachary K. Wickens – Department of Chemistry, University of Wisconsin–Madison, Madison, Wisconsin 53706, United States; orcid.org/0000-0002-5733-5288; Email: wickens@wisc.edu

* Charles S. Yeung – Discovery Chemistry, Merck & Co., Inc., 33 Avenue Louis Pasteur, Boston, MA 02115, United States; orcid.org/0000-0001-7320-7190; email: charles.yeung@merck.com

ACKNOWLEDGMENT

We thank Prof. Alison Wendlandt for helpful suggestions. We thank the Stahl, Weix, Yoon, and Schomaker groups for sharing their chemical inventory. We thank Dr. Wesley Swords (Yoon group) for assistance with Stern-Volmer quenching experiments. We thank Sara Alektiar for cyclic voltammetry data. This work was financially supported by the Office of the Vice Chancellor for Research and Graduate Education at the University of Wisconsin–Madison with funding from the Wisconsin Alumni Research Foundation. Acknowledgement is made to the

Donors of the American Chemical Society Petroleum Research Fund for partial funding of this research (60677-DN11). Spectroscopic instrumentation was supported by a generous gift from Paul J. and Margaret M. Bender, NSF (CHE-1048642), and NIH (S10OD012245 and 1S10OD020022-1).

REFERENCES

- (1) Ashby, E. C. Single-Electron Transfer, a Major Reaction Pathway in Organic Chemistry. An Answer to Recent Criticisms. *Acc. Chem. Res.* **1988**, *21* (11), 414–421.
- (2) Zhang, N.; Samanta, S. R.; Rosen, B. M.; Percec, V. Single Electron Transfer in Radical Ion and Radical-Mediated Organic, Materials and Polymer Synthesis. *Chem. Rev.* **2014**, *114* (11), 5848–5958.
- (3) Broggi, J.; Terme, T.; Vanelle, P. Organic Electron Donors as Powerful Single-Electron Reducing Agents in Organic Synthesis. *Angew. Chem., Int. Ed.* **2014**, *53* (2), 384–413.
- (4) Ebersson, L. Electron-Transfer Reactions in Organic Chemistry. In *Advances in Physical Organic Chemistry*; Gold, V., Bethell, D., Eds.; Academic Press, 1982; Vol. 18, pp 79–185.
- (5) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113* (7), 5322–5363.
- (6) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. Photoredox Catalysis in Organic Chemistry. *J. Org. Chem.* **2016**, *81* (16), 6898–6926.
- (7) Romero, N. A.; Nicewicz, D. A. Organic Photoredox Catalysis. *Chem. Rev.* **2016**, *116* (17), 10075–10166.
- (8) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* **2016**, *116* (17), 10035–10074.
- (9) Narayanam, J. M. R.; Stephenson, C. R. J. Visible Light Photoredox Catalysis: Applications in Organic Synthesis. *Chem. Soc. Rev.* **2010**, *40* (1), 102–113.
- (10) Teegardin, K.; Day, J. I.; Chan, J.; Weaver, J. Advances in Photocatalysis: A Microreview of Visible Light Mediated Ruthenium and Iridium Catalyzed Organic Transformations. *Org. Process Res. Dev.* **2016**, *20* (7), 1156–1163.
- (11) Capaldo, L.; Ravelli, D. The Dark Side of Photocatalysis: One Thousand Ways to Close the Cycle. *Eur. J. Org. Chem.* **2020**, *2020* (19), 2783–2806.
- (12) Arias-Rotondo, D. M.; McCusker, J. K. The Photophysics of Photoredox Catalysis: A Roadmap for Catalyst Design. *Chem. Soc. Rev.* **2016**, *45* (21), 5803–5820.
- (13) Roth, H. G.; Romero, N. A.; Nicewicz, D. A. Experimental and Calculated Electrochemical Potentials of Common Organic Molecules for Applications to Single-Electron Redox Chemistry. *Synlett* **2016**, *27* (05), 714–723.
- (14) Glaser, F.; Kerzig, C.; Wenger, O. S. Multi-Photon Excitation in Photoredox Catalysis: Concepts, Applications, Methods. *Angew. Chem., Int. Ed.* **2020**, *59* (26), 10266–10284.
- (15) Liu, J.; Lu, L.; Wood, D.; Lin, S. New Redox Strategies in Organic Synthesis by Means of Electrochemistry and Photochemistry. *ACS Cent. Sci.* **2020**, *6* (8), 1317–1340.
- (16) Meerwein, H.; Büchner, E.; Emster, K. van. Über Die Einwirkung Aromatischer Diazoverbindungen Auf α,β -Ungesättigte Carbonylverbindungen. *J. Prakt. Chem.* **1939**, *152* (7–10), 237–266.
- (17) Sainsbury, M. Modern Methods of Aryl-Aryl Bond Formation. *Tetrahedron* **1980**, *36* (23), 3327–3359.
- (18) Hanson, P.; Hammond, R. C.; Goodacre, P. R.; Purcell, J.; Timms, A. W. Sandmeyer Reactions. Part 2. Estimation of

- Absolute Rate Constants for Some Hydrogen-Transfer Reactions and for the Transfer of Water Ligands on CuII to Aryl Radicals by Use of a Pschorr Radical Clock. *J. Chem. Soc., Perkin Trans. 2* **1994**, No. 4, 691–696.
- (19) Ghosh, I.; Marzo, L.; Das, A.; Shaikh, R.; König, B. Visible Light Mediated Photoredox Catalytic Arylation Reactions. *Acc. Chem. Res.* **2016**, *49* (8), 1566–1577.
- (20) Wang, C.-S.; Dixneuf, P. H.; Soulé, J.-F. Photoredox Catalysis for Building C–C Bonds from C(Sp²)–H Bonds. *Chem. Rev.* **2018**, *118* (16), 7532–7585.
- (21) Discekici, E. H.; Treat, N. J.; Poelma, S. O.; Mattson, K. M.; Hudson, Z. M.; Luo, Y.; Hawker, C. J.; Alaniz, J. R. de. A Highly Reducing Metal-Free Photoredox Catalyst: Design and Application in Radical Dehalogenations. *Chem. Commun.* **2015**, *51* (58), 11705–11708.
- (22) Devery, J. J.; Nguyen, J. D.; Dai, C.; Stephenson, C. R. J. Light-Mediated Reductive Debromination of Unactivated Alkyl and Aryl Bromides. *ACS Catal.* **2016**, *6* (9), 5962–5967.
- (23) Boyington, A. J.; Riu, M.-L. Y.; Jui, N. T. Anti-Markovnikov Hydroarylation of Unactivated Olefins via Pyridyl Radical Intermediates. *J. Am. Chem. Soc.* **2017**, *139* (19), 6582–6585.
- (24) Cheng, Y.; Gu, X.; Li, P. Visible-Light Photoredox in Homolytic Aromatic Substitution: Direct Arylation of Arenes with Aryl Halides. *Org. Lett.* **2013**, *15* (11), 2664–2667.
- (25) Shirakawa, E.; Hayashi, T. Transition-Metal-Free Coupling Reactions of Aryl Halides. *Chem. Lett.* **2012**, *41* (2), 130–134. <https://doi.org/10.1246/cl.2012.130>.
- (26) Galli, C. Radical Reactions of Arenediazonium Ions: An Easy Entry into the Chemistry of the Aryl Radical. *Chem. Rev.* **1988**, *88* (5), 765–792.
- (27) Kvasovs, N.; Gevorgyan, V. Contemporary Methods for Generation of Aryl Radicals. *Chem. Soc. Rev.* **2021**, *50* (4), 2244–2259.
- (28) Heinrich, M. R. Intermolecular Olefin Functionalisation Involving Aryl Radicals Generated from Arenediazonium Salts. *Chem. Eur. J.* **2009**, *15* (4), 820–833.
- (29) Raviola, C.; Protti, S. Leaving Groups in Metal-Free Arylations: Make Your Choice! *Eur. J. Org. Chem.* **2020**, *2020* (33), 5292–5304.
- (30) Weix and co-workers conducted an analysis of the commercially available aryl halides and found that aryl chlorides comprise more than 2/3 of the commercial substrates, see: Huang, L.; Ackerman, L. K. G.; Kang, K.; Parsons, A. M.; Weix, D. J. LiCl-Accelerated Multimetallic Cross-Coupling of Aryl Chlorides with Aryl Triflates. *J. Am. Chem. Soc.* **2019**, *141* (28), 10978–10983.
- (31) Electron-deficient aryl chlorides are viable substrates for the most reducing visible light photocatalysts. For examples, see refs. 32–35
- (32) Glaser, F.; Larsen, C. B.; Kerzig, C.; Wenger, O. S. Aryl Dechlorination and Defluorination with an Organic Super-Photoreductant. *Photochem. Photobiol. Sci.* **2020**, *19* (8), 1035–1041.
- (33) Shon, J.-H.; Kim, D.; Rathnayake, M. D.; Sittel, S.; Weaver, J.; Teets, T. S. Photoredox Catalysis on Unactivated Substrates with Strongly Reducing Iridium Photosensitizers. *Chem. Sci.* **2021**, *12* (11), 4069–4078.
- (34) Schmalzbauer, M.; Ghosh, I.; König, B. Utilising Excited State Organic Anions for Photoredox Catalysis: Activation of (Hetero)Aryl Chlorides by Visible Light-Absorbing 9-Anthrolate Anions. *Faraday Discuss.* **2019**, *215* (0), 364–378.
- (35) Ghosh, I.; Shaikh, R. S.; König, B. Sensitization-Initiated Electron Transfer for Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2017**, *56* (29), 8544–8549.
- (36) Blanksby, S. J.; Ellison, G. B. Bond Dissociation Energies of Organic Molecules. *Acc. Chem. Res.* **2003**, *36* (4), 255–263.
- (37) Costentin, C.; Robert, M.; Savéant, J.-M. Fragmentation of Aryl Halide π Anion Radicals. Bending of the Cleaving Bond and Activation vs Driving Force Relationships. *J. Am. Chem. Soc.* **2004**, *126* (49), 16051–16057.
- (38) Ghosh, I.; Ghosh, T.; Bardagi, J. I.; König, B. Reduction of Aryl Halides by Consecutive Visible Light-Induced Electron Transfer Processes. *Science* **2014**, *346* (6210), 725–728.
- (39) Neumeier, M.; Sampedro, D.; Májek, M.; de la Peña O’Shea, V. A.; Jacobi von Wangelin, A.; Pérez-Ruiz, R. Dichromatic Photocatalytic Substitutions of Aryl Halides with a Small Organic Dye. *Chem. Eur. J.* **2018**, *24* (1), 105–108.
- (40) MacKenzie, I. A.; Wang, L.; Onuska, N. P. R.; Williams, O. F.; Begam, K.; Moran, A. M.; Dunietz, B. D.; Nicewicz, D. A. Discovery and Characterization of an Acridine Radical Photoreductant. *Nature* **2020**, *580* (7801), 76–80.
- (41) Cole, J. P.; Chen, D.-F.; Kudisch, M.; Pearson, R. M.; Lim, C.-H.; Miyake, G. M. Organocatalyzed Birch Reduction Driven by Visible Light. *J. Am. Chem. Soc.* **2020**, *142* (31), 13573–13581.
- (42) Wang, S.; Wang, H.; Koenig, B. Photo-Induced Thiolate Catalytic Activation of Inert Caryl-Hetero Bonds for Radical Borylation. **2020**. <https://doi.org/10.26434/chemrxiv.13298837.v1>.
- (43) Jin, S.; Dang, Hang. T.; Haug, G. C.; He, R.; Nguyen, V. D.; Nguyen, V. T.; Arman, H. D.; Schanze, K. S.; Larionov, O. V. Visible Light-Induced Borylation of C–O, C–N, and C–X Bonds. *J. Am. Chem. Soc.* **2020**, *142* (3), 1603–1613.
- (44) Zhang, L.; Jiao, L. Visible-Light-Induced Organocatalytic Borylation of Aryl Chlorides. *J. Am. Chem. Soc.* **2019**, *141* (23), 9124–9128.
- (45) Cybularczyk-Cecotka, M.; Szczepanik, J.; Giedyk, M. Photocatalytic Strategies for the Activation of Organic Chlorides. *Nat Catal* **2020**, *3* (11), 872–886.
- (46) Barham, J. P.; König, B. Synthetic Photoelectrochemistry. *Angew. Chem., Int. Ed.* **2020**, *59* (29), 11732–11747.
- (47) Novaes, L. F. T.; Liu, J.; Shen, Y.; Lu, L.; Meinhardt, J. M.; Lin, S. Electrocatalysis as an Enabling Technology for Organic Synthesis. *Chem. Soc. Rev.* **2021**.
- (48) Capaldo, L.; Quadri, L. L.; Ravelli, D. Merging Photocatalysis with Electrochemistry: The Dawn of a New Alliance in Organic Synthesis. *Angew. Chem., Int. Ed.* **2019**, *58* (49), 17508–17510.
- (49) Yu, Y.; Guo, P.; Zhong, J.-S.; Yuan, Y.; Ye, K.-Y. Merging Photochemistry with Electrochemistry in Organic Synthesis. *Org. Chem. Front.* **2019**, *7* (1), 131–135.
- (50) Cowper, N. G. W.; Chernowsky, C. P.; Williams, O. P.; Wickens, Z. K. Potent Reductants via Electron-Primed Photoredox Catalysis: Unlocking Aryl Chlorides for Radical Coupling. *J. Am. Chem. Soc.* **2020**.
- (51) Kim, H.; Kim, H.; Lambert, T. H.; Lin, S. Reductive Electrophotocatalysis: Merging Electricity and Light To Achieve Extreme Reduction Potentials. *J. Am. Chem. Soc.* **2020**, *142* (5), 2087–2092.
- (52) For example, HTE remains particularly difficult for electrochemical processes but is well-established for photochemical systems. Recent work from Lin has advanced an attractive HTE plate for electrophotocatalysis in undivided cells. However, divided cells are required for all reductive electrophotocatalysis reported to-date, which would not be compatible with this technology, see: ref 53

- (53) Rein, J.; Annand, J. R.; Wismer, M. K.; Fu, J.; Siu, J. C.; Klapars, A.; Strotman, N. A.; Kalyani, D.; Lehnher, D.; Lin, S. Unlocking the Potential of High-Throughput Experimentation for Electrochemistry with a Standardized Microscale Reactor. *2021*. <https://doi.org/10.26434/chemrxiv.14173538.v2>.
- (54) ArCl with EWG are in bounds and typically work, albeit in diminished yield. For a selected examples, see: 55 and 56
- (55) Li, H.; Liu, Y.; Chiba, S. Anti-Markovnikov Hydroarylation of Alkenes via Polysulfide Anion Photocatalysis. *Chem. Commun.* **2021**. DOI: 10.1039/d1cc02185k
- (56) Boyington, A. J.; Seath, C. P.; Zearfoss, A. M.; Xu, Z.; Jui, N. T. Catalytic Strategy for Regioselective Arylethylamine Synthesis. *J. Am. Chem. Soc.* **2019**, *141* (9), 4147–4153.
- (57) Andrieux, C. P.; Pinson, J. The Standard Redox Potential of the Phenyl Radical/Anion Couple. *J. Am. Chem. Soc.* **2003**, *125* (48), 14801–14806.
- (58) Chernowsky, C.; Chmiel, A.; Wickens, Z. Photocatalytic Activity of Diverse Organic Radical Anions: Catalyst Discovery Enables Cleavage of Strong C(Sp²)-N and C(Sp²)-O Bonds. *2021*. <https://doi.org/10.26434/chemrxiv.14710398.v1>.
- (59) Targos, K.; Williams, O. P.; Wickens, Z. K. Unveiling Potent Photooxidation Behavior of Catalytic Photoreductants. *J. Am. Chem. Soc.* **2021**, *143* (11), 4125–4132.
- (60) Connell, T. U.; Fraser, C. L.; Czyn, M. L.; Smith, Z. M.; Hayne, D. J.; Doeven, E. H.; Agugiaro, J.; Wilson, D. J. D.; Adcock, J. L.; Scully, A. D.; Gómez, D. E.; Barnett, N. W.; Polyzos, A.; Francis, P. S. The Tandem Photoredox Catalysis Mechanism of [Ir(Ppy)₂(Dtb-Bpy)]⁺ Enabling Access to Energy Demanding Organic Substrates. *J. Am. Chem. Soc.* **2019**, *141*, 17646–17658.
- (61) Chernowsky, C.; Chmiel, A.; Wickens, Z. Photocatalytic Activity of Diverse Organic Radical Anions: Catalyst Discovery Enables Cleavage of Strong C(Sp²)-N and C(Sp²)-O Bonds. *2021*. <https://doi.org/10.26434/chemrxiv.14710398.v1>.
- (62) Hu, J.; Wang, J.; Nguyen, T. H.; Zheng, N. The Chemistry of Amine Radical Cations Produced by Visible Light Photoredox Catalysis. *Beilstein J. Org. Chem.* **2013**, *9* (1), 1977–2001.
- (63) Speckmeier, E.; Fischer, T. G.; Zeitler, K. A Toolbox Approach To Construct Broadly Applicable Metal-Free Catalysts for Photoredox Chemistry: Deliberate Tuning of Redox Potentials and Importance of Halogens in Donor-Acceptor Cyanoarenes. *J. Am. Chem. Soc.* **2018**, *140* (45), 15353–15365.
- (64) Recent reports from Leonori and coworkers have illustrated that α -amino radicals formed in situ from alkylamine oxidation promote aryl radical reactions from aryl iodides and bromides via halogen atom transfer (XAT). However, the stronger and less polarizable C(sp²)-Cl bonds are expected to be less amenable to such a mechanism, see: refs. 65 and 66.
- (65) Constantin, T.; Zanini, M.; Regni, A.; Sheikh, N. S.; Juliá, F.; Leonori, D. Aminoalkyl Radicals as Halogen-Atom Transfer Agents for Activation of Alkyl and Aryl Halides. *Science* **2020**, *367* (6481), 1021–1026.
- (66) Constantin, T.; Juliá, F.; Sheikh, N. S.; Leonori, D. A Case of Chain Propagation: α -Aminoalkyl Radicals as Initiators for Aryl Radical Chemistry. *Chem. Sci.* **2020**, *11* (47), 12822–12828.
- (67) Azzena, U.; Denurra, T.; Melloni, G.; Fenude, E.; Rassu, G. Electron-Transfer-Induced Reductive Demethoxylation of Anisole: Evidence for Cleavage of a Radical Anion. *J. Org. Chem.* **1992**, *57* (5), 1444–1448.
- (68) Xu, H.; Yu, B.; Zhang, H.; Zhao, Y.; Yang, Z.; Xu, J.; Han, B.; Liu, Z. Reductive Cleavage of Inert Aryl C-O Bonds to Produce Arenes. *Chem. Commun.* **2015**, *51* (61), 12212–12215.
- (69) The oxidation potential of formate, +1.25 V vs SCE, is within the bound of the excited state reduction potential of each catalyst. For the potential of formate, see supporting information. The Oxidation Potential of Formate, +1.25 V vs SCE, Is within the Bound of the Excited State Reduction Potential of Each Catalyst. For the CV of Formate, See Supporting Information.
- (70) Gu, X.; Lu, S.; Fu, X.; Qiu, Z.; Sui, Q.; Guo, X. Carbon Dioxide Radical Anion-Based UV/S₂O₈²⁻/HCOOH Reductive Process for Carbon Tetrachloride Degradation in Aqueous Solution. *Sep. Purif. Technol.* **2017**, *172*, 211–216.
- (71) Jacobsen, E.; Roberts, J. L.; Sawyer, D. T. Electrochemical Oxidation of Formate in Dimethylsulfoxide at Gold and Platinum Electrodes. *J. Electroanal. Chem. Interf. Electrochem.* **1968**, *16* (3), 351–360.
- (72) Grills, D. C.; Lyman, S. V. Radiolytic Formation of the Carbon Dioxide Radical Anion in Acetonitrile Revealed by Transient IR Spectroscopy. *Phys. Chem. Chem. Phys.* **2018**, *20* (15), 10011–10017.
- (73) Rosso, J. A.; Bertolotti, S. G.; Braun, A. M.; Mártire, D. O.; Gonzalez, M. C. Reactions of Carbon Dioxide Radical Anion with Substituted Benzenes. *J. Phys. Org. Chem.* **2001**, *14* (5), 300–309.
- (74) Koppenol, W. H.; Rush, J. D. Reduction Potential of the Carbon Dioxide/Carbon Dioxide Radical Anion: A Comparison with Other C1 Radicals. *J. Phys. Chem.* **1987**, *91* (16), 4429–4430.
- (75) Sevrin, M. J.; Furst, L.; Nguyen, J. D.; Collins, J. L.; Stephenson, C. R. J. Lithium Bis-Catechol Borate as an Effective Reductive Quencher in Photoredox Catalysis. *Tetrahedron* **2018**, *74* (26), 3246–3252.
- (76) Beatty, J. W.; Stephenson, C. R. J. Amine Functionalization via Oxidative Photoredox Catalysis: Methodology Development and Complex Molecule Synthesis. *Acc. Chem. Res.* **2015**, *48* (5), 1474–1484.
- (77) Singh, P. P.; Srivastava, V. Recent Advances in Using 4DPAIPN in Photocatalytic Transformations. *Org. Biomol. Chem.* **2021**, *19* (2), 313–321.
- (78) Németh, G.; Greff, Z.; Sipos, A.; Varga, Z.; Székely, R.; Sebestyén, M.; Jászay, Z.; Béni, S.; Nemes, Z.; Pirat, J.-L.; Volle, J.-N.; Virieux, D.; Gyuris, Á.; Kelemenics, K.; Áy, É.; Minarovits, J.; Szathmary, S.; Kéri, G.; Örfi, L. Synthesis and Evaluation of Phosphorus Containing, Specific CDK9/CycT1 Inhibitors. *J. Med. Chem.* **2014**, *57* (10), 3939–3965.
- (79) Fyfe, J. W. B.; Watson, A. J. B. Recent Developments in Organoboron Chemistry: Old Dogs, New Tricks. *Chem* **2017**, *3* (1), 31–55.
- (80) We found for most of the reactions studied herein that the photocatalyst bleached over the course of the reaction and that adding photocatalyst in two batches provided improved yields, see SI for details. For an example of isophthalonitrile catalysts being attacked by radical intermediates, see: Grotjahn, S.; König, B. *Org. Lett.* **2021**, *23*, 3146–3150
- (81) For an Example of Unactivated Alkene Hydroarylation, See: Ref. 55.
- (82) While both alkylamine and formate reductants can act as hydrogen atom sources, we noted that sodium formate exhibits limited solubility under the reaction conditions. We

- suspect that the limited concentration of the reductant diminishes premature HAT to the aryl radical relative to homogeneous alkylamine reductants.
- (83) Ito, O. Flash Photolysis Study on Reversible Addition Reactions of Thiyl Radicals. *Res. Chem. Intermed.* **1995**, *21* (1), 69–93.
- (84) For a proposed mechanism for CO₂^{•-} generation from PhSSPh, see supporting information. PhSSPh was used as the thiyl radical source to initiate the slightly endothermic HAT (BDE PhSH = 79 kcal/mol, BDE HCO₂⁻ = 86 kcal/mol, see: ref 84).
- (85) Dénès, F.; Pichowicz, M.; Povie, G.; Renaud, P. Thiyl Radicals in Organic Synthesis. *Chem. Rev.* **2014**, *114* (5), 2587–2693.
- (86) McDaniel, K. A.; Blood, A. R.; Smith, G. C.; Jui, N. T. Dearomatization of Unactivated Arenes via Catalytic Hydroalkylation. *ACS Catal.* **2021**, *11* (9), 4968–4972.
-