

Branch-Selective Addition of Unactivated Olefins into Imines and Aldehydes

Jeishla L. M. Matos, Suhelen Vásquez-Céspedes, Jieyu Gu, Takuya Oguma, Ryan A. Shenvi*

Department of Chemistry, The Scripps Research Institute, 10550 N Torrey Pines Rd, La Jolla, CA 92037, United States

Supporting Information Placeholder

Radical hydrofunctionalization occurs with ease using metal-hydride atom transfer (MHAT) catalysis to couple alkenes and competent radicalophilic electrophiles. Traditional two-electron electrophiles have remained unreactive. Herein we report the addition of electronically-unbiased olefins into imines and aldehydes. Iron-catalysis allows addition of alkyl-substituted olefins into imines through the intermediacy of free-radicals, whereas a combination of catalytic $\text{Co}(\text{Sal}^{\text{t-Bu},\text{t-Bu}})$ and chromium salts enable a branch-selective coupling of olefins and aldehydes through the formation of a putative alkyl chromium intermediate.

Branch-selective reactions of alkyl-substituted olefins via carbocationic¹ or radical intermediates² benefits from an abundance of methods, but the analogous transformation into branched-carbanion equivalents remains underdeveloped (Figure 1). A common way to transform an olefin into a carbanion equivalent is via hydrometallation of a double bond. However, such branch-selective hydrometallation of alkenes is generally limited to styrenes, allenes, and dienes³—all electronically biased systems that stabilize a developing carbon-metal bond.⁴ In the absence of electronic bias, steric constraints dominate: canonical metal hydrides favor linear selectivity and linear hydrometallation is predominately observed.⁵ To obtain branch-selectivity with electronically-unbiased alkenes, we have investigated M—H hydrogen atom transfer (MHAT) catalysis and subsequent capture of the nascent intermediates by a second metal complex.^{6,7,8} For example, we recently established that nickel complexes intercept $\text{Co}(\text{Sal}^{\text{t-Bu},\text{t-Bu}})$ -catalyzed MHAT cycles in a direct organocobalt to organonickel transmetallation.^{7,7} Similar alkyl transmetallations have been reported in non-catalytic systems between alkyl— $\text{Co}(\text{dmgBF}_2)_2\text{Py}_2$ and inorganic nickel,⁹ and proposed for bioorganometallic¹⁰ and catalytic processes.¹¹ This alkyl transfer does not appear limited to nickel: vitamin B12-mimetics (such as $\text{Co}(\text{salen})$ derivatives) can undergo alkyl transfer to palladium,¹² rhodium,¹³ other cobalt,¹⁴ platinum,¹⁵ gold,¹⁶ chromium,¹⁷ and zinc¹⁸ salts and organometallics species. Yet despite the apparent generality of this transformation, there is a paucity of preparative cross-coupling methods which leverage this reactivity.




Figure 1: Transformation of olefins into carbanion equivalents by a radical/ polar crossover.

The capacity to form cobalt organometallics via MHAT followed by cage-collapse^{7,19,20,21} prompted us to explore transmetalation partners that might lead to otherwise inaccessible branched products. Here we show that olefins can be added to imines and aldehydes to form sp^3 — sp^3 bonds. The former reaction with an activated electrophile occurs under standard MHAT catalysis, whereas the latter reaction requires interception of MHAT intermediates with chromium salts (Figure 1e). This transformation expands the current scope of olefins as carbanion surrogates²² which has heretofore required the use of electronically-activated olefins, such as styrene, allenyl, or dienes. Alkyl-substituted olefins, in contrast, react with carbonyls at the least-substituted position through a Prins mechanism,²³ or undergo iron-catalyzed hydromagnesiation reactions to form linear nucleophiles.^{24,25,26}

We initially investigated the Markovnikov addition of alkenes into carbonyl derivatives by utilizing the intermediacy of the free radicals and noticed that productive reactions





Figure 2. Alkyl radicals generated by MHAT add to chiral sulfinimines. d.r. of two major diastereomers reported. ^astereochemistry at the α -carbon is (*S*). ^bstereochemistry in the α -carbon is (*R*). ^cMn(dpm)₃ instead of Fe(acac)₃ was used. ^dcontains 15% unrearranged pinene product and its diastereomer.

were only obtained with standard radicalophiles, such as radical-stabilizing imines. Glyoxyimines are preceded as radical acceptors,²⁷ and chiral sulfinyl auxiliaries can be used to impart stereocontrol. Although the competitive reduction of these electrophiles by the metal hydrides or the stoichiometric silane was observed, this could be minimized by using a slight excess of the olefin and Fe⁺³ salts as the catalyst.²⁸ Several feedstock alkenes served as competent nucleophilic components and delivered unnatural amino acids derivatives with good to excellent diastereoselectivities. (Figure 2). The early transition state of radical reactions allows facile formation of sterically hindered unnatural α -amino acids bearing β -quaternary carbons, and reactive groups like free-hydroxyls (**13**) or two-electron electrophiles such as esters, epoxides, or aldehydes (**8**, **15**, and **17**) are tolerated. Complex feedstock terpenes can engage the sulfinylimines to deliver adducts **12**, **14**, and **15**, and even glycans deliver amino esters with good diastereocontrol (**16**). Comparison of the optical rotation obtained from our reaction to that of *t*-butyl glycine derivatives shows that sulfinimines with the (*S*)-configuration affords the (*S*)-amine whereas the (*R*)-sulfinimine affords the (*R*)-amine.²⁸ Better diastereoselectivity is obtained with the more hindered mesitylene or tri-isopropyl arene-derived sulfinamide. Given the ease with which these compounds are made (i.e. no prefunctionalization is necessary for radical formation) we anticipate that this method will find application in the synthesis of unnatural amino acids.²⁹

Addition of the free radical to aldehydes, however, proved challenging (see Table 1), as may be expected due to the higher instability of an *O*-centered radical relative to a *C*-centered radical, which is reflected by the more facile C–C bond scission than C–C bond formation.³⁰ Strategies to

drive this energetically unfavorable addition include sequestering the unstable *O*-centered radical as an alkoxide (which cannot undergo homolytic β -scission) in an intramolecular setting³¹ or accessing excited-states via photochemistry.³² However, neither strategy may be used for intermolecular

Table 1. Conversion of *C*-centered radicals to 2-electron nucleophiles.



Entry	Deviations from above	Yield (%) ^a
1	Fe(dpm) ₃ , Fe(acac) ₃ , Co(acac) ₂ or Mn(dpm) ₃ instead of Co(Sal ^{<i>t</i>-Bu,<i>t</i>-Bu})	< 10%
2	CrCl ₂ instead of CrCl ₃	trace
3	with Zn ⁰ or Mn ⁰	11%
4	Co(salen)Cl and no [O]	45% ^c
5	no [O]	–
6	0.2 equiv. of CrCl ₃ instead of 1 equiv.	22%
7	0.2 equiv. of CrCl ₃ and TMSCl (1 equiv.)	–
8	in DMF instead of THF/CH ₃ CN	–
9	without CH ₃ CN	35%
10	under air	46%
11	with 1 equiv. of H ₂ O	–
12	No Co(Sal ^{<i>t</i>-Bu,<i>t</i>-Bu})	–
13	No CrCl ₃	– ^d

^ayield determined by GC/FID using a calibrated internal standard; ^b isolated yield with 20 mol % of Co(salen)Cl and CrCl₃(THF)₃; ^c1 equiv. of NaBF₄ added; ^dsome isomerization and hydrogenation was observed; d.r. 1:1 in all cases.




Figure 3. Conversion of alkenes into carbanion surrogates with branched-selectivity. d.r. at the formed bond is close to 1:1 unless otherwise noted. See Supporting Information for the d.r. of the isolated compounds. ^a20 mol% of [Co] and 20 mol% of [F⁺] were used.

addition with alkyl-substituted olefins.³³ In light of the precedence for alkyl-cobalt complexes to transmetallate other metal species and the apparent facility with which organocobalt species can form from olefins,^{7,19} we wondered if a two-electron nucleophile equivalent could arise from unactivated olefins via sequential one-electron reductions via interception of organocobalt with chromium species.

We were drawn to chromium chemistry for several reasons: 1) organochromium reagents are known to add into carbonyls in a 1,2-fashion² 2) Cr⁺² salts are proposed to intercept alkyl-radicals to form organochromium species with bimolecular rate constants on the order of 10⁷ M⁻¹s⁻¹,³⁴ 3) alkyl-cobalamines and -cobaloximes can also be intercepted by Cr⁺²,^{17,35,36} and 4) chromium salts are inexpensive and of low toxicity in the +2 and +3 oxidation states.³⁷ Furthermore, the weak Brønsted acidity of organochromium complexes

allows for a high functional group tolerance and for their use in late-stage functionalization for complex molecule synthesis.³⁸


Initially, attempts to merge MHAT catalysis and chromium chemistry met with poor results. β -diketonate complexes of Co and Mn were not productive, although iron salts afforded the product in low yield (Table 1, Entry 1).³⁹ We discovered, however, that use of Co(Sal^{*t*-Bu,*t*-Bu}) and equimolar amounts of 1-fluoro-2,4,6-trimethylpyridinium tetrafluoroborate in the presence of phenylsilane and CrCl₃ could couple the terminal olefin in **19** to 3-trifluoromethyl benzaldehyde in good yields. Given that Cr⁺² is typically the active species in the addition of alkyl halides into carbonyls, we initially explored the reaction using CrCl₂ or CrCl₃ alongside an external metal reductant only to discover that these conditions lead to less product formed than the amount of [Co]

pre-catalyst added (Entries 2 and 3). One explanation is that the external reductants impede the Co-cycle by unproductive reduction of Co^{+3} intermediates.⁴⁰ Our optimized conditions appear to circumvent this problem by reductive formation of Cr^{+2} *in situ* (see below). Although it is possible to perform this reaction with the pre-oxidized $\text{Co}(\text{Sal}^{t\text{-Bu}, t\text{-Bu}})\text{Cl}$, use of Co^{+2} and an external oxidant generally afforded higher yields (Entry 4), thereby allowing use of the more convenient +3 and +2 oxidation states of Cr and Co, respectively. Control experiments indicate that both metals are necessary for product formation (Entries 12-13).⁴¹


Evaluation of the scope (Figure 3) revealed that both aromatic and alkyl aldehydes are competent electrophiles, and a wide range of electronic variation is tolerated. In general, electron withdrawn substrates afford higher yields than electron-rich electrophiles, yet even vanillin-derived aldehydes such as **33** and **34** react in high yield. Various heteroaromatic aldehydes may be employed (**37** – **39**), as well as terpene-derived substrates (**41**). Esters (**45**), tosylates (**44**), and chlorides (**46**) are orthogonal electrophiles, but competitive reduction of bromides, and iodides was observed. A switch in solvent from THF to DME allows 1,2-disubstituted olefins to be engaged (**54** – **57**), although trisubstituted olefins are not yet competent. Modest diastereocontrol is imparted by a chiral directing group (**49**),⁴² and sterically bulky substrates (**35**, **43**, **50** and **52**).⁴³

Figure 4. Delayed addition and stoichiometric reactions suggest transmetalation of alkyl-Co^{+3} to alkyl-Cr^{+3} .

a. Effect of delayed addition of CrCl_3^a



b. Oxidation state of active Cr reagent



^areactions ran with 20 mol% of [Co] and 20 mol% of [O].

Although we currently do not have a complete mechanistic understanding, several observations are worth noting. First, the yield of the product formed does not vary as a function of delayed Cr^{+3} addition, which is consistent with intermediate formation of an organocobalt species that is engaged by the Cr, and inconsistent with an alternative hypothesis of Cr capture by a freely-diffusing C-centered radical;⁴⁴ analogy can be drawn to our previously reported Ni/Co hydroarylation⁷ and the mechanistic studies of Espenson and coworkers.³⁵ Stoichiometric experiments support a transmetalation and suggest reaction with Cr^{+2} rather than the Cr^{+3} species. In these experiments, a *sec*-alkyl cobalt was formed *in situ* by displacement of 2-bromopropane by $\text{Co}(\text{Sal}^{t\text{-Bu}, t\text{-Bu}})$; addition of CrCl_3 and aldehyde **20** yields no product, whereas CrCl_2 produces 11% of adduct **58**.⁴⁵ We suspect reduction of Cr^{+3} to Cr^{+2} occurs via the stoichiometric silane reductant necessary for the MHAT catalytic cycle.^{46,47} By analogy to the proposal of Espenson and coworkers in similar systems,³⁵ a possible mechanism for the alkyl transfer could involve electron transfer from a Cr^{+2} to an alkyl-Co^{+3} intermediate to form an unstable alkyl-Co^{+2} species which is known to homolyze to afford an alkyl radical that could escape the solvent cage and capture a second Cr^{+2} species, a kinetically facile process ($k = 10^7 \text{ M}^{-1}\text{s}^{-1}$).^{34,48,49}

In summary, we have discovered divergent reactivity available to alkenes that enables branch-selective (Markovnikov) addition to radicalophilic and non-radicalophilic electrophiles. First, carbon-centered radicals generated by MHAT are competent to add to chiral sulfinimines, which stabilize the incipient *N*-centered radical, and impart stereocontrol. The products of these reactions are valuable, unnatural amino acid derivatives. Second and complementarily, although these same radicals do not productively add into aldehydes, the addition of Cr^{+3} salts allows coupling to occur. This latter method circumvents the poor reactivity of free radicals towards carbonyl intermediates while maintaining the Markovnikov reactivity and chemoselectivity of MHAT. Overall, this work enables cross-coupling of abundant chemical feedstocks (aldehydes and olefins) without the need for pre-functionalization. Mechanistic experiments and analogy to the literature is consistent with alkyl-Co^{+3} transmetalation to alkyl-Cr^{+3} , mediated by Cr^{+2} . This second example^{7,7} of catalytic MHAT organocobalt transmetalation calls attention to the potentially general use of these alkyl-cobalt complexes as catalytically-generated organometallic species capable of transferring their alkyl ligands to various other transition metals (including Ni and Cr) for previously inaccessible branch-selective bond-forming processes from olefins. This reactivity complements catalytically-generated organocuprate species which can also engage in hydrometallation/ transmetalation, but do so with linear selectivity.⁵⁰

ASSOCIATED CONTENT

Supporting Information.

The Supporting Information is available free of charge on

the ACS Publications website.

Detailed experimental procedures, compound characterization and spectral data.

AUTHOR INFORMATION

Corresponding Author

rshenvi@scripps.edu

Funding Sources

Generous support was provided by the National Institutes of Health (R35 GM122606) and the NSF (GRFP to J. L. M. M.). We thank Dr. L. Pasternack and Dr. D.-H. Huang for NMR assistance, Dr. J. Chen and Brittany Sánchez for HRMS measurements, and Dr. R. Gianatassio for providing the starting material for **9**.

REFERENCES

1. Schevenels, F. T.; Shen, M.; Snyder, S. A. Isolable and Readily Handled Halophosphonium Pre-reagents for Hydro- and Deuteriohalogenation. *J. Am. Chem. Soc.* **2017**, *139*, 6329.
2. Crossley, S. W. M.; Martinez, R. M.; Obradors, C.; Shenvi, R. A. Mn-, Fe-, and Co-Catalyzed Radical Hydrofunctionalizations of Olefins. *Chem. Rev.* **2016**, *116*, 8912.
3. Or π -systems proximal to a directing group. See: Murphy, S. K., Coulter, M. M., & Dong, V. M. β -hydroxy ketones prepared by regioselective hydroacylation. *Chem. Sci.*, **2012**, *3*, 355.
4. Coombs, J. R.; Morken, J. P. Catalytic Enantioselective Functionalization of Unactivated Terminal Alkenes. *Angew. Chem. Int. Ed.* **2016**, *55*, 2636.
5. Crabtree, R. H.; *The Organometallic Chemistry of the Transition Metals*, Sixth Edition. John Wiley & Sons: New Jersey, 2014.
6. Green, S. A.; Matos, J. L. M.; Yagi, A.; Shenvi, R. A. Branch-Selective Hydroarylation: Iodoarene-Olefin Cross-Coupling. *J. Am. Chem. Soc.* **2016**, *138*, 12779.
7. Shevick, S. L.; Obradors, C.; Shenvi, R. A. Mechanistic Interrogation of Co/Ni-Dual Catalyzed Hydroarylation. *J. Am. Chem. Soc.* **2018**, *140*, 12056.
8. Green, S. A.; Vásquez-Céspedes, S.; Shenvi, R. A. Iron-Nickel Dual-Catalysis: A New Engine for Olefin Functionalization. *J. Am. Chem. Soc.* **2018**, *140*, 11317.
9. (a) Ram, M. S.; Yap, G. P. A.; Liable-Sands, L.; Rheingold, A. L.; Marchaj, A.; Norton, J. R. Kinetics and Mechanism of Alkyl Transfer from Organocobalt(III) to Nickel(I): Implications for the Synthesis of Acetyl Coenzyme A by CO Dehydrogenase. *J. Am. Chem. Soc.* **1997**, *119*, 1648. (b) Ram, M. S.; Riordan, C. G. Methyl Transfer From a Cobalt Complex to Ni(tmc)⁺ Yielding Ni(tmc)Me⁺: A model for Methylcobalamin Alkylation of CO Dehydrogenase. *J. Am. Chem. Soc.* **1995**, *117*, 2365.
10. Can, M.; Armstrong, F. A.; Ragsdale, S. W. Structure, Function, and Mechanism of the Nickel Metalloenzymes, CO Dehydrogenase, and Acetyl-CoA Synthase. *Chem. Rev.*, **2014**, *114*, 4149-4174.
11. Komeyama, K.; Yamahata, Y.; Osaka, I. Nickel and Nucleophilic Cobalt-Catalyzed Trideuteriomethylation of Aryl Halides Using Trideuteriomethyl p-Toluenesulfonate. *Org. Lett.* **2018**, *20*, 4375.
12. Scovell, W. M. Kinetics and Mechanism of Methyl Transfer from Methylcobalamin to Palladium(II). *J. Am. Chem. Soc.* **1974**, *96*, 3451.
13. Dodd, D.; Johnson, M. D. Bimolecular Nucleophilic Displacement as a Mechanism of Alkyl Transfer from Cobalt. *J. Chem. Soc. D*, **1971**, 1371.
14. (a) Stolzenberg, A. M.; Cao, Y. Alkyl Exchange Reactions of Organocobalt Porphyrins. A Bimolecular Homolytic Substitution Reaction. *J. Am. Chem. Soc.* **2001**, *123*, 9078-9090. (b) Dodd, D.; Johnson, M. D.; Lockman, B. L. Kinetics and Mechanism of Apparent Alkyl Transfer from Alkylcobaloximes to Cobaloxime(I), Cobaloxime(II), and Cobaloxime(III) reagents. *J. Am. Chem. Soc.* **1977**, *99*, 3664.
15. Fanchiang, Y. T.; Ridley, W. P.; Wood, J. M. Methylation of Platinum Complexes by Methylcobalamin. *J. Am. Chem. Soc.* **1979**, *101*, 1442.
16. Agnes, G.; Bendle, S.; Hill, H. A. O.; Williams, F. R.; Williams, R. J. P. Methylation by Methyl Vitamin B12. *J. Chem. Soc., Chem. Commun.* **1971**, 850.
17. Espenson, J. H.; Shveima, J. S. Alkyl Transfer from Cobalt to Chromium. *J. Am. Chem. Soc.* **1973**, *95*, 4468.
18. Witman, M. W.; Weber, J. H. Methylation of zinc(II), cadmium(II), and lead(II) by a trans-dimethylcobalt complex. *Inorg. Chem.*, **1976**, *15*, 2375.
19. Crossley, S. W. M.; Barabe, F.; Shenvi, R. A. Simple, Chemoselective, Catalytic Olefin Isomerization. *J. Am. Chem. Soc.* **2014**, *136*, 16788.
20. Eisenberg, D. C.; Norton, J. R. Hydrogen-Atom Transfer Reactions of Transition-Metal Hydrides. *Isr. J. Chem.* **1991**, *31*, 55.
21. Formation of an organocobalt from olefins is also postulated on: (a) Shigehisa, H.; Aoki, T.; Yamaguchi, S.; Shimizu, N.; Hiroya, K. Hydroalkoxylation of Unactivated Olefins with Carbon Radicals and Carbocation Species as Key Intermediates. *J. Am. Chem. Soc.* **2013**, *135*, 10306. (b) Shigehisa, H.; Nishi, E.; Fujisawa, M.; Hiroya, K. Cobalt-Catalyzed Hydrofluorination of Unactivated Olefins: A Radical Approach of Fluorine Transfer. *Org. Lett.*, **2013**, *15*, 5158. (c) Tokuyasu, T.; Kunikawa, S.; Masuyama, A.; Nojima, M.

- Co(III)–Alkyl Complex- and Co(III)–Alkylperoxo Complex-Catalyzed Triethylsilylperoxidation of Alkenes with Molecular Oxygen and Triethylsilane. *Org. Lett.*, **2002**, *4*, 3595. (d) Waser, J.; Gaspar, B.; Nambu, H.; Carreira, E. M. Hydrazines and Azides via the Metal-Catalyzed Hydrohydrazination and Hydroazidation of Olefins. *J. Am. Chem. Soc.*, **2006**, *128*, 11693. (e) Gaspar, B.; Carreira, E. M. Catalytic Hydrochlorination of Unactivated Olefins with para-Toluenesulfonyl Chloride. *Angew. Chem. Int. Ed.* **2008**, *47*, 5758.
22. (a) Nguyen, K. D.; Park, B. Y.; Luong, T.; Sato, H.; Garza, V. J.; Krische, M. J. Metal-catalyzed reductive coupling of olefin-derived nucleophiles: Reinventing carbonyl addition. *Science*, **2016**, *354*, 6310.
 23. Zheng, Y.-L.; Liu, Y.-Y.; Wu, Y.-M.; Wang, Y.-X.; Lin, Y.-T.; Ye, M. Iron-Catalyzed Regioselective Transfer Hydrogenative Couplings of Unactivated Aldehydes with Simple Alkenes. *Angew. Chem. Int. Ed.* **2016**, *55*, 6315.
 24. Shirakawa, E.; Ikeda, D.; Masui, S.; Yoshida, M.; Hayashi, T. Iron-Copper Cooperative Catalysis in the Reactions of Alkyl Grignard Reagents: Exchange Reaction with Alkenes and Carbometalation of Alkynes. *J. Am. Chem. Soc.* **2012**, *134*, 272.
 25. α -olefins have also been added to activated α -hydroxy ketones or diols via cyclometallation (a) Yamaguchi, E.; Mowat, J.; Luong, T.; Krische, M. J. Regio- and Diastereoselective C-C Coupling of α -Olefins and Styrenes to 3-Hydroxy-2-oxindoles by Ru-Catalyzed Hydrohydroxyalkylation. *Angew. Chem. Int. Ed.* **2013**, *52*, 8428. (b) Park, B. Y.; Luong, T.; Sato, H.; Krische, M. J. Osmium(0)-Catalyzed C–C Coupling of Ethylene and α -Olefins with Diols, Ketols, or Hydroxy Esters via Transfer Hydrogenation. *J. Org. Chem.*, **2016**, *81*, 8585.
 26. Branch-selective hydroformylations of alkyl olefins have been reported in a number of cases, but at generally low ratios, with the exception of idiosyncratic substrates. See: (a) Pittaway, R.; Fuentes, J. A.; Clarke, M. L. Diastereoselective and Branched-Aldehyde-Selective Tandem Hydroformylation–Hemiaminal Formation: Synthesis of Functionalized Piperidines and Amino Alcohols. *Org. Lett.* **2017**, *19*, 2845. (b) Kuil, M.; Soltner, T.; van Leeuwen, P. W. N. M.; Reek, J. N. H. High-Precision Catalysts: Regioselective Hydroformylation of Internal Alkenes by Encapsulated Rhodium Complexes. *J. Am. Chem. Soc.* **2006**, *128*, 11344.
 27. (a) Garrido-Castro, A. F.; Choubane, H.; Daaou, M.; Maestro, M. C. Alemán, J. Asymmetric radical alkylation of N-sulfinimines under visible light photocatalytic conditions. *Chem. Commun.* **2017**, *53*, 7764. (b) Fernández-Salas, J. A.; Maestro, M. C.; Rodríguez-Fernández, M. M.; García-Ruano, J. L.; Alonso, I. Intermolecular Alkyl Radical Additions to Enantiopure N-tert-Butanesulfinyl Aldimines. *Org. Lett.* **2013**, *15*, 1658. (c) Zhong, Y.-W.; Dong, Y.-Z.; Fang, K.; Izumi, K.; Xu, M.-H.; Lin, G.-Q. A Highly Efficient and Direct Approach for Synthesis of Enantiopure β -Amino Alcohols by Reductive Cross-Coupling of Chiral N-tert-Butanesulfinyl Imines with Aldehydes. *J. Am. Chem. Soc.*, **2005**, *127*, 11956–11957. (d) Robak, M. T.; Herbage, M. A.; Ellman, J. A. Synthesis and Applications of tert-Butanesulfinamide. *Chem. Rev.*, **2010**, *110*, 3600–3740.
 28. See Supporting Information
 29. For research performed concurrent with this work: Ni, S.; Garrido-Castro, A. F.; Merchant, R. R.; deGruyter, J. N.; Schmitt, D. C.; Mousseau, J. J.; Gallego, G. M.; Yang, S.; Collins, M. R.; Qiao, J. X.; Yeung, K.; Langley, D. R.; Poss, M. A.; Scola, P. M.; Qin, T.; Baran, P. S. A General Amino Acid Synthesis Enabled by Innate Radical Cross-Coupling. *Angew. Chem. Int. Ed.* **2018**, *10.1002/anie.201809310*
 30. Beckwith, A. L. J.; Hay, B. P. Kinetics of the reversible β -scission of the cyclopentyl radical. *J. Am. Chem. Soc.*, **1989**, *111*, 230.
 31. Saladrigas, M.; Bosch, C.; Saborit, G. V.; Bonjoch, J.; Bradshaw, B. Radical Cyclization of Alkene-Tethered Ketones Initiated by Hydrogen-Atom Transfer. *Angew. Chem. Int. Ed.* **2018**, *57*, 182.
 32. Pitzer, L.; Sandfort F.; Strieth-Kalthoff, F.; Glorius, F. Intermolecular Radical Addition to Carbonyls Enabled by Visible Light Photoredox Initiated Hole Catalysis. *J. Am. Chem. Soc.*, **2017**, *139*, 13652.
 33. Reference 32 has a single example.
 34. Kochi, J. K.; Powers, J. W.; Mechanism of reduction of alkyl halides by chromium(II) complexes. Alkylchromium species as intermediates. *J. Am. Chem. Soc.* **1970**, *92*, 137.
 35. (a) Bakac, A.; Espenson, J. H. Unimolecular and bimolecular homolytic reactions of organochromium and organocobalt complexes. Kinetics and equilibria. *J. Am. Chem. Soc.* **1984**, *106*, 5197. (b) Espenson, J. H.; Sellers, T. D. Kinetics and mechanism of alkylchromium formation in the reductive cobalt-carbon bond cleavage of alkylcorrins by chromium(II). *J. Am. Chem. Soc.*, **1974**, *96*, 94.
 36. This reactivity has also been used for the coupling of unactivated alkyl halides and aldehydes using Cr^{+2} : Takai, K.; Nitta, K.; Fujimura, O.; Utimoto, K. Preparation of alkylchromium reagents by reduction of alkyl halides with chromium(II) chloride under cobalt catalysis. *J. Org. Chem.*, **1989**, *54*, 4732.; as well as the coupling of dienes with aldehydes: Takai, K.; Toratsu, C. B12-Catalyzed Generation of Allylic Chromium Reagents from 1,3-Dienes, CrCl_2 , and Water. *J. Org. Chem.* **1998**, *63*, 6450. In more recent examples, combination of Co/Cr have been used in: Xiong, Y.; Zhang, G. Enantioselective 1,2-Difunctionalization of 1,3-Butadiene by Sequential Alkylation and Carbonyl Allylation. *J. Am. Chem. Soc.* **2018**, *140*, 2735.
 37. (a) Steib, A. K.; Kuzmina, O. M.; Fernandez, S.; Flubacher, D.; Knochel, P. Efficient chromium(II)-catalyzed cross-coupling reactions between Csp² centers. *J. Am. Chem. Soc.* **2013**, *135*, 15346. (b) Oral-rat LD 50 for Cr_2O_3 > 2700 mg/kg. Chromium(III) and its Inorganic Compounds. **2014**, The MAK-Collection for Occupational Health and Safety. doi:10.1002/3527600418.mb1606583vee4614.
 38. Gil, A.; Albericio, F. and Álvarez, M. Role of the Nozaki–Hiyama–Takai–Kishi Reaction in the Synthesis of Natural Products. *Chem. Rev.* **2017**, *117*, 8420.

39. Kurosu, M.; Lin, M. H.; Kishi, Y. Fe/Cr- and Co/Cr-mediated catalytic asymmetric 2-haloallylations of aldehydes. *J. Am. Chem. Soc.* **2004**, *126*, 12248.
40. Co^{+3} is the oxidation state necessary for formation of the Co-H with the silane. Use of external metal reductants typically lead to formation of an inactive Co^{+2} .
41. Different attempts to make the chromium catalytic have been so far unsuccessful; presumably the additives necessary to dissociate the Cr-O bond, as proposed in standard Cr-catalyzed reactions, can also react with the salen ligand from the cobalt (see: Namba, K.; Kishi Y. New catalytic cycle for couplings of aldehydes with organochromium reagents. *Org. Lett.* **2004**, *6*, 5031.) Recent examples of combining Ir-photocatalysis with Cr where dissociation of the putative Cr-O bond seems to occur without an external agent, could in a future be explored for these endeavors (see: Schwarz, J. L.; Schäfers, F.; Tlahuext-Aca, A.; Lückemeier, L.; Glorius, F.; Diastereoselective Allylation of Aldehydes by Dual Photoredox and Chromium Catalysis. *J. Am. Chem. Soc.* **2018**, *140*, 12705.)
42. The chiral lactone auxiliary has been used in Ref. 21d.
43. Addition of Lewis basic ligands did not change diastereoselectivity and tended to suppress reactivity.
44. A mechanism in which subsequent alkyl transfer occurs via a free radical intermediate cannot be ruled out (see Ref. 35). Formation of an intermediate alkyl-chloride that leads to formation of product is unlikely, given that we do not see reaction through the alkyl-chloride in **46**.
45. Control experiments with the alkyl-halide during the same period of time yields no product under these conditions.
46. Aluminum- and borohydrides are known to reduce Cr^{+3} into Cr^{+2} , perhaps reduction by the silane or a derivative could occur through a similar mechanism. See: Fürstner, A. Carbon-Carbon Bond Formations Involving Organochromium(III) Reagents. *Chem. Rev.*, **1999**, *99*, 991.
47. The low yield might be suggestive of possible decomposition of the ligand under these conditions. See Andrez, J.; Guidal, V.; Scopelliti, R.; Pécaut, J.; Gambarelli, S.; Mazzanti, M. Ligand and Metal Based Multielectron Redox Chemistry of Cobalt Supported by Tetradentate Schiff Bases. *J. Am. Chem. Soc.*, **2017**, *139*, 8628.
48. Outer sphere electron transfer from Cr to Co: Candlin, J. P.; Halpern, J.; Trimm, D. L. Kinetics of the Reduction of Some Cobalt(III) Complexes by Chromium(II), Vanadium (II), and Europium (II). *J. Am. Chem. Soc.*, **1964**, *86*, 1019.
49. An alternative mechanism of SH_2 , as finally proposed in Ref. 35a, is not likely operative given that Co^{+2} is not catalytically active in our system (Table 1, entry 5). A mechanism similar to our previously reported Co/Ni dual catalysis involving oxidation of the alkyl- Co^{+3} to an alkyl- Co^{+4} prior homolysis and concomitant reduction of the Cr^{+3} to Cr^{+2} (see Ref. 7) is also unlikely given that control experiments indicate there is no reactivity with Cr^{+3} (Figure 4).
50. Pyea, D. R.; Mankad, N. P. Bimetallic catalysis for C-C and C-X coupling reactions. *Chem. Sci.* **2017**, *8*, 1705.