

## Structure-reactivity relationships on substrates and inhibitors of the lysine deacylase

### Sirtuin 2 from *Schistosoma mansoni* (*SmSirt2*)

Daria Monaldi<sup>1</sup>, Dante Rotili<sup>2\*</sup>, Julien Lancelot<sup>3</sup>, Martin Marek<sup>4</sup>, Nathalie Wössner<sup>1</sup>, Alessia Lucidi<sup>2</sup>, Daniela Tomaselli<sup>2</sup>, Elizabeth Ramos-Morales<sup>4</sup>, Christophe Romier<sup>4</sup>, Raymond J. Pierce<sup>3</sup>, Antonello Mai<sup>2\*</sup> and Manfred Jung<sup>1\*</sup>

<sup>1</sup>Institute of Pharmaceutical Sciences, University of Freiburg, Albertstr. 25, 79104, Freiburg, Germany.

<sup>2</sup>Dipartimento di Chimica e Tecnologie del Farmaco “Sapienza” Università di Roma, 00185 Rome, Italy

<sup>3</sup>Université de Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 8204 - CIIL - Centre d'Infection et d'Immunité de Lille, 59000 Lille, France

<sup>4</sup>Département de Biologie Structurale Intégrative, Institut de Génétique et Biologie Moléculaire et Cellulaire (IGBMC), Université de Strasbourg (UDS), CNRS, INSERM, 67404 Illkirch Cedex, France

\* corresponding authors, E-mail: dante.rotili@uniroma1.it, antonello.mai@uniroma1.it, manfred.jung@pharmazie.uni-freiburg.de

### ABSTRACT

The only drug currently available for treatment of the neglected disease Schistosomiasis is Praziquantel, and the possible emergence of resistance makes research on novel therapeutic agents necessary and urgent. To this end, the targeting of *Schistosoma mansoni* epigenetic enzymes, which regulate the parasitic life cycle, emerged as promising approach. Due to the strong effects of human Sirtuin inhibitors on parasite survival and reproduction, *Schistosoma* sirtuins were postulated as potential therapeutic targets. *In vitro* testing of synthetic substrates of *S. mansoni* Sirtuin 2 (*SmSirt2*) and kinetic experiments on a myristoylated peptide demonstrated

lysine long chain deacylation as an intrinsic *SmSirt2* activity in addition to its known deacetylase activity for the first time. Focused *in vitro* screening of the GSK Kinetobox library and structure-activity relationships (SAR) of identified hits, led to the first *SmSirt2* inhibitors with activity in the low micromolar range. Several *SmSirt2* inhibitors showed potency against both larval schistosomes (viability) and adult worms (pairing, egg laying) in culture without general toxicity to human cancer cells.

## INTRODUCTION

Schistosomiasis is a neglected tropical disease, affecting millions of people in tropical and sub-tropical countries and causing more than 300000 deaths per year.<sup>1</sup> One of its major causative agents is the blood fluke *Schistosoma mansoni*, characterized by a complex life cycle, where the parasite is developed through four morphologically distinct forms, with two of them, schistosomula and adult worms, present in the final human host. Praziquantel is so far the gold standard for the treatment of schistosomiasis, showing several positive features like low cost, high efficacy and low toxicity,<sup>2</sup> that made possible its use for mass treatment campaigns.<sup>3</sup> Long term mass treatment with Praziquantel has led to a reduction in mortality and morbidity in endemic areas,<sup>4,5</sup> but also to a reduction in treatment efficacy and, in some cases, to the isolation of resistant strains.<sup>6,7,8,9</sup> This aspect, in association with the drug inactivity on larval stages and its unknown mechanism of action,<sup>2,7</sup> render relying on only one drug inadequate in the long term and highlights the need for the development of novel therapeutic agents. Different approaches have been already studied, for example the use of artemisinins<sup>10</sup> and benzodiazepines,<sup>11</sup> but none of them is similar to Praziquantel in terms of efficacy, safety and cost.

The publication of the *Schistosoma mansoni* genome sequence<sup>12</sup> made the research and validation of novel therapeutic targets a particularly promising paradigm for the development of novel antischistosomal drugs. A drug repurposing approach,<sup>13</sup> based on targeting orthologues of proteins already targeted in other pathologies,<sup>10</sup> has demonstrated potential for developing compounds able to selectively inhibit parasitic enzymes without affecting the correspondent human isoforms. Moreover, since schistosomes can be considered similar to cancer cells in terms of intensive metabolic activity and invisibility to host immune system,<sup>14</sup> our interest is focused on histone deacetylase enzymes (HDACs), which are able to remove acetyl groups from lysine residues of histones and other proteins and are already known cancer drug targets with clinically

approved drugs. The essentiality of several histone deacetylases for growth and survival has already been demonstrated in a variety of parasite genera, including *Plasmodium*, *Leishmania*, *Trypanosoma* and *Schistosoma*.<sup>15</sup> The complex life cycles of these parasites are subject to complex epigenetic regulation and the selective inhibition of enzymes involved in these processes, including HDACs, represents a valid therapeutic strategy. For example, in the cases of *Plasmodium falciparum* and *Trypanosoma brucei* several molecules display selective *in vitro* and *in vivo* inhibitory activity against parasitic Zn<sup>2+</sup>-dependent HDACs (class I and II).<sup>15</sup> This approach has also led to the development of selective inhibitors of *S. mansoni* HDAC8, including a mercaptoacetamide analogue of suberoylanilide hydroxamic acid (SAHA),<sup>16</sup> alkylhydroxamates<sup>17</sup> and benzoylhydroxamates,<sup>18</sup> characterized by submicro or nanomolar IC<sub>50</sub> values for *SmHDAC8* with good selectivity over hHDAC1, hHDAC6 and, in some cases, also over hHDAC8. Some of these compounds also had low micromolar EC<sub>50</sub> values for killing schistosome larvae and abolished pairing stability and egg laying in adult worms.<sup>16-18</sup>

The situation is different for NAD<sup>+</sup>-dependent HDACs (class III, Sirtuins), where, despite their evident potential as human anticancer and metabolic disease targets,<sup>19</sup> only one compound (Selisistat) has reached clinical trials so far as a potential treatment for Huntington's disease. In parasites, sirtuin isoforms have been identified in *Plasmodium falciparum* (*PfSir2A* and *PfSir2B*),<sup>20,21</sup> in *Trypanosoma cruzi* (*TcSir2rp1* and *TcSir2rp3*)<sup>22,23</sup> and in *Leishmania* (*LmSir2rp1* and *Lsir2rp1*),<sup>24,25</sup> but most *in vitro* tested inhibitors showed modest activity and/or lack of selectivity.<sup>15</sup> An exception is provided by bisnaphthalimidopropyl derivatives that showed significant activity *in vitro* and in mice chronically infected with *Leishmania infantum*.<sup>26</sup>

In 2013 Lancelot *et al.* published the identification and characterisation of five *Schistosoma mansoni* sirtuins (*SmSirt1*, *SmSirt2*, *SmSirt5*, *SmSirt6* and *SmSirt7*) as orthologues of their respective mammalian counterpart isoforms.<sup>27</sup> Furthermore, it was demonstrated that Sirtinol and

Salermide, known inhibitors of human Sirtuin 1 and 2 (hSirt1 and hSirt2),<sup>28,29,30</sup> in addition to being inducers of selective apoptosis in cancer cell lines,<sup>29,31</sup> and showing protective effects in a muscular dystrophy model in nematodes,<sup>32</sup> have pro-apoptotic effects in schistosome larvae (schistosomula), through DNA fragmentation, and markedly reduce pairing stability and egg production in adult worms. These features support the potential of *Schistosoma* sirtuins as drug targets for the development of novel and selective antischisosomal drugs.

Beyond their deacetylase activity, human sirtuins are also implicated in the removal of short, medium and long fatty acyl groups from lysine residues of histones and non-histones proteins.<sup>33,34,35,36,37,38</sup> Lysine acylation has been identified as a posttranslational modification and it is strongly connected to regulation of metabolism. In fact, metabolic intermediates are used for this process and enzymes implicated in energy pathways are, in many cases, subject to these modifications.<sup>36,39,40</sup> Although deacylation of acyllysine is a common feature for all mammalian sirtuins, each isoform is characterised by a different pattern of specificity and efficiency for deacylation that may be quite distinct from the deacetylase activity.<sup>41</sup> In parasites, studies regarding lysine deacylation are lacking so far with the exception of the medium and long fatty acyl chain removal by PfSir2A in *Plasmodium falciparum*.<sup>42</sup> In a previous study, we established a homogeneous in vitro assay for the determination of SmSirt2 deacetylase activity which uses the readily available Z-(Ac)Lys-AMC (ZMAL, **1**) and represents an optimal tool for cost efficient high throughput campaigns.<sup>43</sup> In order to further characterize the function of SmSirt2, we performed and report here the analysis of SmSirt2 lysine deacylation activity by the use of both lysine-derived small molecule and oligopeptidic substrates.


Moreover, with the aim of finding novel and selective drug-like inhibitors of SmSirt2, we present an extensive structure-activity relationship study concerning novel hits identified by an *in vitro* screening of the Kinetobox library,<sup>44</sup> provided by GSK. This library is constituted by compounds

that were shown to be potent and specific inhibitors of growth of *Leishmania donovani*, *Trypanosoma cruzi* and *Trypanosoma brucei* with low human cellular cytotoxicity.<sup>44</sup> The kinetoplastid parasites *Leishmania sp.* and *Trypanosoma sp.* are characterized by complex morphological changes and an involvement of epigenetic regulation during their life cycle, and we postulated that the screening of this diverse library would be a good starting point for our study on *S. mansoni* aimed at identifying novel chemical entities able to interfere with parasite growth. *S. mansoni* is phylogenetically very distant from kinetoplastids, but has similar dynamic phenotypic changes through the different life stages, some of them implying epigenetic modifications.<sup>45</sup> This inhibitor collection is freely available and provides a set of 592 compounds with diverse structural features, potentially providing novel chemical space for chemically yet uncharted targets. Here we present new hits for *SmSirt2* (as well as *hSirt2*) and present initial SAR to further characterize ligand affinity and specificity to *SmSirt2*. In particular, we can show for the first time that *SmSirt2* is a druggable target with selectivity over *hSirt2*. Moreover, several of the characterized *SmSirt2* inhibitors were active against both schistosomula larvae and adult worms in culture. Some of the negative controls showed activity on *Schistosoma* as well, but they were also toxic against human cancer cells in culture. In contrast, selective *SmSirt2* inhibitors did not show toxicity in mammalian cells, further supporting the use of *SmSirt2* as a valuable drug target in schistosomes.

## RESULTS


**Short-, medium-, and long-chain deacylation activity of *SmSirt2*.** To extend the biochemical characterization of *SmSirt2* activity, we studied its ability to deacylate long chain fatty acids from the  $\epsilon$ -amino group of lysine substrates. We synthesized seven analogues of the *SmSirt2* substrate ZMAL (Z-(Ac)Lys-AMC) **1**, ie **3a** (Z-(But)Lys-AMC), **3b** (Z-(Hex)Lys-AMC), **3c** (Z-(Oct)Lys-AMC), **3d** (Z-(Dec)Lys-AMC), **3e** (Z-(Lau)Lys-AMC), **3f** (Z-(Myr)Lys-AMC) and **3g** (Z-(Pal)Lys-AMC), by replacing its acetyl group with short, medium and long acyl chains. Substrate **1** was synthesized according to published procedures.<sup>46</sup> For the preparation of **3a-g**, the appropriate acyl chlorides (commercially available for **3a-d,g**, and synthesized for **3e** and **3f** by reaction of the corresponding lauric or myristic acid with thionyl chloride) were treated with Z-Lys-OH leading to the formation of the  $\epsilon$ -acyl-Z-Lys-OH **2a-g**, which were converted into the desired substrates **3a-g** by treatment with phosphorus oxychloride and 7-amino-4-methylcoumarin (AMC) (Scheme 1).

### Scheme 1. Synthesis of **3a-g**<sup>a</sup>



<sup>a</sup>Reagents and conditions: a) Z-Lys-OH, 1 M sodium hydroxide, water, rt, 20-45 min; b) AMC (structure shown), phosphoryl chloride, dry pyridine, -15 °C, 40 min to 3h.


Then we used the acyl-lysine substrates **3a-g**, in comparison to the acetylated **1**, in the homogeneous fluorescence based assay,<sup>43</sup> in order to evaluate the ability of *SmSirt2* to catalyse their conversion into free lysine substrate. Compounds **1**, **3a-g** were tested at 10.5  $\mu$ M with one time point measurement, according to a published procedure for **1**.<sup>43</sup> As shown in Figure 1A (Table S1 in Supporting Information), the measured substrate conversion increases with increasing chain length from acetyl to hexanoyl analogues (see **1**, **3a** and **3b**), then exhibit an opposite trend from the octanoyl to the palmitoyl analogue (from **3c** to **3g**). When tested with hSirt2 (Figure 1B, Table S2), **1** and **3a-g** showed a somewhat discontinuous pattern of conversion, with **3a** (butyryl) and **3b** (hexanoyl) as the best substrates. Unfortunately, solubility issues did not allow the measurement of  $K_m$  values for any of the ZMAL analogues in order to obtain a more quantitative overview, however we performed a quantitative analysis on peptidic substrates (see below).



**Figure 1.** In vitro conversion of **1** (acetyl) and **3a-g** (see Scheme 1) by *SmSirt2* (A) and hSirt2 (B). A pure AMC sample is measured to simulate a (hypothetical) conversion of 100% for comparison. Errors are represented as standard deviation of the mean (SD, n=3).




Nicotinamide (NA) is a physiological sirtuin inhibitor which, once released from  $\text{NAD}^+$ , stays in a subpocket of the enzyme (pocket C) and, by a rebound mechanism, can block the enzymatic activity.<sup>47</sup> In hSirt2 the lysine acyl chain pocket is in a close contact with the C pocket, determining the possibility for NA potency to be influenced by the substrate acyl chain length.<sup>34</sup> To study the effect of short-, medium- and long-chain lysine substrates on the relative NA potency for *Sm*Sirt2 inhibition, we measured  $\text{IC}_{50}$  values for NA in presence of our ZMAL analogues **1**, **3a-g**. As shown in Figure 2 (Table S3), the  $\text{IC}_{50}$  values of NA are similar using substrates up to the decanoyl group (**3c**), after which they increase with the growth of the acyl chain length. Since an opposite trend has been published for acylated peptides and hSirt2,<sup>34</sup> we can hypothesize that, despite the structural identity<sup>27,48</sup> between the schistosome and the human Sirt2, these two enzymes are characterized by different kinetics. It could be possible that the putative structural differences in the active site/C pocket between hSirt2 and smSirt2 can explain the obtained results, but due to a lack of structural data, this is yet speculative.




**Figure 2.**  $\text{IC}_{50}$  values of NA against *Sm*Sirt2 in the presence of the acetylated substrate **1** and the acylated ZMAL analogues **3a-g** (see Scheme 1). Errors are represented as standard error of the mean (SEM, n=3).  $\text{IC}_{50}$  values are reported in Table S3 (Supporting information).

In order to gain a more quantitative insight into the interplay between *SmSirt2* and its acylated substrates, we performed a kinetic analysis through the use of a HPLC-based assay using more soluble peptides, not tagged with a fluorescent label to rule out potential artifactual effects.  $K_m$ ,  $k_{cat}$  and  $k_{cat}/K_m$  of *SmSirt2* were measured in the presence of an acetylated (PSDKac) and a myristoylated peptide (PSDKmyr) derived from  $\alpha$ -tubulin which is a Sirt2 substrate.<sup>49</sup> As shown in Figure 3, the acyl chain elongation in PSDKmyr determined an increase of affinity for *SmSirt2* without affecting the enzymatic turnover, with consequent threefold higher catalytic efficiency in the presence of PSDKmyr than with PSDKac (Table 1).

Moreover, since in hSirt2 the acylated substrate binds the enzyme before  $NAD^+$ ,<sup>50</sup> the length of the acyl group could influence the  $NAD^+$  binding affinity for *SmSirt2* and have, more generally, an influence on the  $NAD^+$  reaction. Consequently, the  $K_m$  values for  $NAD^+$  were also analyzed in the presence of PSDKac and PSDKmyr, respectively. As shown in Figure 4, the extension of the peptide chain length increases the binding affinity of  $NAD^+$  to *SmSirt2* and the catalytic turnover. In fact, a threefold higher  $k_{cat}$  is the reason for a better catalytic efficiency in presence of PSDKmyr than with PSDKac (Table 1). There was no inhibition of ZMAL conversion by PSDKac in a concentration of 300  $\mu M$  (data not shown).



**Figure 3.** Michaelis-Menten plots for *SmSirt2* dependent deacetylation (A) and demyristoylation (B). Error bars indicate standard error of the mean ( $n \geq 2$ ).



**Figure 4.** Michaelis-Menten plots for  $\text{NAD}^+$  with PSDKac (A) and with PSDKmyr (B) in *SmSirt2*. Error bars indicate standard error of the mean ( $n \geq 2$ ).

**Table 1.** Kinetic parameters for PSDKac, PSDKmyr and  $\text{NAD}^+$  for *SmSirt2*.

	$K_m$ ( $\mu\text{M}$ ) <sup>a</sup>	$k_{\text{cat}}$ ( $\text{min}^{-1}$ ) <sup>a</sup>	$k_{\text{cat}}/K_m$ ( $\text{s}^{-1}\text{M}^{-1}$ ) <sup>a</sup>
<b>PSDKac</b>	$40.9 \pm 12.2$	$(31.8 \pm 2.4) 10^{-2}$	$(1.3 \pm 0.3) 10^2$
<b>PSDKmyr</b>	$15.1 \pm 2.5$	$(33.8 \pm 1.3) 10^{-2}$	$(3.7 \pm 0.9) 10^2$
<b><math>\text{NAD}^+</math> (PSDKac)</b>	$33.9 \pm 8.5$	$(10.3 \pm 0.6) 10^{-2}$	$(5.1 \pm 1.8) 10^1$
<b><math>\text{NAD}^+</math> (PSDKmyr)</b>	$13.3 \pm 2.3$	$(32.3 \pm 1.3) 10^{-2}$	$(4.1 \pm 0.9) 10^2$
<b>ZMML</b>	$37.5 \pm 12.7$	$(50.6 \pm 4.9) 10^{-4}$	$(1.3 \pm 0.4) 10^2$
<b><math>\text{NAD}^+</math> (ZMAL)</b>	$75.8 \pm 8.1$	$(71.3 \pm 3.2) 10^{-4}$	$(9.4 \pm 4.0) 10^1$
<b><math>\text{NAD}^+</math> (ZMML)</b>	$9.3 \pm 1.9$	$(51.8 \pm 1.6) 10^{-4}$	$(5.6 \pm 0.9) 10^2$

<sup>a</sup>Reported as  $\pm$  SE.  $K_m$  for ZMAL (59  $\mu\text{M}$ ) taken from lit<sup>43</sup>

To further characterize the deacylation reactivity, we compared the potency of selected reference hSirt2 deacetylation inhibitors (NA, SirReal1 and 2,<sup>51</sup> AEM2<sup>52</sup>) with their ability to inhibit the demyristoylation reaction. As reported in Table 2, while in hSirt2 these compounds show higher

inhibitory potency for deacetylation than for demyristoylation, they did not inhibit either reaction catalysed by *SmSirt2*, with the only exception of NA.


**Table 2. Activity of hSirt2 standard inhibitors with 1 or 3f as substrates.**

Compound	<i>SmSirt2</i> IC <sub>50</sub> [μM] <sup>a</sup> or % inhibition at 20 μM		hSirt2 IC <sub>50</sub> [μM] <sup>a</sup> or % inhibition at 20 μM	
	1	3f	1	3f
NA	23.1 ± 1.8	420 ± 11.6	49.8 ± 4.6 <sup>b</sup>	8.7 ± 0.5 %
SirReal1	24.2 ± 0.8 %	< 15 %	3.7 ± 0.8 <sup>c</sup>	< 15 %
SirReal2	< 15 %	< 15 %	0.4 ± 0.1 <sup>c</sup>	< 15 %
AEM2	< 15 %	< 15 %	2.5 ± 0.2 <sup>b</sup>	42.8 ± 3.0%

<sup>a</sup>IC<sub>50</sub> are reported as ± SEM (n=3), for inhibition % ± SD (n = 3). <sup>b</sup>Reference <sup>53</sup>. <sup>c</sup>Reference <sup>51</sup>.

***SmSirt2* inhibition.** Using the *in vitro*<sup>43</sup> assay described above we initially screened the FDA-approved drugs library (provided by Enzo Life Sciences (ELS) AG, Switzerland) that had already been tested on schistosomula and adult worms.<sup>54,55</sup> Idebenone (**4**) emerged as an overlapping compound that was known to block the growth of schistosomula, showed good *SmSirt2* inhibition potency and selectivity over hSirt2<sup>56</sup> (Figure 5). Although Idebenone probably has a pleiotropic mode of action, this lent credence to our approach to identify druggable molecules from a diverse set of compounds that also show antischistosomal activity. We next proceeded with the GSK Kinetobox library, which was initially tested at the fixed assay concentration of 25 μM. Since we were looking for compounds with an IC<sub>50</sub> in the low micromolar range, we focused our attention only on compounds with inhibitory potency higher than 50% at that concentration. Furthermore, to test whether hits quench the AMC signal, inhibit the detection enzyme trypsin or are

autofluorescent compounds, all promising candidates were subjected to counter-analysis in order to exclude false positives and false negatives (data not shown). Compounds showing no assay interference and a dose-dependent activity were then tested on hSirt2 to evaluate their selectivity. Among the 592 tested compounds, we identified three hits characterized by potency in the low micromolar range: TCMDC-143159 (**5**), TCMDC-143362 (**6**) and TCMDC-143295 (**7**) (Figure 5). Due to reported toxicity of the anabasine ring,<sup>57</sup> present in **5**, we decided initially to resynthesize **6** and **7** to confirm their activity, and to modify their structures in order to obtain analogues with improved activity and selectivity profiles. Since the inhibition of hSirt2 is an undesired feature in new antischistosomal, but might identify new scaffolds for optimization of human sirtuin inhibitors we followed the lead of **6** as an unselective compound to some extent but focused mostly on the selective compound **7**.



**Figure 5. A) Structure of antischistosomal treatment standard Praziquantel (PZQ)**  
**B) Structures and *in vitro* data of idebenone (4) and hits from the Kinetobox library (5-7).** For IC<sub>50</sub> values, errors are represented as  $\pm$  SEM (n = 3), while for inhibition % as  $\pm$  SD (n = 3).


### **Synthesis and initial SAR of 1,2,4-oxadiazolyl compound 6.**

As previously cited, Feldmann *et al.* recently published the crystal structure of hSirt2 in complex with myristoylated peptides, showing the placement of the acyl chain in a hydrophobic pocket and the conservation of this localization during the entire catalytic reaction.<sup>34,51</sup> Since this could also be the case in *SmSirt2*, after establishing its synthesis, initial optimization of **6** was attempted with the replacement of the acetyl group with a longer fatty acyl group, such as an octanoyl (**11**) or a decanoyl (**12**) chain, in order to potentially address such a pocket. The synthesis of **6** and its analogues **11** and **12** is outlined in Scheme 2. A nucleophilic substitution between the commercially available 2-(piperidin-3-yl) acetic acid ethyl ester and 3-fluoro-2-methylbenzyl bromide was performed in order to obtain the intermediate **8**. The reaction between the requisite acyl chlorides and 4-aminobenzonitrile yielded the 4-cyanoanilides **9b** and **9c** which, together with the commercially available 4-cyanoacetanilide **9a**, were treated with hydroxylamine hydrochloride to give the benzamidoximes **10a-c**. Subsequent condensation between **10a,b** and **8** afforded the required compounds **6** and **11**, and the reaction between **10c** and the acyl chloride of **8**, obtained by hydrolysis of the ethyl ester and subsequent chlorination, gave the final compound **12**.

When tested against *SmSirt2* and hSirt2 at 25  $\mu$ M, the hit **6** was confirmed to have inhibitory activity from the newly synthesized sample whereas the octanoyl and decanoyl analogues **11** and **12** were practically inactive against both parasite and human enzymes (**6**: *SmSirt2* IC<sub>50</sub> = 14.5  $\pm$

0.6  $\mu\text{M}$ ; hSirt2  $\text{IC}_{50} = 8.0 \pm 1.1 \mu\text{M}$ ; **11**: *SmSirt2* inh.  $30.5 \pm 3.5\%$ ; hSirt2 inh.  $<15\%$ . **12**: *SmSirt2* inh.  $<15\%$ ; Sirt2 inh.  $<15\%$ ).

### Scheme 2. Synthesis of compounds **6**, **11** and **12**.<sup>a</sup>




<sup>a</sup> Reagents and conditions: (a) 3-fluoro-2-methylbenzylbromide, triethylamine, toluene,  $0^\circ\text{C} \rightarrow \text{rt}$ , 18 h; (b) 4-aminobenzonitrile, triethylamine, dry DCM,  $0^\circ\text{C} \rightarrow \text{rt}$ , 2 h,  $\text{N}_2$ ; (c) hydroxylamine hydrochloride, sodium carbonate, water/ethanol, reflux, 6-8 h; (d) **8**, **10a**, potassium carbonate, pyridine, reflux 8h, then  $\text{rt}$  72 h; (e) **8**, **10b**, potassium carbonate, pyridine, microwave,  $180^\circ\text{C}$ , 10 min, 300 Watt, then reflux 43 h and finally  $\text{rt}$  12 h; (f) from **8**: i) 1M lithium hydroxide, ethanol, 20 h,  $\text{rt}$ ; ii) thionyl chloride, DMF, dry DCM,  $-15^\circ\text{C} \rightarrow \text{r.t.}$ , 3h,  $\text{N}_2$ ; iii) **10c**, triethylamine, DMF, dry DCM,  $\text{rt}$ , 22 h, then  $150^\circ\text{C}$  3 h,  $\text{N}_2$ .

### Synthesis and SAR of pyrimidopyrimidine **7**.

To work on the structure of hit **7**, we started with a fragment-based approach with the development of analogues with a simplified structure, in order to identify the substructure(s) responsible of the inhibitory activity. In particular, since the 2,4,7-triaminopyrimidopyrimidine moiety of **7** (“part A”, Figure 6) could mimic the adenosine part of  $\text{NAD}^+$ , we synthesized a first generation of analogues keeping this portion fixed and introducing several modifications at the

Na (Figure 6). More precisely, we purchased parent pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine **13** and prepared derivatives where the Na is a secondary (**14**, **18** and **19**), tertiary (**20-23**), or inserted in a cyclic amine (**15-17**). While **14-17** are characterized by small amine substitution at the C7 position, the *N*<sup>7</sup>-benzylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamines **18-23** represent the most substituted compounds of the series, with changes at the Na, C $\beta$  and C4 benzyl positions (“part B”, Figure 6).



**Figure 6.** Overview of analogues of **7**: fragment-based approach and SAR studies.

As shown in Table 3 (see below), compounds **13-22** show complete loss of activity on *SmSirt2* and remain inactive on the human enzyme. Among the simplified analogues, only **23**, bearing a large phenyl substituent at the right part of the molecule (4-biphenyl portion, “part C”, Figure 6), displayed low *SmSirt2* inhibition. As a consequence, we designed a second generation of




analogues in which small focused modifications were applied, without leading to an excessive alteration of the prototype structure, but maintaining a high similarity to it. Specifically, we focused on the N $\alpha$ , C $\beta$  and C $\gamma$  positions of **7** (Figure 6) through the removal of the methyl or methoxy groups (**24-26**). We increased the steric hindrance of the C $\beta$  substituent (“part B”, Figure 6), by replacing the methyl with an ethyl, *isopropyl*, or phenyl group (**27-29**, **33**, **41**), and we explored the effect of replacement of the 4-methoxyphenoxy portion of **7** (“part C”, Figure 6) with other substituted phenoxy, 1-naphthyloxy, benzyloxy, phenylthio, aniline and benzamide moieties (**30-44**).

*Chemistry.* The pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine **13** is commercially available. The final compounds **14-44** were prepared by reaction between the commercial 4-amino-2-bromopyrimidine-5-carbonitrile and the appropriate amines **46** in anhydrous 2-methoxyethanol, in the presence of triethylamine at 80 °C. The obtained pyrimidine intermediates **47** were converted into the desired compounds **14-44** through condensation reactions performed at 150 °C with the guanidine free base in dry 2-methoxyethanol (Scheme 3). The amines **46a-h** used for the synthesis of the pyrimidine intermediates **47a-h** are commercially available and were purchased from vendors. The amines **46i-f'**, useful synthons for the synthesis of the final compounds **22-44**, were prepared by reaction of the corresponding phenyl ketones or aldehydes **45a-w** with 7 M ammonia (for secondary amines) or 2 M methylamine (for tertiary amines) solution in anhydrous methanol in the presence of titanium *isopropoxide* at room temperature, followed by the addition, after the formation of the imine intermediates, of the reducing agent sodium borohydride at 0 °C under nitrogen atmosphere (Scheme 4A). The phenyl ketones/aldehydes **45a-e** are commercially available. The ketones **45f-q,s-u** were synthesized by reaction between the appropriate aromatic alcohols or thiols and the 1-(4-fluorophenyl)ethan-1-one, -propan-1-one, -2-methylpropan-1-one, and -(phenyl)methanone in the presence of anhydrous potassium carbonate in dry DMF at 175 °C

(Scheme 4B). The 1-(4-((4-methoxybenzyl)oxy)phenyl)ethan-1-one **45r**,<sup>58</sup> the 1-(4-((4-methoxyphenyl)amino)phenyl)ethan-1-one **45v**,<sup>59</sup> and the N-(4-acetylphenyl)-4-methoxybenzamide **45w**<sup>60</sup> were prepared according to the literature. For the structures of the intermediates **45-47** see Supporting Information.


### Scheme 3. Synthesis of compounds **7**, **14-44**.<sup>a</sup>



R = 4-OMe; 4-Ph; 3-OMe-, 4-OMe-, 4-OCF<sub>3</sub>-, 4-SMe-, 4-OBn-, 3,4-(OMe)<sub>2</sub>-, 3,5-(OMe)<sub>2</sub>-, 3,4,5-(OMe)<sub>3</sub>-PhO;  
 4-OMe-1-naphthyl-O, 4-OMe-BnO; 4-OMe-, 3,5-(OMe)<sub>2</sub>-PhS; 4-OMe-PhNH; 4-OMe-PhCONH;  
 R<sub>1</sub> = H, Me, Et, Ph; R<sub>2</sub> = H, Me

<sup>a</sup>Reagents and conditions: (a) triethylamine, dry 2-methoxyethanol, 80 °C, 2.5-8 h; (b) guanidine free base, dry 2-methoxyethanol, 150 °C, 1.5-4.5 h.








**Scheme 4. Synthesis of the intermediate compounds 45 and 46.<sup>a</sup>**

















<sup>a</sup>Reagents and conditions: (a) 2 M methylamine in methanol, titanium *isopropoxide*, dry THF, N<sub>2</sub>, 5-6 h, rt, then sodium borohydride, N<sub>2</sub>, 2 h, rt; (b) 7 M ammonia in methanol, titanium *isopropoxide*, N<sub>2</sub>, 5 h, rt, then sodium borohydride, N<sub>2</sub>, 2 h, rt; c) anhydrous potassium carbonate, dry DMF, 175 °C, 5 h;

All the above compounds were tested *in vitro* against *SmSirt2*<sup>43</sup> and hSirt2, to study their selectivity for the parasitic enzyme (Table 3).<sup>56</sup> An amine and a pyrimidine synthetic intermediates (**46k** and **47m**, see Figure 6 and Table S4 for their structures) were also included in the list, to confirm the importance of the intact pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine scaffold for the inhibiting activity. The percentages of inhibition at 25 μM or the IC<sub>50</sub> values of the new compounds tested against *SmSirt2* as well as hSirt2 are reported in Table 3.

**Table 3. *In vitro* inhibition of *SmSirt2* and *hSirt2* by 7, 13-44, 46k, and 47m.**

					
compd	R	R <sub>1</sub>	R <sub>2</sub>	<i>SmSirt2</i> <sup>a</sup> % inhibition at 25 μM or IC <sub>50</sub> [μM]	<i>hSirt2</i> <sup>a</sup> % inhibition at 25 μM or IC <sub>50</sub> [μM]
7		Me	Me	23.7 ± 9.6	21.9 ± 2.5%
13				<15%	<15%
14				<15%	23.2 ± 0.8%
15				<15%	<15%
16				<15%	<15%
17				<15%	<15%
18	-OMe	H	H	<15%	<15%
19	H	Me	H	<15%	<15%
20	H	H	Me	<15%	<15%
21	H	Me	Me	<15%	<15%

22	-OMe	Me	Me	<15%	<15%
23	-Ph	Me	Me	24.8 ± 2.2%	<15%
24		Me	Me	27.6 ± 2.8%	<15%
25		Me	H	37.5 ± 5.6%	21.0 ± 3.4%
26		H	Me	<15%	<15%
27		Et	Me	12.8 ± 0.8	36.7 ± 7.7
28		<i>i</i> -Pr	Me	27.7 ± 3.8	57.4 ± 5.2%
29		Ph	Me	2.34 ± 0.2 <sup>b</sup>	22.1 ± 3.4%
30		Me	Me	23.1 ± 1.4	33.8 ± 0.4%
31		Me	Me	44.7 ± 4.4	35.3 ± 4.9%
32		Me	Me	12.5 ± 1.1	7.4 ± 3.9%
33		Ph	Me	3.3 ± 0.2	29.6 ± 1.9%
34		Me	Me	30.8 ± 3.0	29.2 ± 3.9%
35		Me	Me	20.3 ± 2.1%	37.5 ± 4.8%
36		Me	Me	49 ± 1.1%	<15%
37		Me	Me	40.3 ± 5.2%	70.4 ± 4.2%

<b>38</b>		Me	Me	18.4 ± 0.9%	30.6 ± 0.2%
<b>39</b>		Me	Me	46.2 ± 4.3%	61.7 ± 5.3%
<b>40</b>		Me	Me	14.9 ± 0.9	13.3 ± 1.6
<b>41</b>		Me	Me	4.3 ± 0.4	27.9 ± 1.8 %
<b>42</b>		Ph	Me	2.0 ± 0.1	21.5 ± 3.1
<b>43</b>		Me	Me	65.1 ± 7.2	40.8 ± 4.9%
<b>44</b>		Me	Me	41.9 ± 2.2%	<15%
<b>46k</b>				<15%	<15%
<b>47m</b>				<15%	<15%

<sup>a</sup>IC<sub>50</sub> ± SEM are reported (n=3), while inhibition % are reported as ± SD (n = 3). <sup>b</sup>maximum of observed inhibition: 60%.

As stated above, none of the fragments or simplified molecules **13-22** showed inhibitory activity on either the parasite or human enzymes. Only the derivatives carrying the 1-([1,1'-biphenyl]-4-yl)ethyl (**23**) or 1-(4-phenoxyphenyl)ethyl (**24**) substituent at N $\alpha$  (Figure 6) partially retained inhibition against *SmSirt2*, demonstrating the crucial role of the substituted phenoxyphenyl or

1,1'-biphenyl portion at the right side of the molecule (“part C”, Figure 6). The removal of the methyl group at either the 7 N $\alpha$  (see **25**) or C $\beta$  position (see **26**) (Figure 6) decreased or totally abolished the *SmSirt2* inhibitory activity, respectively. In contrast, the insertion at C $\beta$  of groups bigger than methyl [ethyl (**27**), *isopropyl* (**28**), and phenyl (**29**)], generally improved the *SmSirt2* inhibitory potency up to 10-fold, reaching with **29** the single-digit micromolar level (IC<sub>50</sub> = 2.34  $\mu$ M). Interestingly, **29** is more selective than **7** for *SmSirt2* over hSirt2 as judged by IC<sub>50</sub> values, but it reaches only a maximum of inhibition of 60 % which complicates the interpretation of the selectivity data. The *isopropyl* derivative **28** was an exception, displaying similar inhibitory potency as **7** against *SmSirt2*, and higher potency against hSirt2.

At the “part C” of the molecule (4-methoxyphenoxy moiety, Figure 6), no particular increase in activity or selectivity was obtained with the shift of the methoxy group from *para* to *meta* position (**30**), and the introduction of 3,4-dimethoxy (**31**) or 3,4,5-trimethoxy (**34**) groups at the phenoxy portion, as well as the replacement of the 4-methoxy with a 4-trifluoromethoxy (**35**), 4-methylthio (**36**), or 4-benzyloxy (**37**) group reduced the inhibitory potency against *SmSirt2*. However, the insertion of two methoxy groups at 3,5 position of the phenoxy moiety (**32**) led to 2-fold increase of potency against *SmSirt2* and improved selectivity over hSirt2. In this last compound, the further replacement of the methyl group at C $\beta$  with a phenyl ring (**33**) determined an additional increase of potency against *SmSirt2* (IC<sub>50</sub> = 3.3  $\mu$ M, 7-fold higher potency compared to **7**), confirming the positive SAR about C $\beta$  substitution with large groups. The replacement of the 4-methoxyphenoxy moiety of **7** with the larger 4-methoxy-1-naphthyloxy (**38**) or the longer 4-methoxybenzyloxy (**39**) group led to a decrease in potency against *SmSirt2* and, in the case of **39**, improved hSirt2 inhibition. Again in the 4-methoxyphenoxy group, the isosteric change oxygen-sulphur atom led to the 4-methoxyphenylthio analogue **40**, which produced a 1.6-

fold enhancement of *SmSirt2* inhibition ( $IC_{50} = 14.9 \mu\text{M}$ ), combined with improved inhibition of the human enzyme ( $IC_{50} = 13.3 \mu\text{M}$ ). The further change from the 4-methoxy to the 3,5-dimethoxy substitution at the phenylthio moiety of **40** provided **41**, that showed improved potency and selectivity against *SmSirt2* (*SmSirt2*  $IC_{50} = 4.3 \mu\text{M}$ , hSirt2 inhibition = 27.9 % at 25  $\mu\text{M}$ ). Moreover, the combination of the positive SAR of the series with the replacement of the C $\beta$  methyl of **41** with the C $\beta$  phenyl group gave **42**, the most potent compound of the series vs *SmSirt2* ( $IC_{50} = 2 \mu\text{M}$ ) and 10-fold selective over the human counterpart hSirt2 ( $IC_{50} = 21.5 \mu\text{M}$ ). The replacement of the 4-methoxyphenoxy portion with a 4-methoxyaniline (**43**) or 4-methoxybenzamide (**44**) group led to a decrease in potency against *SmSirt2*. Finally, the total absence of inhibitory activity against both parasitic and human enzymes by the intermediates **46k** and **47m** confirmed the importance of the intact pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine scaffold (“part A”, Figure 6) for the inhibitory activity of such compounds. The inhibiting activity of selected compounds **33**, **36**, **41**, **42** and **44** was also evaluated in the presence of the myristoylated substrate **3f** instead of the acetylated **1**. As reported in Table 4, in many cases no significant inhibition was measured in the presence of *SmSirt2* and hSirt2 with the exception of **33**, **41** and **42**, which show an inhibitory potency for the hSirt2 catalyzed demyristoylation in the low micromolar range. This may present a new starting point for the development of human Sirtuin inhibitors with a selectivity for long chain acyl removal, as so far a preference, if any, has been observed only for deacetylation.

**Table 4. Inhibitory activity of selected compounds 33, 36, 41, 42, and 44 against *SmSirt2* and hSirt2 with myristoylated 3f as substrate.**

Compound	<i>SmSirt2</i> (3f)	hSirt2 (3f)
	$IC_{50} [\mu\text{M}]^a$ or %	$IC_{50} [\mu\text{M}]^a$ or %



	inhibition at 20 μM	inhibition at 20 μM
<b>7</b>	16.8 ± 1.14 <sup>b</sup>	<15%
<b>33</b>	<15%	13.1 ± 2.2
<b>36</b>	<15%	38.1 ± 5.2%
<b>41</b>	15%	25.1 ± 4.1
<b>42</b>	<15%	15.6 ± 1.9
<b>44</b>	<15%	<15%

<sup>a</sup>IC<sub>50</sub> values are reported as ± SEM (n=3), while inhibition % as ± SD (n = 3). <sup>b</sup>maximum of inhibition: 45 %.


We also tested compounds **26**, **33**, **39** and **41** on human isoforms Sirt1 and 3 regarding deacetylation. All compounds showed less than 15% inhibition at 25 μM (data not shown) while 500 μM of nicotinamide led to complete inhibition as expected.<sup>51</sup> Evenmore, hSirt2 is the closest human homologue to *SmSirt2*<sup>48</sup> (see also Table S5, data not shown for hSirt1, 3).

#### ***In vitro* effects on schistosomula and adult worms.**

In order to evaluate whether these optimized compounds have effects on the parasite, **29**, **32**, **33**, **41** and **42** together with the prototype **7** were tested on schistosomula and adult worms. Two potential negative controls **26** and **39** were also included in the series. As shown in Figure S6, all tested compounds reduce the viability of cultured schistosomula at 10 and 20 μM, including **26** and **39**, which show low or no activity on the recombinant enzyme (Table 3).

Experiments on adult worms report a similar trend (Figure 7), where again the *SmSirt2* inhibitors as well as the potential negative controls reduce worm pairing and egg laying already after 24 hours of culture. Whilst the possibility that the enzyme inhibitors do exert their effects via


inhibition of *SmSirt2* cannot be ruled out, the strong effects that **26** and **39** also have on parasite viability and reproduction indicate that the effects of compounds shown in Figure 7 on the parasite may be at least in part due to the modulation of other targets (off-target effects).



**Figure 7.** Effects of **7** and its analogues on adult worms (AW) pairing at 10  $\mu$ M and 20  $\mu$ M (**A** and **B** respectively). Assay used: microscopy examination. Results after 24, 48 and 72 hours are represented with black, white and chequered bars respectively. Effect of these compounds on egg laying (**C**). Compounds **26** and **39** do not block *SmSirt2* in-vitro.

To get some further insight, we measured selected compounds **7**, **26**, **29**, **32**, **33**, **39** and **41** for toxicity on human cultured cancer cells using a MTS viability assay. In this assay, the compounds

that are inactive on *SmSirt2* but active on schistosomes also show toxicity on the human cells while the *SmSirt2* inhibitors with activity on the worms do not show this general toxicity (see Figure 8).



**Figure 8.** Growth inhibition of HL-60 acute myeloid leukaemia cells by compound 7 and selected analogues at 10 μM assay concentration.

## DISCUSSION AND CONCLUSION

The identification of a robust deacylase activity, supported by kinetic investigations, expands the range of possible biological activities relevant for *SmSirt2*. The high demyristoylation efficiency of this enzyme suggests the merit of undertaking future investigations on deacylation inhibition, in addition to deacetylation inhibition, as a new strategy to kill both schistosomula and adult worms. For inhibitors of hSirt2 with a thiomyristoylated lysine core (also called TM), it has been proposed that a dual deacetylase/demyristoylase activity is beneficial for cellular potency in cancer cells and selectivity over non-cancer cells.<sup>61</sup>

With the aim of finding potent and selective inhibitors of *SmSirt2*, we performed screening of the GSK-kinetobox library, composed of 592 compounds with proven activity against cultured kinetoplastids, namely *Leishmania donovani*, *Trypanosoma cruzi* and *Trypanosoma brucei*. From an initial test we identified **5**, **6** and **7** as valuable hits for further optimization. As outlined above, we focused our attention on **6** and **7**. Regarding **6**, the attempt to potentially target the enzymatic (possibly extended) C pocket through an octanoylated and decanoylated analogue, led to a loss in activity. Extensive SAR studies performed on **7** highlighted the crucial role of the presence of the intact pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (“part A” of the molecules), disubstituted at N7 (N $\alpha$  in Figure 6) with a methyl and a substituted 1-phenylethyl moiety (“part B”), further carrying a substituted phenoxy/phenylthio group at its C4 position (“part C”), recognized as crucial to elicit *SmSirt2* inhibitory potency and selectivity over hSirt2 (Figure 6). In particular, the presence at “part C” of the 4-methoxyphenoxy, 3,5-dimethoxyphenoxy and 3,5-dimethoxyphenylthio portion, combined with the the phenyl group at C $\beta$  position (“part B”) led to the most potent derivatives **29**, **33**, **41**, and **42** with IC<sub>50</sub> values in the single-digit micromolar range for *SmSirt2*. Experiments with **7** and its analogues on schistosomula and adult worms showed strong activity which was also present in enzymatically inactive compounds. However, while compounds **26** and

**39**, inactive on *SmSirt2*, also showed unspecific toxicity to human cancer cells, the selective *SmSirt2* inhibitors **7**, **29**, **32**, **33** and **41** were non-toxic to human cancer cells. Generally, the compounds show good properties in terms of druglikeness, e.g. logP values are between 2.7 (**32**) and 4.6 (**42**) (see Table S7). Molecular weight is mostly above 400 but only up to 525 Da maximum.

Thus, for the first time, we can show that *SmSirt2* can be drugged with selectivity over the human isotype (especially **29**, **33** and **41**) and the inhibitors block *Schistosoma* growth without general toxicity to human cells. This can be used as starting point for further optimization studies. In addition, leads for new inhibitors of hSirt2 have been identified, interestingly some with a preference for demyristoylation over deacetylation.

## EXPERIMENTAL SECTION

**Recombinant production and purification of *SmSirt2*.** Recombinant expression and purification of *SmSirt2* was done as previously described.<sup>43</sup> Briefly, overexpression was carried out in *E. coli* BL21(DE3) cells in 2 x Luria Broth (2 x LB) medium. The cells were grown to O.D.<sub>600</sub> of 1.2 at 37°C, then the culture was cooled down to 25°C and induction of expression was done by adding 0.5 mM final isopropyl-1-thio- $\beta$ -D-galactopyranoside (IPTG, Euromedex), in presence of 100  $\mu$ M ZnCl<sub>2</sub>. Harvested bacteria were resuspended in lysis buffer (400 mM NaCl, 10 mM Tris-HCl pH 8.0) and lysed by sonication. The lysate was clarified by centrifugation (17,000 rpm, JA-25.50 Beckman) for 1 h. The supernatant was loaded onto Talon Metal affinity resin (Clontech) pre-equilibrated with the lysis buffer. The 3C protease treatment was used to remove the His-tag from the recombinant protein, which was subsequently loaded onto HiLoad 16/60 Superdex 200 gel filtration column (Amersham Bioscience) equilibrated in 400 mM NaCl, 10 mM Tris-HCl pH 8.0, and 2 mM DTT. Finally, the protein was concentrated with an Amicon Ultra centrifugal filter units (Millipore) to reach a final concentration of 1.5 mg/ml as assayed by the A280 measurement (NanoDrop).

**Long chain deacylation.** Conversion in comparison to the theoretical maximal conversion (100% ZMAL conversion, obtained by measuring 10.5  $\mu$ M AMC) of **3a-g** by *SmSirt2* and hSirt2 and their inhibition were evaluated by homogeneous assay using 10.5  $\mu$ M assay concentration of potential substrates (prepared from 12.6 mM stock solution in DMSO and diluted with assay buffer) instead of ZMAL. Regarding hSirt2, 25 mM HEPES, 137 mM NaCl, 2.7 mM KCl, 1 mM MgCl<sub>2</sub>, 0.015% Triton X-100, pH= 8.0 was used as assay buffer. OriginPro 9.0 G and GraphPad 7.0 were used for the analysis of results.

**Kinetic analysis of PSDKac and PSDKmyr.** Deacylation reactions were evaluated by reversed phase HPLC (Kinetex XB-C18 column, 100 Å, 5  $\mu$ m, 250 x 4.6 mm) by monitoring the

formation of the deacylated product at 214 nm. Linear deacylation rates were determined by incubation of 47  $\mu\text{L}$  of *SmSirt2* solution (104 ng/ $\mu\text{L}$ , final assay concentration 80 ng/ $\mu\text{L}$ ) in assay buffer (25 mM HEPES, 137 mM NaCl, 2.7 mM KCl, 1 mM MgCl<sub>2</sub>, 0.015% Triton X-100, pH = 8.0) with 3  $\mu\text{L}$  of DMSO, 5  $\mu\text{L}$  of PSDK<sub>ac</sub> (Ac-Pro-Ser-Asp-Lys(acetyl)-Tyr-Ile-Gly-Gly-Trp-Trp-NH<sub>2</sub> custom synthesized by PSL, Heidelberg, Germany) or PSDK<sub>myr</sub> (Ac-Pro-Ser-Asp-Lys(myristoyl)-Tyr-Ile-Gly-Gly-Trp-Trp-NH<sub>2</sub> custom synthesized by PSL, Heidelberg, Germany) solution (prepared from 3.6 mM stocks in DMSO and diluted with assay buffer, concentration ranges 10-300  $\mu\text{M}$ ) and 5  $\mu\text{L}$  of NAD<sup>+</sup> (prepared from 6 mM stock in assay buffer and diluted in assay buffer; final assay concentration 500  $\mu\text{M}$ ). At 0, 1, 3, 5, 10, 20 and 30 minutes the deacylation was quenched with 6.7  $\mu\text{L}$  of TFA (10% in assay buffer, final assay concentration 1%), incubated for 5 min at 37 °C and then centrifuged for 10 minutes at 14000g. 55  $\mu\text{L}$  of the supernatant were transferred into HPLC vials and analyzed. Each experiment was done twice in duplicate. HPLC method to evaluate the deacylation: eluent A, H<sub>2</sub>O + 0.05% TFA; eluent B, acetonitrile + 0.05% TFA; 0 - 4 min, linear increase from B = 10% to B = 40%; 4 - 10 min, linear increase to B = 60%; 10 min, linear increase to B = 100%; 10-14 min, B = 100%; 14-16 min, linear decrease to B = 10%; 16 - 25 min, B = 10%) with a flowrate of 1 mL/min. The following method has been used to follow demyristoylation reactions: eluent A, H<sub>2</sub>O + 0.05% TFA; eluent B, acetonitrile + 0.05% TFA; 0 - 5 min, linear increase from B = 10% to B = 40%; 5-8.5 min, linear increase to B = 80%; 8.5-9.5 min, linear increase to B = 90%; 9.5-10.5 min, linear increase to B = 100%; 10.5-15 min, B = 100%; 15 min, B = 100%; 15 - 18 min, linear decrease to B = 10%) with a flowrate of 1 mL/min. The quantification of product peaks allowed the determination of deacylation rates and the data have been fitted to the Michaelis-Menten equation. GraphPad Prism has been used to determine  $K_m$ ,  $k_{cat}$ ,  $k_{cat}/K_m$ .

**Kinetic analysis of NAD<sup>+</sup><sub>PSDKac</sub> and NAD<sup>+</sup><sub>PSDKmyr</sub>.** Deacylation reactions were evaluated by reversed phase HPLC (Kinetex XB-C18 column, 100 Å, 5 µm, 250 x 4.6 mm) by monitoring the formation of the deacylated product at 214 nm. Linear deacylation rates have been determined by incubation of 47 µL of *SmSirt2* solution (104 ng/µL, final assay concentration 80 ng/µL) in assay buffer (25 mM HEPES, 137 mM NaCl, 2.7 mM KCl, 1 mM MgCl<sub>2</sub>, 0.015% Triton X-100, pH= 8.0) with 3 µL of DMSO, 5 µL of PSDKac or PSDKmyr (prepared respectively from 2.4 mM and 0.96 mM stock in DMSO and diluted with assay buffer, saturating assay concentrations) and 5 µL of NAD<sup>+</sup> (prepared from 6 mM stock in assay buffer and diluted with assay buffer. NAD<sup>+</sup> assay concentration range 10-250 µM). At 0, 1, 3, 5, 10, 20 and 30 minutes the deacylation was quenched with 6.7 µL of TFA(10% in assay buffer, final assay concentration 1%), incubated for 5 min at 37 °C and then centrifuged for 10 minutes at 14000g. 55 µL of the supernatant have been transferred into HPLC vials and analysed. Each experiment was done twice in duplicate. Deacetylation and demyristoylation were evaluated using the same HPLC methods described for PSDKac and PSDKmyr respectively. The quantification of product peaks allowed the determination of deacylation rates, and the data have been fitted to the Michaelis-Menten equation.  $K_{m,NAD^+}$ ,  $k_{cat,NAD^+}$  and  $k_{cat}/K_m$  have been determined by the use of GraphPad Prism 7.0.

**Kinetic analysis of ZMML.** Deacylation reactions were evaluated by a homogeneous fluorescence based assay. 14 µL of *SmSirt2* solution (104 ng/µL, final assay concentration 80 ng/µL) in assay buffer (25 mM Tris-HCl, 137 mM NaCl, 2.7 mM KCl, 1 mM MgCl<sub>2</sub>, 0.5 mM DTT, 0.015% of TritonX-100, pH = 8.0) with 1 µL DMSO, 2.5 µL of ZMML solution (prepared from 12.6 mM stocks in DMSO and diluted with assay buffer, concentration ranges 10-250 µM) and 2.5 µL NAD<sup>+</sup> (prepared from 4 mM stock in assay buffer and diluted in assay buffer; final assay concentration 500 µM) was added to start the reaction. At 0, 1, 3, 5, 10, 20 and 30 minutes the deacylation was quenched with 20 µL of stop solution (50 mM Tris, 100 mM NaCl, 6.7%



(v/v) DMSO, trypsin 1 mg/mL, 8 mM nicotinamide, pH = 8.0). The plate was incubated for 20 min at 37°C and 250 rpm and the fluorescence intensity was measured in a microplate reader ( $\lambda_{\text{Ex}}=390$  nm,  $\lambda_{\text{Em}}=460$  nm, BMG POLARstar Optima, BMG Labtech, Germany). Each experiment was done in duplicate. Deacylation rates were evaluated in relation to theoretical maximal conversion (100% conversion of ZMML obtained by measuring respective concentration of AMC) the data have been fitted to the Michaelis-Menten equation. Origin 9.0 G has been used to determine  $K_m$ ,  $k_{\text{cat}}$  and  $k_{\text{cat}}/K_m$ .

**Kinetic analysis of  $\text{NAD}^+(\text{ZMAL})$  and  $\text{NAD}^+(\text{ZMML})$ .** Deacylation reactions were evaluated by a homogeneous fluorescence based assay. 14  $\mu\text{L}$  of *SmSirt2* solution (104 ng/ $\mu\text{L}$ , final assay concentration 80 ng/ $\mu\text{L}$ ) in assay buffer (25 mM Tris-HCl, 137 mM NaCl, 2.7 mM KCl, 1 mM  $\text{MgCl}_2$ , 0.5 mM DTT, 0,015% of TritonX-100, pH = 8.0) with 1  $\mu\text{L}$  DMSO, 2.5  $\mu\text{L}$  of ZMAL or ZMML solution (prepared respectively from 12.6 mM stocks in DMSO and diluted with assay buffer, saturating assay concentrations) and 2.5  $\mu\text{L}$   $\text{NAD}^+$  (prepared from 4 mM stock in assay buffer and diluted in assay buffer; assay concentration ranges 10-250  $\mu\text{M}$ ). At 0, 1, 3, 5, 10, 20 and 30 minutes the deacylation was quenched with 20  $\mu\text{L}$  of stop solution (50 mM Tris, 100 mM NaCl, 6.7% (v/v) DMSO, trypsin 1 mg/mL, 8 mM nicotinamide, pH = 8.0). The plate was incubated for 20 min at 37°C and 250 rpm and the fluorescence intensity was measured in a microplate reader ( $\lambda_{\text{Ex}}=390$  nm,  $\lambda_{\text{Em}}=460$  nm, BMG POLARstar Optima, BMG Labtech, Germany). Each experiment was done in duplicate. Deacylation rates were evaluated in relation to theoretical maximal conversion (100% conversion of ZMAL or ZMML obtained by measuring respective concentration of AMC) the data have been fitted to the Michaelis-Menten equation. Origin 9.0 G has been used to determine  $K_{m,\text{NAD}^+}$ ,  $k_{\text{cat},\text{NAD}^+}$  and  $k_{\text{cat}}/K_m$ .

***In vitro* Kinetobox screening (*SmSirt2*).** For the screening of the Kinetobox library a homogeneous fluorescence based assay, developed in our group, was used to determine *SmSirt2*<sup>43</sup> activity. OriginPro 9.0 G was employed to determine IC<sub>50</sub> values. Absence of eventual assay interference due to trypsin inhibition was confirmed according published procedures<sup>43</sup> (data not shown), while, to exclude any quenching of the AMC signal, 2.5 μL of a AMC solution (prepared from 12.6 mM stock solution in DMSO and diluted with assay buffer, final assay concentration, 10.5 μM) were used instead of ZMAL in the homogeneous assay. Active compounds were tested for known classes of assay interference compounds with the publicly available online tool “False Positive Remover” ([www.cbligand.org](http://www.cbligand.org)). The only compound flagged to have PAINS characteristics was the approved drug Idebenone but none of our leads or analogues in the optimization campaign.

***In vitro* inhibition of hSirt2.** The activity of potential hits on hSirt2 was measured according published procedures.<sup>56</sup> All compounds were initially tested at 25 μM and, for candidates that showed a hSirt2 inhibition equal or higher than 50% at this concentration, IC<sub>50</sub> values have been measured and determined using OriginPro 9.0 G.

**General chemistry conditions.** Reagents, starting materials and solvents were used without further purification of the purchased form. All reactions were monitored by Thin-layer chromatography (TLC) with Merck precoated silica gel 60 F<sub>254</sub> plates and analysed under UV light (254 nm) using different mobile phases. Microwave assisted reactions were performed using a Discover S-1863 microwave system (CEM GmbH, Germany) and Biotage Initiator (Uppsala, Sweden) high frequency microwave synthesizer working at 2.45 GHz, fitted with magnetic stirrer and sample processor; reaction vessels were Biotage microwave glass vials sealed with applicable cap; temperature was controlled through the internal IR sensor of the microwave apparatus. Synthesized compounds were purified by flash column chromatography with a

Biotage Isolera One automated flash purification system with UV-vis detector. TELOS Flash-LL silica columns 60 M were used as stationary phase with a mobile phase as specified in the following description. Yields were not optimized. Proton ( $^1\text{H}$ ), carbon ( $^{13}\text{C}$ ) and fluorine ( $^{19}\text{F}$ ) spectra were recorded on Bruker Avance III HD spectrometer at 400, 100 and 376 MHz respectively, in reference to the solvents reported in the description. Chemical shifts  $\delta$  are given in parts per million (ppm) and the peak assignment was supported by COSY and HSQC experiments. The purity of compounds **2**, **3**, **6**, **8-12** was determined by HPLC (UV detection at  $\lambda = 210$  nm) and was equal to or higher than 95 % using the following conditions: eluent A,  $\text{H}_2\text{O}$  + 0.05% TFA; eluent B, acetonitrile + 0.05% TFA; linear gradient conditions (0-29 min, linear increase from A = 100% and B = 0% to A = 0% and B = 100%; 29-31 min, B = 100%; 31 min, decrease to B = 10%; 31-40 min B = 10%) with a flow rate of 1 mL/min; analytical column: Phenomenex Synergi 4  $\mu\text{m}$  HYDRO-RP 80  $\text{\AA}$ , 250 mm X 4.6 mm for **3e**, **3f**, **3b**, **3c**, **3g** and **11**; Phenomenex Kinetex 5  $\mu\text{m}$  XB-C18 100  $\text{\AA}$ , 250 mm X 4.6 mm for **3a**, **3d**, **6** and **12**. Low resolution mass spectra of compounds **7**, **14-44** were recorded on an API-TOF Mariner by Perspective Biosystem (Stratford, Texas, USA), samples were injected by a Harvard pump using a flow rate of 5-10  $\mu\text{L}/\text{min}$ , infused in the Electrospray system. High resolution mass spectrometry (HR-MS) with electrospray ionization (ESI) analysis was performed using an Thermo Scientific Exactive mass spectrometer and for low resolution mass spectrometry with electrospray ionization (ESI) analysis was performed using Advion expression CMS spectrometer, with electron ionization (EI) on an Agilent Technologies 6890 N Network GC-MS system. Elemental analysis was used to determine the purity of compounds **7**, **14-44** that was always >95%. Analytical results are within  $\pm 0.40\%$  of the theoretical values. All chemicals were purchased from Sigma Aldrich srl, Milan (Italy) or from TCI Europe NV, Zwijndrecht (Belgium), and were of the highest purity. As a rule, samples prepared for physical and biological

studies were dried in high vacuum over phosphorus pentoxide for 20 h at temperatures ranging from 25 to 40 °C, depending on the sample melting point. General procedures, chemical, physical and spectral data for the syntheses of the final compounds **7**, **14-44** and of all unknown compounds among intermediates **45-47** are described below

**General procedure for the synthesis of 2-Benzyloxycarbonylamino-6-acylamino-hexanoic acid compounds (2a-d and 2g).** To an ice-cold solution of *Z*-lysine (1 equiv, 5.35-6.24 mmol, 1.50-1.75 g) in 5.35-6.24 mL of 1M NaOH (1 equiv, 5.35-6.24 mmol) and water (8 equiv, 42.80-49.92 mmol, 42.80-49.92 mL), 1 equiv of acylchloride (5.35-6.24 mmol, 0.57-1.19 g, 0.55-1.29 mL) in dry THF (1 equiv, 5.35-6.24 mmol, 5.35-6.24 mL) and 5.35-6.24 mL 1M NaOH (1equiv, 5.35-6.24 mmol) were added drop by drop. After 20-45 minutes, the reaction was saturated with NaCl<sub>ss</sub>, cooled below 0 °C and acidified to pH = 1.0 with 2M HCl. The product was extracted with ethyl acetate (four times, 60 mL each) and the combined organic layers extracted with 5% Na<sub>2</sub>CO<sub>3</sub> (four times, 60 mL each). The bicarbonate solution was acidified with 2M HCl to pH = 1.0 and extracted with ethyl acetate (four times, 60 mL each). The combined organic extracts were washed with NaCl<sub>ss</sub>, dried over Na<sub>2</sub>SO<sub>4</sub> followed by evaporation of the solvent.

**2-Benzyloxycarbonylamino-6-butyrylamino-hexanoic acid (2a).** The crude product was purified by flash column chromatography on SiO<sub>2</sub> gel with DCM-methanol (96%-4%). Colorless oil; yield 5% (0.09mmol, 31.7 mg); rf = 0.09 (DCM-methanol 96%-4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 7.34 (s, 5H, *CH* aromatic ring), 5.82 (s, 1H, *NH* amide), 5.72 (s, 1H, *NH* amide), 5.10 (s, 2H, *CH*<sub>2</sub> benzyl), 4.38–4.33 (m, 1H, *CHNH*), 3.33–3.17 (m, 2H, *CH*<sub>2</sub>NH), 2.15 (t, <sup>3</sup>*J*(H,H) = 7.5 Hz, 2H, *COCH*<sub>2</sub>), 1.91–1.73 (m, 2H,*CH*<sub>2</sub>CH), 1.61 (m, 2H, *CH*<sub>2</sub>CH<sub>3</sub>), 1.56–1.47 (m, 2H, *CH*<sub>2</sub>CH<sub>2</sub>), 1.39 (m, 2H, *CH*<sub>2</sub>CH<sub>2</sub>), 0.91 ppm (t, <sup>3</sup>*J*(H,H) = 7.4 Hz, 3H, *CH*<sub>2</sub>CH<sub>3</sub>). MS (ESI), *m/z*: 349.0 [M - H]<sup>-</sup>.

**2-Benzyloxycarbonylamino-6-hexanoylamino-hexanoic acid (2b).** The crude product was directly used for the next step without NMR.

**2-Benzyloxycarbonylamino-6-octanoylamino-hexanoic acid (2c).** Yield 95% (5.93 mmol, 2.41 g) of crude viscous oil; rf = 0.18 (DCM – methanol 96%-4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 7.37–7.28 (m, 5H, *CH* aromatic ring), 6.04 (s, 1H, *NH* amide), 5.67 (d, <sup>3</sup>*J*(H,H) = 8.0 Hz, 1H,

*NH* amide), 5.10 (s, 2H,  $CH_2$  benzyl), 4.40-4.33 (m, 1H,  $CHNH$ ), 3.29–3.18 (m, 2H,  $CH_2NH$ ), 1.94–1.70 (m, 2H,  $CH_2CH$ ), 1.65-1.58 (m, 2H,  $COCH_2$ ), 1.55–1.48 (m, 2H,  $CH_2CH_2$ ), 1.44–1.35 (m, 2H,  $CH_2CH_2$ ), 1.30–1.24 (m, 10H,  $CH_2CH_2$  octanoyl chain), 0.86 ppm (t,  $^3J(H,H) = 6.4$  Hz, 3H,  $CH_2CH_3$ ). MS (ESI),  $m/z$ : 405.1 [M - H]<sup>-</sup>.

**2-Benzyloxycarbonylamino-6-decanoylamino-hexanoic acid (2d).** The crude product was purified by flash column chromatography on SiO<sub>2</sub> gel with DCM-methanol (methanol gradient from 1% to 10%). Colorless oil; yield 39% (2.31 mmol, 1.05 g);  $rf = 0.60$  (DCM-methanol 95%-5%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta = 7.38$ – $7.29$  (m, 5H, *CH* aromatic ring), 5.85 (s, 1H, *NH* amide), 5.66 (s, 1H, *NH* amide), 5.10 (s, 2H,  $CH_2$  benzyl), 4.40-4.36 (m, 1H,  $CHNH$ ), 3.38–3.10 (m, 2H,  $CH_2NH$ ), 2.18 (t,  $^3J(H,H) = 7.6$  Hz, 2H,  $COCH_2$ ), 1.62–1.52 (m, 2H,  $CH_2CH$ ), 1.63-1.58 (m, 2H,  $CH_2CH_2$ ), 1.54–1.50 (m, 2H,  $CH_2CH_2$ ), 1.47–1.35 (m, 2H,  $CH_2CH_2$ ), 1.32–1.19 (m, 12H,  $CH_2CH_2$  decanoyl chain), 0.87 ppm (t,  $^3J(H,H) = 6.9$  Hz, 3H,  $CH_2CH_3$ ). MS (ESI),  $m/z$ : 433.3 [M - H]<sup>-</sup>.

**2-Benzyloxycarbonylamino-6-palmitoylamino-hexanoic acid (2g).** Yield 73% (4.59 mmol, 2.38 g) of crude product as colorless oil;  $rf = 0.13$  (DCM-acetonitrile 60%-40%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta = 7.35$ – $7.28$  (m, 5H, *CH* aromatic ring), 5.94 (s, 1H, *NH* amide), 5.10 (s, 1H, *NH* amide), 4.54 (s, 2H,  $CH_2$  benzyl), 4.41–4.33 (m, 1H,  $CHNH$ ), 3.29–3.17 (m, 2H,  $CH_2NH$ ), 1.92–1.70 (m, 2H,  $CH_2CH$ ), 1.62–1.55 (m, 2H,  $COCH_2$ ), 1.54–1.47 (m, 2H,  $CH_2CH_2$ ), 1.44-1.35 (m, 2H,  $CH_2CH_2$ ), 1.31–1.25 (m, 26H,  $CH_2CH_2$  palmitoyl chain), 0.88 ppm (t,  $^3J(H,H) = 6.9$  Hz, 3H,  $CH_2CH_3$ ). MS (ESI),  $m/z$ : 517.5 [M - H]<sup>-</sup>.

**General procedure for the synthesis of 2-Benzyloxycarbonylamino-6-acylamino-hexanoic acid compounds (2e and 2f).** 1 equiv. of myristic acid (3.50 mmol, 800 mg) or lauric acid (7.49 mmol, 1.5 g) were dissolved in 1 mL of dry DCM. Then 10 equiv of SOCl<sub>2</sub> (35.0-74.9 mmol, 4.16-8.91 g, 2.54-5.43 mL) were added. After 3 hours at 90 °C, the reaction was cooled to r.t. and

SOCl<sub>2</sub> was removed by evaporation. The crude product was then dissolved in 4 mL of DCM dry and 1 equiv of Z-Lysine (3.50-7.49 mmol, 0.981-2.01 g) was added at 0 °C with 2 equiv of 2M NaOH (7.0-14.98 mmol, 280-599 mg, 0.26-0.55mL). The reaction was left stirring for 20-72 h at rt. The reaction was then quenched by adding of 2N HCl to a pH of 2.0 and extracted three times with 20 mL of DCM. The combined organic phase was washed with NaCl<sub>ss</sub> and filtered over Na<sub>2</sub>SO<sub>4</sub>.

**2-Benzoyloxycarbonylamino-6-lauroylamino-hexanoic acid (2e).** The crude product was purified by flash chromatography on SiO<sub>2</sub> gel with DCM-acetonitrile (acetonitrile gradient from 1% to 40%). Yellow oil; yield 12.7% (0.95 mmol, 439 mg); rf = 0.21 (DCM-acetonitrile 55%-45%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 7.37–7.26 (m, 5H, CH aromatic ring), 5.81 (s, 1H, NH amide), 5.73 (d, <sup>3</sup>J(H,H) = 7.8 Hz, 1H, NH amide), 5.09 (s, 2H, CH<sub>2</sub> benzyl), 4.45–4.33 (m, 1H, CHNH), 3.30–3.12 (m, 2H, CH<sub>2</sub>NH), 2.19–2.10 (m, 2H, COCH<sub>2</sub>), 1.93–1.71 (m, 2H, CH<sub>2</sub>CH), 1.63–1.55 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.53–1.46 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.45–1.36 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.31–1.24 (m, 16H, CH<sub>2</sub>CH<sub>2</sub> lauroyl chain), 0.88 ppm (t, <sup>3</sup>J(H,H) = 6.9 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). ESI-MS (-): 461.2 [M - H]<sup>-</sup>.

**2-Benzoyloxycarbonylamino-6-myristoylamino-hexanoic acid (2f).** A mixture of DCM-acetonitrile (acetonitrile gradient from 1% to 80%) was used to purify the crude product by flash chromatography on SiO<sub>2</sub> gel. Colorless oil; yield 13.4% (0.470 mmol, 231 mg); rf = 0.35 (ethyl acetate–methanol 80%-20%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 7.37–7.31 (m, 5H, CH aromatic ring), 5.91 (s, 1H, NH amide), 5.70 (d, <sup>3</sup>J(H,H) = 7.8 Hz, 1H, NH amide), 5.14 (s, 2H, CH<sub>2</sub> benzyl), 4.42-4.36 (m, 1H, CHNH), 3.34–3.14 (m, 2H, CH<sub>2</sub>NH), 2.20 (t, 2H, <sup>3</sup>J(H,H) = 7.2 Hz, COCH<sub>2</sub>), 1.96–1.73 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.65–1.58 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.56–1.51 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.46–1.36 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.32–1.27 (m, 20H, CH<sub>2</sub>CH<sub>2</sub> myristoyl chain), 0.90 ppm (t, <sup>3</sup>J(H,H) = 6.9 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). ESI-MS (-): 489.7 [M - H]<sup>-</sup>.

**General procedure for the synthesis of [5-Acylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyle)pentyl]carbamic acid benzyl esters (3a-g).** 1 equiv of **2a-g** (0.09-4.53 mmol, 0.03-1.72 g) was dissolved in 52 equiv of dry pyridine (4.71-235 mmol, 0.372-18.62 g, 0.38-18.98 mL) and then 2.7 equiv of 7-amino,4-methylcoumarin (0.18-9.06 mmol, 0.03-1.60 g) were added at -15 °C. Then 2.7 equiv (0.24-12.2 mmol, 0.04-1.90 g, 0.02-1.14 mL) of POCl<sub>3</sub> were added by syringe resulting in a red-orange solution. After 40 min-3 h the mixture has been poured into a ten-fold volume of H<sub>2</sub>O/ice and extracted with ethyl acetate (four times, 50 mL each). The combined organic layers were washed with NaCl<sub>ss</sub> (50 mL), 2M HCl (50 mL), NaCl<sub>ss</sub> (50 mL), 5% NaHCO<sub>3</sub> (50 mL) and NaCl<sub>ss</sub> (30 mL). The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent evaporated.

**[5-Butyrylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyle)pentyl] carbamic acid benzyl ester (3a).** The resulting product was purified by flash chromatography on SiO<sub>2</sub> gel with DCM-methanol 96%-4%. White crystal; yield 44% (0.04 mmol, 20.1 mg); rf = 0.22 (DCM-methanol 96%-4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 9.33 (s, 1H, NH amide), 7.67 (s, 1H, CH coumarin), 7.55 (d, <sup>3</sup>J(H,H) = 8.7 Hz, 1H, CH coumarin), 7.49 (d, <sup>3</sup>J(H,H) = 8.7 Hz, 1H, CH coumarin), 7.35 (s, 5H, CH benzyl), 6.18 (s, 1H, NH amide), 5.84 (s, 1H, NH amide), 5.82 (s, 1H, COCH coumarin), 5.13 (s, 2H, CH<sub>2</sub> benzyl), 4.37-4.33 (m, 1H, CHNH), 3.33-3.23 (m, 2H, CH<sub>2</sub>NH), 2.41 (s, 3H, CH<sub>3</sub> coumarin), 2.15 (t, <sup>3</sup>J(H,H) = 7.5 Hz, 2H, COCH<sub>2</sub>), 2.17-1.74 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.65 (m, 2H, CH<sub>2</sub>CH<sub>3</sub>), 1.58 (quint, <sup>3</sup>J(H,H) = 6.1 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.46 (quint, <sup>3</sup>J(H,H) = 6.2 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 0.91 ppm (t, <sup>3</sup>J(H,H) = 7.4 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>): δ = 173.8 (2C, CO amide), 170.9 (CO ester), 161.2 (CO carbamate), 154.0 (CCH<sub>3</sub> coumarin), 152.3 (CO coumarin), 141.5 (CNH coumarin), 136.0 (CCH<sub>2</sub> benzyl), 128.5 (2C, CH benzyl), 128.2 (CH benzyl), 128.0 (2C, CH benzyl), 125.0 (CH coumarin), 116.0 (CH coumarin), 115.7 (CCH coumarin), 113.3 (CH coumarin), 107.2 (COCH coumarin), 67.2 (CH<sub>2</sub> benzyl), 55.3



(CHNH), 38.6 (CH<sub>2</sub>NH), 37.9 (COCH<sub>2</sub>), 31.0 (CH<sub>2</sub>CH), 28.6 (CH<sub>2</sub>CH<sub>2</sub>), 22.0 (CH<sub>2</sub>CH<sub>2</sub>), 19.0 (CCH<sub>3</sub>), 18.5 (CH<sub>2</sub>CH<sub>3</sub>), 14.0 ppm (CH<sub>2</sub>CH<sub>3</sub>). HRMS (ESI): *m/z* calculated for C<sub>28</sub>H<sub>33</sub>N<sub>3</sub>O<sub>6</sub> + H<sup>+</sup> [M + H]<sup>+</sup>: 508.2442. Found: 508.2444. HPLC analysis: retention time = 19.496 min; peak area, 97%.

**[5-Hexanoylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyl)pentyl] carbamic acid benzyl ester (3b)**. Purification of the resulting crude product by flash chromatography on SiO<sub>2</sub> gel with DCM-methanol 96%-4%. Colorless powder; yield 7% (0.30 mmol, 162 mg); *rf* = 0.25 (DCM-methanol 96%-4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 9.35 (s, 1H, NH amide), 7.68 (s, 1H, CH coumarin), 7.54 (d, <sup>3</sup>*J*(H,H) = 8.8 Hz, 1H, CH coumarin), 7.49 (d, <sup>3</sup>*J*(H,H) = 8.6 Hz, 1H, CH coumarin), 7.34 (s, 5H, CH benzyl), 6.18 (s, 1H, CH coumarin), 5.89 (s, 1H, COCH), 5.84 (d, <sup>3</sup>*J*(H,H) = 7.9 Hz, 1H, NH amide), 5.12 (s, 2H, CH<sub>2</sub> benzyl), 4.39–4.33 (m, 1H, CHNH), 3.36–3.19 (m, 2H, CH<sub>2</sub>NH), 2.40 (s, 3H, CCH<sub>3</sub>), 2.17 (t, <sup>3</sup>*J*(H,H) = 7.1 Hz, 2H, COCH<sub>2</sub>), 2.12–1.99 (m, 2H, CH<sub>2</sub>CH), 1.61–1.60 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.60–1.57 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.50–1.43 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.34–1.21 (m, 4H, CH<sub>2</sub>CH<sub>2</sub> and CH<sub>2</sub>CH<sub>3</sub>), 0.87 ppm (t, <sup>3</sup>*J*(H,H) = 6.9 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>): δ = 174.1 (2C, CO amide), 171.1 (CO ester), 161.2 (CO carbamate), 154.0 (CCH<sub>3</sub> coumarin), 152.5 (CO coumarin), 141.5 (CNH coumarin), 136.0 (CCH<sub>2</sub> benzyl), 128.5 (2C, CH benzyl), 128.2 (CH benzyl), 127.9 (2C, CH benzyl), 125.0 (CH coumarin), 115.9 (CH coumarin), 115.8 (CCH coumarin), 113.2 (CH coumarin), 107.2 (COCH coumarin), 67.1 (CH<sub>2</sub> benzyl), 55.3 (CHNH), 38.3 (CH<sub>2</sub>NH), 36.6 (COCH<sub>2</sub>), 31.3 (2C, CH<sub>2</sub>CH<sub>2</sub>), 28.6 (CH<sub>2</sub>CH<sub>2</sub>), 25.4 (2C, CH<sub>2</sub>CH<sub>2</sub>), 22.3 (CH<sub>2</sub>CH<sub>3</sub>), 18.5 (CCH<sub>3</sub>), 13.9 ppm (CH<sub>2</sub>CH<sub>3</sub>). HRMS (ESI): *m/z* calculated for C<sub>30</sub>H<sub>37</sub>N<sub>3</sub>O<sub>6</sub> + H<sup>+</sup> [M + H]<sup>+</sup>: 536.2755. Found: 536.2751. HPLC analysis: retention time = 22.108 min; peak area, 98%.

**[5-Octanoylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyl)pentyl] carbamic acid benzyl ester (3c).** Purification by flash chromatography on SiO<sub>2</sub> gel with DCM-methanol 96%-4%. White crystal; yield 10% (0.39 mmol, 218 mg); *rf* = 0.28 (DCM-methanol 96%-4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  = 9.41 (s, 1H, NH amide), 7.70 (s, 1H, CH coumarin), 7.54 (d, <sup>3</sup>*J*(H,H) = 8.3 Hz, 1H, CH coumarin), 7.49 (d, <sup>3</sup>*J*(H,H) = 8.3 Hz, 1H, CH coumarin), 7.34 (s, 5H, CH benzyl), 6.18 (s, 1H, NH amide), 6.14 (s, 1H, COCH coumarin), 5.87 (s, 1H, NH amide), 5.12 (s, 2H, CH<sub>2</sub> benzyl), 4.41–4.33 (m, 1H, CHNH), 3.34–3.23 (m, 2H, CH<sub>2</sub>NH), 2.40 (s, 3H, CCH<sub>3</sub>), 2.20 (t, <sup>3</sup>*J*(H,H) = 7.1 Hz, 2H, COCH<sub>2</sub>), 2.08–1.73 (m, 4H, CH<sub>2</sub>CH<sub>2</sub> and CH<sub>2</sub>CH), 1.60 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.47 (quint, <sup>3</sup>*J*(H,H) = 7.2 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.25 (s, 8H, CH<sub>2</sub>CH<sub>2</sub> octanoyl chain), 0.85 ppm (t, <sup>3</sup>*J*(H,H) = 6.8 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>):  $\delta$  = 174.2 (2C, CO amide), 171.2 (CO ester), 161.2 (CCH coumarin), 154.0 (CO coumarin), 152.5 (CCH<sub>3</sub> coumarin), 141.5 (CNH coumarin), 136.0 (CO carbamate), 128.5 (2C, CH benzyl), 128.2 (CH benzyl), 128.0 (2C, CH benzyl), 125.0 (CH coumarin), 115.9 (CH coumarin), 115.8 (CCH coumarin), 113.1 (CH coumarin), 107.1 (COCH coumarin), 67.1 (CH<sub>2</sub> benzyl), 55.3 (CHNH), 38.3 (CH<sub>2</sub>NH), 36.5 (COCH<sub>2</sub>), 31.6 (CH<sub>2</sub>CH<sub>2</sub>), 31.4 (CH<sub>2</sub>CH<sub>2</sub>), 29.2 (CH<sub>2</sub>CH), 28.9 (CH<sub>2</sub>CH<sub>2</sub>), 28.6 (CH<sub>2</sub>CH<sub>2</sub>), 25.8 (CH<sub>2</sub>CH<sub>2</sub>), 22.5 (CH<sub>2</sub>CH<sub>2</sub>), 22. (CH<sub>2</sub>CH<sub>3</sub>), 18.5 (CCH<sub>3</sub>), 14.0 ppm (CH<sub>2</sub>CH<sub>3</sub>). HRMS (ESI): *m/z* calculated for C<sub>32</sub>H<sub>41</sub>N<sub>3</sub>O<sub>6</sub> + Na<sup>+</sup> [M + Na]<sup>+</sup>: 563.2888. Found: 586.2885. HPLC analysis: retention time = 24.323 min; peak area, 99%.

**[5-Decanoylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyl)pentyl] carbamic acid benzyl ester (3d).** The crude product was purified by column chromatography on SiO<sub>2</sub> gel with DCM-methanol 96%-4%. White powder; yield 5% (0.12 mmol, 69.40 mg); *rf* = 0.33 (DCM-methanol 96%-4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  = 9.36 (s, 1H, NH amide), 7.69 (s, 1H, CH coumarin), 7.53 (d, <sup>3</sup>*J*(H,H) = 8.8 Hz, 1H, CH coumarin), 7.49 (d, <sup>3</sup>*J*(H,H) = 8.6 Hz, 1H, CH coumarin), 7.34 (s, 5H, CH benzyl), 6.18 (s, 1H, NH amide), 5.93 (s, 1H, COCH coumarin), 5.85

(s, 1H, *NH* amide), 5.13 (s, 2H,  $CH_2$  benzyl), 4.38–4.34 (m, 1H, *CHNH*), 3.33–3.23 (m, 2H,  $CH_2NH$ ), 2.40 (s, 3H,  $CCH_3$ ), 2.18 (t,  $^3J(H,H) = 7.7$  Hz, 2H,  $COCH_2$ ), 2.07–1.87 (m, 2H,  $CH_2CH$ ), 1.81–1.72 (m, 2H,  $CH_2CH_2$ ), 1.60 (quint,  $^3J(H,H) = 7.2$  Hz, 2H,  $CH_2CH_2$ ), 1.47 (quint,  $^3J(H,H) = 7.2$  Hz, 2H,  $CH_2CH_2$ ), 1.23 (s, 12H,  $CH_2CH_2$  deacanoyl chain), 0.86 ppm (t,  $^3J(H,H) = 6.9$  Hz, 3H,  $CH_2CH_3$ ).  $^{13}C$  NMR (101 MHz,  $CDCl_3$ ):  $\delta = 207.1$  (2C, CO amide), 174.1 (CO ester), 161.2 (CO carbamate), 154.0 ( $CCH_3$  coumarin), 152.5 (COCH coumarin), 141.5 (CNH coumarin), 136.0 ( $CCH_2$  benzyl), 128.5 (2C, CH benzyl), 128.2 (CH benzyl), 128.0 (2C, CH benzyl), 125.0 (CH coumarin), 115.9 (CH coumarin), 115.8 ( $CCH$  coumarin), 113.2 (CH coumarin), 107.2 (CH coumarin), 67.1 ( $CH_2$  benzyl), 55.3 (*CHNH*), 38.3 ( $CH_2NH$ ), 36.7 ( $COCH_2$ ), 31.8 ( $CH_2CH_2$ ), 30.9 ( $CH_2CH_2$ ), 29.4 ( $CH_2CH$ ), 29.3 ( $CH_2CH_2$ ), 29.2 ( $CH_2CH_2$ ), 29.2 ( $CH_2CH_2$ ), 28.7 ( $CH_2CH_2$ ), 25.7 ( $CH_2CH_2$ ), 22.6 ( $CH_2CH_2$ ), 22.3 ( $CH_2CH_3$ ), 18.5 ( $CCH_3$ ), 14.1 ppm ( $CH_2CH_3$ ). HRMS (ESI):  $m/z$  calculated for  $C_{34}H_{45}N_3O_6 + Na^+$  [ $M + Na$ ] $^+$ : 591.3201. Found: 614.3201. HPLC analysis: retention time = 26.061 min; peak area, 99%.

**[5-Lauroylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyl)pentyl] carbamic acid benzyl ester (3e).** Purification of crude product by flash column chromatography DCM–methanol (methanol gradient from 1% to 8%). White powder; yield 4% (0.04 mmol, 25 mg);  $rf = 0.30$  (DCM–methanol 96%–4%).  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta = 9.29$  (s, 1H, *NH* amide), 7.69 (s, 1H, *CH* coumarin), 7.54 (s, 1H, *CH* coumarin), 7.50 (s, 1H, *CH* coumarin), 7.35 (s, 5H, *CH* benzyl), 6.19 (s, 1H, *NH* amide), 5.80 (s, 2H, COCH coumarin and *NH* amide), 5.13 (s, 2H,  $CH_2$  benzyl), 4.32–4.28 (m, 1H, *CHNH*), 3.40–3.17 (m, 2H,  $CH_2NH$ ), 2.41 (s, 3H,  $CCH_3$ ), 2.23–1.98 (m, 2H,  $CH_2CH$ ), 1.79–1.70 (m, 4H,  $COCH_2$  and  $CH_2CH_2$ ), 1.61 (m, 4H,  $CH_2CH_2$ ), 1.54–1.38 (m, 2H,  $CH_2CH_2$ ), 1.24 (s, 14H,  $CH_2CH_2$  lauroyl chain), 0.87 ppm (t,  $^3J(H,H) = 6.8$  Hz, 3H,  $CH_2CH_3$ ).  $^{13}C$  NMR (101 MHz,  $CDCl_3$ ):  $\delta = 171.3$  (CO amide), 170.9 (CO amide), 161.2 (CO ester), 154.1 (CO carbamate), 152.3 (CNH coumarin), 141.4 (CO coumarin), 136.0 ( $CCH_3$

coumarin), 128.5 (2C, CH benzyl), 128.2 (CH benzyl), 128.0 (2C, CH benzyl), 125.2 (CH coumarin), 125.1 (CCH coumarin), 120.6 (CCH<sub>2</sub> benzyl), 116.0 (CH coumarin), 113.3 (CH coumarin), 107. (COCH coumarin), 70.04 (CH<sub>2</sub> benzyl), 67.3 (CHNH), 38.4 (CH<sub>2</sub>NH), 37.3 (COCH<sub>2</sub>), 31.9 (CH<sub>2</sub>CH<sub>2</sub>), 31.4 (CH<sub>2</sub>CH<sub>2</sub>), 29.6 (CH<sub>2</sub>CH<sub>2</sub>), 29.5 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.4 (CH<sub>2</sub>CH<sub>2</sub>), 29.3 (CH<sub>2</sub>CH<sub>2</sub>), 28.8 (CH<sub>2</sub>CH<sub>2</sub>), 25.9 (CH<sub>2</sub>CH<sub>2</sub>), 22.6 (CH<sub>2</sub>CH<sub>2</sub>), 22.4 (CH<sub>2</sub>CH<sub>2</sub>), 18.6 (CH<sub>2</sub>CH<sub>3</sub>), 14.8 (CCH<sub>3</sub>), 14.1 ppm (CH<sub>2</sub>CH<sub>3</sub>). HRMS (ESI): *m/z* calculated for C<sub>36</sub>H<sub>49</sub>N<sub>3</sub>O<sub>6</sub> + Na<sup>+</sup> [M + Na]<sup>+</sup>: 642.3514. Found: 642.3510. HPLC analysis: retention time = 25.175 min; peak area, 97%.

**[5-Myristoylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyl)pentyl] carbamic acid benzyl ester (3f).** Purification of crude product by flash column chromatography DCM–methanol (methanol gradient from 1% to 10%). White powder; yield 11% (0.05mmol, 34 mg), *r<sub>f</sub>* = 0.32 (DCM–methanol 96%–4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 9.56 (s, 1H, *NH* amide), 7.75 (s, 1H, *CH* coumarin), 7.57 (d, <sup>3</sup>*J*(H,H) = 8.3Hz, 1H, *CH* coumarin), 7.50 (d, <sup>3</sup>*J*(H,H) = 8.6 Hz, 1H, *CH* coumarin), 7.35 (s, 5H, *CH* benzyl), 6.89 (s, 1H, *NH* amide), 6.20 (s, 1H, COCH), 6.01 (s, 1H, *NH* amide), 5.12 (s, 2H, CH<sub>2</sub> benzyl), 4.45–4.37 (m, 1H, CHNH), 3.41–3.24 (m, 2H, CH<sub>2</sub>NH), 2.4 (s, 3H, CCH<sub>3</sub>), 2.30 (t, <sup>3</sup>*J*(H,H) = 7.3Hz, 2H, COCH<sub>2</sub>), 2.12–1.64 (m, 2H, CH<sub>2</sub>CH), 1.67–1.64 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.62–1.58 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.54–1.47 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.33–1.23 (m, 20H, CH<sub>2</sub>CH<sub>2</sub> myristoyl chain), 0.89 ppm (t, <sup>3</sup>*J*(H,H) = 6.9 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR (101 MHz, (CD<sub>3</sub>)<sub>2</sub>SO): δ = 172.4 (CO amide), 160.5 (CO amide), 156.6 (CO ester), 154.1 (CO carbamate), 153. (CCH<sub>3</sub> coumarin), 142.7 (COCH coumarin), 128.8 (CNH coumarin), 128.3 (CCH<sub>2</sub> benzyl), 128.2 (2C, CH benzyl), 126.4 (CH benzyl), 115.7 (2C, CH benzyl), 115.5 (CH coumarin), 112.8 (CH coumarin), 106.1 (CCH coumarin), 65.9 (CH coumarin), 56.0 (COCH coumarin), 38.5 (CH<sub>2</sub> benzyl), 35.9 (CHNH), 31.8 (CH<sub>2</sub>NH), 29.5 (COCH<sub>2</sub>), 29.5 (CH<sub>2</sub>CH), 29.4 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.3 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.2 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.2 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.1 (2C, CH<sub>2</sub>CH<sub>2</sub>),

25.8 (CH<sub>2</sub>CH<sub>2</sub>), 23.4 (CH<sub>2</sub>CH<sub>2</sub>), 22.6 (CH<sub>2</sub>CH<sub>3</sub>), 18.4 (CCH<sub>3</sub> coumarin), 14.4 ppm (CH<sub>2</sub>CH<sub>3</sub>). MS (ESI), *m/z*: 646.7 [M - H]<sup>-</sup>. HPLC analysis: retention time = 30.424 min; peak area, 95%.

**[5-Palmitoylamino-1-(4-methyl-2-oxo-2H-chromen-7-ylcarbamoyl)pentyl] carbamic acid benzyl ester (3g)**. Purification of crude product by flash column chromatography DCM–methanol (methanol gradient from 1% to 8%). White powder; yield 10% (0.29 mmol, 200mg); *rf* = 0.60 (DCM–methanol 96%–4%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 9.30 (s, 1H, NH amide), 7.68 (s, 1H, CH coumarin), 7.54 (d, <sup>3</sup>*J*(H,H) = 8.8 Hz, 1H, CH coumarin), 7.49 (d, <sup>3</sup>*J* = 8.6 Hz, 1H, CH coumarin), 7.35 (s, 5H, CH benzyl), 6.19 (s, 1H, NH amide), 5.79 (d, <sup>3</sup>*J*(H,H) = 8.8 Hz, 1H, NH amide), 5.75 (s, 1H, COCH), 5.13 (s, 2H, CH<sub>2</sub> benzyl), 4.38–4.30 (m, 1H, CHNH), 3.37–3.19 (m, 2H, CH<sub>2</sub>NH), 2.41 (s, 3H, CCH<sub>3</sub>), 2.16 (t, <sup>3</sup>*J*(H,H) = 7.1 Hz, 2H, COCH<sub>2</sub>), 2.08–1.72 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.64–1.60 (m, 2H, CH<sub>2</sub>CH), 1.59–1.55 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.50–1.42 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.31–1.19 (m, 24H, CH<sub>2</sub>CH<sub>2</sub> palmitoyl chain), 0.88 ppm (t, <sup>3</sup>*J*(H,H) = 6.9 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>): δ = 174.0 (CO amide), 174.0 (CO amide), 171.0 (CO ester), 161.2 (CO carbamate), 154.0 (CCH<sub>3</sub> coumarin), 152.3 (COCH coumarin), 141.5 (CNH coumarin), 136.0 (CCH<sub>2</sub> benzyl), 128.5 (2C, CH benzyl), 128.2 (CH benzyl), 127.9 (2C, CH benzyl), 125.0 (CH coumarin), 115.9 (CH coumarin), 115.7 (CCH coumarin), 113.2 (CH coumarin), 107.2 (COCH), 67.2 (CH<sub>2</sub> benzyl), 65.8 (CHNH), 38.1 (COCH<sub>2</sub>), 36.7 (CH<sub>2</sub>CH<sub>2</sub>), 31.9 (CH<sub>2</sub>CH<sub>2</sub>), 31.2 (CH<sub>2</sub>CH), 29.6 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.6 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.6 (CH<sub>2</sub>CH<sub>2</sub>), 29.5 (2C, CH<sub>2</sub>CH<sub>2</sub>), 29.3 (CH<sub>2</sub>CH<sub>2</sub>), 29.2 (CH<sub>2</sub>CH<sub>2</sub>), 28.6 (CH<sub>2</sub>CH<sub>2</sub>), 25.7 (CH<sub>2</sub>CH<sub>2</sub>), 22.6 (CH<sub>2</sub>CH<sub>2</sub>), 22.1 (CH<sub>2</sub>CH<sub>2</sub>), 18.5 (CH<sub>2</sub>CH<sub>3</sub>), 15.2 (CCH<sub>3</sub> coumarin), 14.1 ppm (CH<sub>2</sub>CH<sub>3</sub>). MS (ESI), *m/z*: 674.9 [M - H]<sup>-</sup>. HPLC analysis: retention time = 26.635 min; peak area, 98%.

**Synthesis of *N*-(4-(5-((1-(3-fluoro-2-methylbenzyl)piperidin-3-yl)methyl)-1,2,4-oxadiazol-3-yl)phenyl)acetamide (6)**. 1 equiv of **8** (0.51 mmol, 150 mg), 1 equiv of **10a** (0.51 mmol, 98.8

mg) and 3.7 equiv of  $K_2CO_3$  (1.89 mmol, 261.3mg) were mixed in 5.00 mL of pyridine and refluxed for eight hours. The reaction mixture was left for 72 h at r.t., and then was diluted with 15 mL of ethyl acetate, washed with  $H_2O$  (two times, 10 mL each) and  $NaCl_{ss}$  (10 mL) and dried over  $Na_2SO_4$ . The solvent was evaporated and the crude product purified by flash column chromatography on  $SiO_2$  gel with DCM–methanol (98%–2%). Yellow oil; yield 5% (0.03mmol, 12mg);  $rf= 0.28$  (DCM-methanol 95%–5%).  $^1H$  NMR (400 MHz,  $d_6$ -DMSO, 50°C):  $\delta = 10.12$  (s, 1H, *NH* anilide), 7.88 (d,  $^3J(H,H) = 8.8$  Hz, 2H, *CH* anilide), 7.75 (d,  $^3J(H,H) = 8.8$  Hz, 2H, *CH* anilide), 7.15–6.97 (m, 3H, *CH* benzyl), 3.44 (s, 2H,  $CH_2$  benzyl), 3.00–2.92 (m, 2H,  $CHCH_2$ ), 2.77–2.74 (m, 1H, *CH* piperidine), 2.63–2.56 (m, 1H, *CH* piperidine), 2.23 (s, 3H,  $CCH_3$  benzyl), 2.15–2.13 (m, 1H, *CH* piperidine), 2.12–2.11 (m, 1H, *CH* piperidine), 2.09 (s, 3H,  $CH_3$  amide), 2.00–1.92 (m, 1H, *CH* piperidine), 1.76–1.68 (m, 1H, *CH* piperidine), 1.67–1.61 (m, 1H, *CH* piperidine), 1.53–1.42 (m, 1H, *CH* piperidine), 1.21–1.11 ppm (m, 1H, *CH* piperidine).  $^{13}C$  NMR (101 MHz,  $(CD_3)_2SO$ , 50°C):  $\delta = 179.4$  (CONH amide), 169.1 (CN oxadiazole), 167.6 (CO oxadiazole), 141.2 (d,  $^1J(C,F) = 264.1$  Hz, CF), 128.1 (*CH* anilide), 126.7 (*CH* benzyl), 125.8 (*CH* benzyl), 124.3 (2C, CCH anilide and  $CCH_2$  benzyl), 124.1 (CNH anilide), 121.0 ( $CCH_3$  benzyl), 120.6 (CCH anilide), 119.5 (2C, *CH* anilide), 113.7 (*CH* benzyl), 60.5 ( $NCH_2$  benzyl), 58.7 ( $NCH_2$  piperidine), 53.8 ( $NCH_2$  piperidine), 34.50 ( $CH_2CH$  piperidine), 34.47 ( $CHCH_2$  piperidine) 30.3 ( $COCH_3$ ), 30.1 ( $CH_2CH$  piperidine), 24.5 ( $CH_2CH_2$  piperidine), 10.3 ppm ( $CCH_3$  benzyl).  $^{19}F$  (376 MHz,  $(CD_3)_2SO$ , 50°C):  $\delta = -117.73$  ppm (s, CF). HRMS (ESI):  $m/z$  calculated for  $C_{24}H_{27}FN_4O_2 + H^+$  [M + H] $^+$ : 423.2191.

Found: 423.2195. HPLC analysis: retention time = 16.559 min; peak area, 99%.

**Synthesis of ethyl 2-(1-(3-fluoro-2-methylbenzyl)piperidin-3-yl)acetate (8).** 2-(piperidin-3-yl) acetic acid ethyl ester (1 equiv, 5.84 mmol, 1 g) was dissolved in 2.5 mL of toluene, followed by the addition of 2.4 equiv of triethylamine (11.7 mmol, 1.62 mL) and 1 equiv of 3-fluoro-2-

methylbenzyl bromide (5.84 mmol, 1.18 g) at 0 °C. After 10 min, the mixture was warmed up to room temperature. After 18 h the precipitate was removed by filtration, washed with cyclohexane and the filtrate has been concentrated. The crude product was dried by the use of a vacuum pump for 2-5 h and then purified by flash column chromatography on SiO<sub>2</sub> gel with DCM-methanol (99%-1%). Colorless oil; yield 85-90% (2.45-2.89 mmol, 723-767 mg); rf = 0.67 (DCM-methanol 95%-5%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 7.10-7.0 (m, 2H, CH benzyl), 6.94-6.89 (m, 1H, CH benzyl), 4.15-4.03 (m, 2H, CH<sub>2</sub>CH<sub>3</sub>), 3.41 (s, 2H, CH<sub>2</sub> benzyl), 2.75-2.59 (m, 2H, CH<sub>2</sub>COO), 2.25 (s, 3H, CH<sub>3</sub> benzyl), 2.24-2.22 (m, 1H, CH piperidine), 2.20-2.16 (m, 1H, CH piperidine), 2.13-1.99 (m, 1H, CH piperidine), 1.89-1.80 (m, 1H, CH piperidine), 1.76-1.74 (m, 1H, CH piperidine), 1.67-1.59 (m, 1H, CH piperidine), 1.59-1.50 (m, 2H, CH piperidine), 1.21 (t, <sup>3</sup>J(H,H) = 7.1 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>), 1.09-1.00 ppm (m, 1H, CH piperidine). MS (ESI), *m/z*: 294.32 [M + H]<sup>+</sup>.

**General procedure for the synthesis of *N*-(4-cyanophenyl)acylamide (9b-c).** 1 equiv of 4-aminobenzonitrile (4.23 mmol, 500 mg) was dissolved in 15 mL of dry DCM at 0 °C under nitrogen atmosphere, followed by addition of 7 equiv of triethylamine and 4 equiv of acyl chloride. After 2 h at r.t. the reaction was quenched with water and extracted (three times, 15 mL of water each). Then the organic phase was washed with 2M HCl (once, 15 mL), NaCl<sub>ss</sub> (once, 15 mL), 5% Na<sub>2</sub>CO<sub>3</sub> (once, 15 mL), NaCl<sub>ss</sub> (once, 15 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated to dryness and the crude product purified by flash column chromatography on SiO<sub>2</sub> gel with DCM-methanol (98%-2%).

***N*-(4-cyanophenyl)octanamide (9b).** White solid; yield 62-65% (2.60-2.73 mmol, 635-666 mg); rf = 0.47 (DCM-methanol 98%-2%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 7.67 (d, <sup>3</sup>J(H,H) = 8.8 Hz, 2H, CH aromatic ring), 7.61 (d, <sup>3</sup>J(H,H) = 8.8 Hz, 2H, CH aromatic ring), 7.38 (s, 1H, NH amide), 2.39 (t, <sup>3</sup>J(H,H) = 7.2 Hz, 2H, COCH<sub>2</sub>), 1.73 (quint, <sup>3</sup>J(H,H) = 7.2 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>),

1.42–1.23 (m, 8H, CH<sub>2</sub>CH<sub>2</sub>), 0.88 ppm (t, <sup>3</sup>J(H,H) = 7.0 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). MS (ESI), *m/z*: 243.3 [M - H]<sup>-</sup>.

***N*-(4-cyanophenyl)decanamide (9c)**. Yellow solid; yield 71% (5.98 mmol, 1.63 g); *rf* = 0.63 (DCM-methanol 99%-1%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 7.67 (d, <sup>3</sup>J(H,H) = 8.9 Hz, 2H, CH aromatic ring), 7.61 (d, <sup>3</sup>J(H,H) = 8.9 Hz, 2H, CH aromatic ring), 7.35 (s, 1H, NH amide), 2.39 (t, <sup>3</sup>J(H,H) = 7.6 Hz, 2H, COCH<sub>2</sub>), 1.73 (quint, <sup>3</sup>J(H,H) = 7.6 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.39-1.27 (m, 12H, CH<sub>2</sub>CH<sub>2</sub>), 0.88 ppm (t, <sup>3</sup>J(H,H) = 6.9 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). MS (ESI), *m/z*: 271.4 [M - H]<sup>-</sup>.

**General procedure for the synthesis of (Z)-*N*-(4-(*N*'-hydroxycarbamimidoyl)-phenyl)acylamide (10a-c)**. 1 equiv of *N*-(4-cyanophenyl)acylamide (3.74-5.98 mmol, 0.60-1.63 g), 3.7 equiv of NH<sub>2</sub>OH·HCl (13.84-22.14 mmol, 0.96-1.50 g) and 1.7 equiv of Na<sub>2</sub>CO<sub>3</sub> (6.36-10.17 mmol, 0.67-1.07 g) were dissolved in a mixture of water and EtOH. After stirring for 6-8 h at reflux, the reaction was cooled on ice and a yellow-orange precipitate formed. The precipitate was collected by filtration and dried under vacuum.

***N*-(4-(*N*'-hydroxycarbamimidoyl)phenyl)acetamide (10a)**. The reaction was performed in a mixture of 5 mL ethanol and 20 mL water. **3a** was obtained as white crystal; yield 86% (4.03 mmol, 778 mg); *rf* = 0.83 (DCM-methanol 80%-20%). <sup>1</sup>H NMR (400 MHz, *d*<sub>6</sub>-DMSO): δ = 10.04 (s, 1H, NH amide), 9.53 (s, 1H, NOH), 7.60 (d, <sup>3</sup>J(H,H) = 9.1 Hz, 2H, CH aromatic ring), 7.56 (d, <sup>3</sup>J(H,H) = 9.1 Hz, 2H, CH aromatic ring), 5.76 (s, 2H, CNH<sub>2</sub>), 2.05 ppm (s, 3H, COCH<sub>3</sub>). MS (ESI), *m/z*: 194.2 [M + H]<sup>+</sup>.

**(Z)-*N*-(4-(*N*'-hydroxycarbamimidoyl)phenyl)octanamide (10b)**. A mixture of 15 mL water and 45 mL ethanol was used for the reaction. **3b** is a white crystal; yield 77% (4.04 mmol, 1.12 g); *rf* = 0.07 (DCM-methanol 95%-5%). <sup>1</sup>H NMR (400 MHz, *d*<sub>6</sub>-DMSO): δ = 10.01 (s, 1H, NH amide), 9.54 (s, 1H, NOH), 7.86-7.23 (m, 4H, CH aromatic ring), 5.78 (s, 2H, CNH<sub>2</sub>), 2.31 (t,



$^3J(\text{H,H}) = 7.4$  Hz, 2H,  $\text{COCH}_2$ ), 1.65- 1.52 (m, 2H,  $\text{CH}_2\text{CH}_2$ ), 1.36-1.20 (m, 8H,  $\text{CH}_2\text{CH}_2$ ), 0.86 ppm (t,  $^3J(\text{H,H}) = 6.5$  Hz, 3H,  $\text{CH}_2\text{CH}_3$ ). MS (ESI),  $m/z$ : 278.2  $[\text{M} + \text{H}]^+$ .

**(Z)-N-(4-(N'-hydroxycarbamimidoyl) phenyl)decanamide (10c).** 20 mL of water and 45 mL of ethanol were used in this case as reaction solvent. **3c** is a white crystal; yield 87% (5.17 mmol, 1.58 g);  $r_f = 0.07$  (DCM-methanol 98%-2%).  $^1\text{H}$  NMR (400 MHz,  $d_6$ -DMSO):  $\delta = 9.95$  (s, 1H, NH amide), 9.52 (s, 1H, NOH), 7.85-7.26 (m, 4H, CH aromatic ring), 5.74 (s, 2H,  $\text{CNH}_2$ ), 2.30 (t,  $^3J(\text{H,H}) = 7.4$  Hz, 2H,  $\text{COCH}_2$ ), 1.62-1.55 (quint,  $^3J(\text{H,H}) = 7.3$  Hz, 2H,  $\text{CH}_2\text{CH}_2$ ), 1.33-1.19 (m, 12H,  $\text{CH}_2\text{CH}_2$ ), 0.86 ppm (t,  $^3J(\text{H,H}) = 7.1$  Hz, 3H,  $\text{CH}_2\text{CH}_3$ ). MS (ESI),  $m/z$ : 306.4  $[\text{M} + \text{H}]^+$ .

**Synthesis of N-(4-(5-((1-(3-fluoro-2-methylbenzyl)piperidin-3-yl)methyl)-1,2,4-oxadiazol-3-yl)phenyl)octanamide (11).** 1 equiv of **8** (0.82 mmol, 240 mg), 2 equiv of **10b** (1.64 mmol, 453.5 mg) and 4 equiv of  $\text{K}_2\text{CO}_3$  (3.27 mmol, 452 mg) in 9 mL of pyridine were mixed and the reaction was performed with microwaves (10 minutes, 180 °C, 300 W). Then the mixture was refluxed at 160 °C for 43 h, and stirred at r.t. for 12 h. After that the pyridine was removed and the reaction mixture dissolved in 40 mL of ethyl acetate, extracted with water (three times, 15 mL each), washed with  $\text{NaCl}_{\text{ss}}$  (once, 15 mL) and dried over  $\text{Na}_2\text{SO}_4$ . The solvent was evaporated and the crude product purified by flash column chromatography on  $\text{SiO}_2$  gel with DCM-methanol (98%-2%) and with DCM-methanol (95%-5%) obtaining **11** as colorless oil. Yield 28.9% (0.02 mmol, 12 mg);  $r_f = 0.36$  (DCM-methanol 95%-5%).  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ , 50°C):  $\delta = 10.06$  (s, 1H, NH anilide), 7.93 (d,  $^3J(\text{H,H}) = 8.6$  Hz, 2H, CH anilide), 7.75 (d,  $^3J(\text{H,H}) = 8.6$  Hz, 2H, CH anilide), 7.24-6.97 (m, 3H, CH benzyl), 3.80-3.64 (m, 2H,  $\text{CH}_2\text{CH}$ ), 3.13-3.02 (m, 1H, CH piperidine), 3.00-2.97 (m, 1H, CH piperidine), 2.95-2.92 (m, 1H, CH piperidine), 2.44-2.42 (m, 2H,  $\text{COCH}_2$ ), 2.40-2.37 (m, 1H, CH piperidine), 2.30 (s, 3H,  $\text{CCH}_3$  benzyl), 2.29-2.22 (m, 1H, CH piperidine), 2.18 (s, 2H,  $\text{CH}_2$  benzyl), 1.89-1.84 (m, 1H, CH piperidine), 1.82-1.79

(m, 1H, CH piperidine), 1.78-1.76 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.73-1.68 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.67-1.60 (m, 1H, CH piperidine), 1.40-1.19 (m, 7H, CH piperidine and CH<sub>2</sub>CH<sub>2</sub> octanoyl chain), 0.92 ppm (t, <sup>3</sup>J(H,H) = 6.4 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR (400 MHz, CD<sub>3</sub>OD, 50°C): δ = 178.7 (CONH amide), 173.6 (CN oxadiazole), 167.6 (CO oxadiazole), 141.5 (CNH anilide), 127.6 (2C, CH anilide), 126.3 (CH benzyl), 126.2 (2C, CCH anilide and CCH<sub>2</sub> benzyl), 126.0 (CH benzyl), 120.6 (d, <sup>1</sup>J(C-F) = 212.6 Hz, CF), 119.5 (2C, CH anilide), 114.2 (CCH<sub>3</sub> benzyl), 113.9 (CH benzyl), 100.0 (CH<sub>2</sub> benzyl), 59.5 (CH<sub>2</sub>CH), 57.7 (NCH<sub>2</sub> piperidine), 53.5 (NCH<sub>2</sub> piperidine), 36.7 (COCH<sub>2</sub>), 31.5 (CH<sub>2</sub>CH<sub>2</sub>), 29.9 (CHCH<sub>2</sub>), 29.1 (CH<sub>2</sub>CH piperidine), 28.9 (CH<sub>2</sub>CH<sub>2</sub>), 28.8 (CH<sub>2</sub>CH<sub>2</sub>), 25.4 (CH<sub>2</sub>CH<sub>2</sub>), 23.6 (CH<sub>2</sub>CH<sub>2</sub> piperidine), 22.3 (CH<sub>2</sub>CH<sub>3</sub>), 13.0 (CH<sub>2</sub>CH<sub>3</sub>), 9.3 ppm (CCH<sub>3</sub>). <sup>19</sup>F (376 MHz, CDCl<sub>3</sub>, 50°C): δ = -117.40 ppm (s, CF). HRMS (ESI): *m/z* calculated for C<sub>30</sub>H<sub>39</sub>FN<sub>4</sub>O<sub>2</sub> + H<sup>+</sup> [M + H]<sup>+</sup>: 507.3130. Found: 507.3129. HPLC analysis: retention time = 22.798 min; peak area, 96%.

**Synthesis of *N*-(4-(5-((1-(3-fluoro-2-methylbenzyl)piperidin-3-yl)methyl)-1,2,4-oxadiazol-3-yl)phenyl)decanamide (12).** 1 equiv of **8** (3.41 mmol, 1 g) was dissolved in 23 mL of ethanol followed by drop by drop addition of 4 equiv (13.63 mmol, 13.6 mL) of 1M LiOH solution. The reaction mixture was stirred for 20 h at r.t. (TLC-control), acidified with 2M HCl solution with consequent removal of water by evaporation. Under N<sub>2</sub> atmosphere, the previously formed carboxylic acid was dissolved in 200 μL of *N,N*-dimethylformamide and 2.5 mL of dry DCM. 5 equiv of SOCl<sub>2</sub> (9.31 mmol, 1.1 g) were added drop by drop at -15 °C. The reaction mixture was stirred for 3 h (TLC-control) and then SOCl<sub>2</sub> was removed by stream of N<sub>2</sub>. 1.3 equiv of **10c** (2.42 mmol, 739 mg), solubilized in 3mL of *N,N*-dimethylformamide and 5 mL of dry DCM, were added to the acyl chloride, followed by 3 equiv of triethylamine (5.58 mmol, 0.77 mL). After 22 h, the solvent was evaporated and 5 mL of *N,N*-dimethylformamide were added. The mixture was heated at 150 °C for 3 h, basified with 2 M KOH and extracted with ethyl acetate

(three times, 10 mL each), washed with NaCl<sub>ss</sub> (once, 10 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. The product was purified by flash chromatography DCM-methanol (methanol gradient from 2% to 25%) followed by preparative HPLC obtaining **12** as white powder. Yield 2.3% (0.08 mmol, 41.8 mg); rf = 0.33 (DCM-methanol 98%-2%). <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD, 50°C): δ = 10.08 (s, 1H, NH amide), 7.90 (d, <sup>3</sup>J(H,H) = 8.3 Hz, 2H, CH anilide), 7.76 (d, <sup>3</sup>J(H,H) = 8.4 Hz, 2H, CH anilide), 7.36-7.20 (m, 3H, CH benzyl), 4.53-4.40 (m, 2H, CHCH<sub>2</sub>), 3.87-3.79 (m, 1H, CH piperidine), 3.65-3.55 (m, 1H, CH piperidine), 3.16-3.12 (m, 1H, CH piperidine), 3.10-3.09 (m, 1H, CH piperidine), 3.03-3.01 (m, 1H, CH piperidine), 2.98-2.93 (m, 1H, CH piperidine), 2.49-2.43 (m, 2H, COCH<sub>2</sub>), 2.40 (s, 3H, CCH<sub>3</sub> benzyl), 2.09-1.97 (m, 3H, CH piperidine and CH<sub>2</sub> benzyl), 1.95-1.85 (m, 1H, CH piperidine), 1.77-1.66 (quint, <sup>3</sup>J(H,H) = 7.8 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 1.52-1.32 (m, 13H, CH piperidine and CH<sub>2</sub>CH<sub>2</sub> decanoyl chain), 0.91 ppm (t, <sup>3</sup>J(H,H) = 6.2 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR (400 MHz, CD<sub>3</sub>OD, 50°C): δ = 177.5 (CONH amide), 173.7 (CN oxadiazole), 167.6 (CO oxadiazole), 162.8 (CH benzyl), 141.6 (CNH anilide), 127.7 (CCH<sub>3</sub> benzyl), 127.6 (CH benzyl), 127.5 (2C, CCH anilide and CCH<sub>2</sub> benzyl), 120.6 (d, <sup>1</sup>J(C-F) = 188.1 Hz, CF), 119. (CH<sub>2</sub> benzyl), 119.5 (2C, CH anilide), 116.6 (2C, CH anilide), 116.3 (CH benzyl), 59.3 (CHCH<sub>2</sub>), 55.7 (NCH<sub>2</sub> piperidine), 52.7 (NCH<sub>2</sub> piperidine), 36.8 (COCH<sub>2</sub>), 32.4 (CH<sub>2</sub>CH piperidine), 31.6 (CHCH<sub>2</sub> piperidine), 29.3 (CH<sub>2</sub>CH<sub>2</sub>), 29.2 (CH<sub>2</sub>CH<sub>2</sub>), 29.0 (CH<sub>2</sub>CH<sub>2</sub> piperidine), 28.9 (CH<sub>2</sub>CH<sub>2</sub> piperidine), 27.8 (CH<sub>2</sub>CH<sub>2</sub>), 25.4 (CH<sub>2</sub>CH<sub>2</sub>), 22.3 (CH<sub>2</sub>CH<sub>2</sub>), 22.2 (CH<sub>2</sub>CH<sub>3</sub>), 13.0 (CH<sub>2</sub>CH<sub>3</sub>), 9.8 ppm (CCH<sub>3</sub>). <sup>19</sup>F (376 MHz, (CDCl<sub>3</sub>, 50°C): δ = -112.65 ppm (s, CF). HRMS (ESI): *m/z* calculated for C<sub>32</sub>H<sub>43</sub>FN<sub>4</sub>O<sub>2</sub> + H<sup>+</sup> [M + H]<sup>+</sup>: 535.3443. Found: 535.3441. HPLC analysis: retention time = 13.378 min; peak area, 98%.

**General Procedure for the Synthesis of the N<sup>7</sup>-Substituted pyrimido[4,5-d]pyrimidine-2,4,7-triamines 7, 14-44.** 2-Substituted-4-aminopyrimidin-5-carbonitriles **47** (1 equiv, 0.83 mmol) were stirred with 0.73 M free base guanidine solution in dry 2-methoxyethanol (3.5 equiv, 2.90

mmol, 3.97 mL) at 150 °C for 1.5-4.5 h. After the completion of the reaction, the mixture was concentrated, quenched with water and extracted with ethyl acetate. The organic phase was dried over anhydrous sodium sulfate, filtered and evaporated giving a crude product purified by silica gel column chromatography eluting with a mixture of chloroform/methanol/ammonia and then triturated with a mixture of petroleum ether/diethyl ether to afford the final compounds **7**, **14-44** as a white powder.

***N*<sup>7</sup>-(1-(4-(4-methoxyphenoxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (**7**)**. Recryst. Solvent: acetonitrile. Yield: 60.0 %. mp: 181-184 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.50-1.52 (d, 3H, CHCH<sub>3</sub>), 2.82 (br s, 3H, NCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 6.28 (br s, 1H, CHCH<sub>3</sub>), 6.49 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.87-6.90 (d, 2H, CH benzene ring), 6.94-7.00 (m, 4H, CH benzene rings), 7.25-7.27 (d, 2H, CH benzene rings), 7.41 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.95 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, DMSO) δ 16.8, 29.2, 51.3, 55.9, 96.4, 115.5 (2C), 117.6 (2C), 121.1 (2C), 128.8 (2C), 136.3, 149.9, 156.0, 156.7, 157.3, 163.3, 163.4, 165.9, 166.0. MS (ESI), *m/z*: 418 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>22</sub>H<sub>23</sub>N<sub>7</sub>O<sub>2</sub>: C 63.30, H 5.55, N 23.49. Found: C 63.38, H 5.57, N 23.41.

***N*<sup>7</sup>-(1-benzylpiperidin-4-yl)pyrimido[4,5-d]pyrimidine-2,4,7-triamine (**14**)**. Recryst. Solvent: acetonitrile/methanol. Yield: 75.3 %. mp: 205-208 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.46-1.54 (m, 2H, 2 x CH piperidine ring), 1.82-1.85 (m, 2H, 2 x CH piperidine ring), 1.99-2.04 (m, 2H, 2 x CH piperidine ring), 2.79-2.81 (m, 2H, 2 x CH piperidine ring), 3.46 (s, 2H, NCH<sub>2</sub>Ph), 3.74 (br m, 1H, NHC<sub>4</sub>-H-piperidine ring), 6.40 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 7.18-7.35 (m, 8H, CH benzene ring, NH and C<sub>4</sub>-NH<sub>2</sub>), 8.84 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, DMSO) δ 31.8 (2C), 48.2, 52.8 (2C), 62.7, 96.2, 127.3, 128.6 (2C), 129.2 (2C), 139.2, 157.0, 163.3, 163.4, 165.7, 166.2. MS (ESI), *m/z*: 351 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>18</sub>H<sub>22</sub>N<sub>8</sub>: C 61.70, H 6.33, N 31.98. Found: C 61.80, H 6.35, N 31.85.

**7-(2-methylpiperidin-1-yl)pyrimido[4,5-d]pyrimidine-2,4-diamine (15).** Recryst. Solvent: methanol. Yield: 52.8 %. mp: > 300 °C. <sup>1</sup>H-NMR (400 MHz; DMSO) δ ppm: 1.12-1.14 (d, 3H, CHCH<sub>3</sub>), 1.36 (m, 1H, CH piperidine ring), 1.58-1.70 (m, 5H, 5 x CH piperidine ring), 2.85-2.92 (t, 1H, CH piperidine ring), 4.66-4.69 (m, 1H, CH piperidine ring), 5.10 (br s, 1H, CH-CH<sub>3</sub>-piperidine ring), 6.43 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 7.34 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.90 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, DMSO) δ 15.4, 19.1, 25.9, 30.3, 38.3, 45.4, 96.1, 156.7, 162.6, 163.3, 165.9, 166.1. MS (ESI), *m/z*: 260 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>12</sub>H<sub>17</sub>N<sub>7</sub>: C 55.58, H 6.61, N 37.81. Found: C 55.68, H 6.63, N 37.69.

**7-(3,4-dihydroisoquinolin-2(1H)-yl)pyrimido[4,5-d]pyrimidine-2,4-diamine (16).** Recryst. Solvent: methanol. Yield: 70.6 %. mp: 278-280 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.85-2.88 (m, 2H, CH piperidine ring), 4.03 (m, 2H, CH piperidine ring), 4.92 (s, 2H, CH piperidine), 6.51 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 7.18-7.25 (m, 4H, CH isoquinoline ring), 7.43 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.97 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 28.6, 41.6, 46.1, 96.5, 126.5, 126.7, 126.8, 129.0, 134.7, 135.4, 156.8, 162.8, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 294 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>15</sub>H<sub>15</sub>N<sub>7</sub>: C 61.42, H 5.15, N 33.43. Found: C 61.53, H 5.17, N 33.30.

**7-(4-phenylpiperazin-1-yl)pyrimido[4,5-d]pyrimidine-2,4-diamine (17).** Recryst. Solvent: methanol. Yield: 62.8 %. mp: > 300 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 3.19 (m, 4H, 2 x CH<sub>2</sub> piperazine ring), 3.95 (m, 4H, 2 x CH<sub>2</sub> piperazine ring), 6.52 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.79-6.83 (t, 1H, CH benzene ring), 6.99-7.01 (m, 2H, CH benzene ring), 7.22-7.26 (m, 2H, CH benzene ring), 7.43 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 43.7 (2C), 48.9 (2C), 96.6, 116.3 (2C), 119.6, 129.4 (2C), 151.5, 156.8, 162.8, 163.4,

166.0, 166.1. MS (ESI),  $m/z$ : 323  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{16}H_{18}N_8$ : C 59.61, H 5.63, N 34.76. Found: C 59.72, H 5.65, N 34.63.

***N*<sup>7</sup>-(4-methoxybenzyl)pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (18).** Recryst. Solvent: methanol. Yield: 82.7 %. mp: > 300 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 3.71 (s, 3H, OCH<sub>3</sub>), 4.42 (br m, 2H, NHCH<sub>2</sub>Ph), 6.46 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.84-6.86 (d, 2H, CH benzene ring), 7.24-7.31 (br m, 4H, 2 x CH benzene ring and C<sub>4</sub>-NH<sub>2</sub>), 7.66-7.76 (br m, 1H, NHCH<sub>2</sub>Ph), 8.86 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 43.7, 55.5, 96.4, 113.9 (2C), 129.0 (2C), 132.8, 157.0, 158.5, 163.3, 164.1, 165.9, 166.3. MS (ESI),  $m/z$ : 298  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{14}H_{15}N_7O$ : C 56.56, H 5.09, N 32.98. Found: C 56.67, H 5.11, N 32.85.

***N*<sup>7</sup>-(1-phenylethyl)pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (19).** Recryst. Solvent: methanol. Yield: 68.3 %. mp: > 300 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.41-1.43 (d, 3H, CHCH<sub>3</sub>), 5.14 (br s, 1H, CHCH<sub>3</sub>), 6.44 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 7.16-7.19 (m, 1H, CH benzene ring), 7.27-7.39 (m, 6H, 4 x CH benzene ring and C<sub>4</sub>-NH<sub>2</sub>), 7.79 (br m, 1H, NH), 8.85 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 23.4, 50.0, 96.4, 126.5 (2C), 126.8, 128.5 (2C), 146.2, 156.9, 163.3, 163.4, 165.9, 166.2. MS (ESI),  $m/z$ : 282  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{14}H_{15}N_7$ : C 59.77, H 5.37, N 34.85. Found: C 59.89, H 5.38, N 34.73.

***N*<sup>7</sup>-benzyl-*N*<sup>7</sup>-methylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (20).** Recryst. Solvent: methanol. Yield: 65.2 %. mp: 275-277 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 3.09 (s, 3H, NCH<sub>3</sub>), 4.89 (s, 2H, CH<sub>2</sub>Ph), 6.49 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.24-7.33 (m, 7H, 5 x CH benzene ring and C<sub>4</sub>-NH<sub>2</sub>), 8.94 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 35.1, 51.9, 96.4, 127.3 (2C), 127.7, 128.9 (2C), 139.0, 156.8, 163.4, 163.6, 166.0, 166.1. MS (ESI),  $m/z$ : 282  $[M +$

H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>14</sub>H<sub>15</sub>N<sub>7</sub>: C 59.77, H 5.37, N 34.85. Found: C 59.87, H 5.39, N 34.74.

***N*<sup>7</sup>-methyl-*N*<sup>7</sup>-(1-phenylethyl)pyrimido[4,5-d]pyrimidine-2,4,7-triamine (21).** Recryst. Solvent: acetonitrile/methanol. Yield: 76.8 %. mp: 192-194 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.53-1.55 (d, 3H, CHCH<sub>3</sub>), 2.82 (s, 3H, NCH<sub>3</sub>), 6.30 (br m, 1H, CHCH<sub>3</sub>), 6.48 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 7.28-7.34 (m, 7H, 5 x CH benzene ring and C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.7, 29.3, 51.8, 96.4, 127.2 (2C), 127.3, 128.8 (2C), 142.2, 156.7, 163.4, 163.4, 165.9, 166.0. MS (ESI), *m/z*: 296 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>15</sub>H<sub>17</sub>N<sub>7</sub>: C 61.00, H 5.80, N 33.20. Found: C 61.11, H 5.82, N 33.07.

***N*<sup>7</sup>-(1-(4-methoxyphenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (22).** Recryst. Solvent: acetonitrile/methanol. Yield: 60.1 %. mp: 184-187 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.49 (d, 3H, CHCH<sub>3</sub>), 2.78 (s, 3H, NCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.26 (br s, 1H, CHCH<sub>3</sub>), 6.48 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.88-6.90 (d, 2H, CH benzene ring), 7.20-7.22 (d, 2H, CH benzene ring), 7.39 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.95 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.8, 29.1, 51.2, 55.5, 96.4, 114.2 (2C), 128.4 (2C), 134.0, 156.7, 158.6, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 326 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>16</sub>H<sub>19</sub>N<sub>7</sub>O: C 59.06, H 5.89, N 30.13. Found: C 59.17, H 5.91, N 30.01.

***N*<sup>7</sup>-(1-([1,1'-biphenyl]-4-yl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (23).** Recryst. Solvent: methanol. Yield: 57.1 %. mp: > 300 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.57-1.59 (d, 3H, CHCH<sub>3</sub>), 2.87 (s, 3H, NCH<sub>3</sub>), 6.35 (br s, 1H, CHCH<sub>3</sub>), 6.51 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 7.34-7.48 (m, 7H, 5 x CH benzene rings and C<sub>4</sub>-NH<sub>2</sub>), 7.63-7.66 (m, 4H, CH benzene rings), 8.97 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.9, 29.4, 51.7, 96.5, 127.0 (2C), 127.1 (2C), 127.8 (2C), 127.9, 129.4 (2C), 139.2, 140.3, 141.5, 156.7, 163.4,

163.4, 166.0, 166.1. MS (ESI),  $m/z$ : 372  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{21}H_{21}N_7$ : C 67.90, H 5.70, N 26.40. Found: C 68.01, H 5.72, N 26.27.

***N*<sup>7</sup>-methyl-*N*<sup>7</sup>-(1-(4-phenoxyphenyl)ethyl)pyrimido[4,5-d]pyrimidine-2,4,7-triamine (24).**

Recryst. Solvent: acetonitrile/methanol. Yield: 41.9 %. mp: 185-187 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.52-1.54 (d, 3H, CHCH<sub>3</sub>), 2.84 (s, 3H, NCH<sub>3</sub>), 6.30 (br s, 1H, CHCH<sub>3</sub>), 6.49 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.97-7.01 (m, 4H, CH benzene rings), 7.11-7.15 (t, 1H, CH benzene ring), 7.30-7.40 (m, 6H, CH benzene rings and C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.8, 29.2, 51.4, 96.5, 118.9 (2C), 119.0 (2C), 123.8, 128.9 (2C), 130.5 (2C), 137.3, 155.9, 156.7, 157.2, 163.3, 163.4, 166.0, 166.1. MS (ESI),  $m/z$ : 388  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{21}H_{21}N_7O$ : C 65.10, H 5.46, N 25.31. Found: C 65.20, H 5.48, N 25.20.

***N*<sup>7</sup>-(1-(4-(4-methoxyphenoxy)phenyl)ethyl)pyrimido[4,5-d]pyrimidine-2,4,7-triamine (25).**

Recryst. Solvent: methanol. Yield: 82.5 %. mp: >300 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.40-1.42 (d, 3H, CHCH<sub>3</sub>), 3.91 (s, 3H, OCH<sub>3</sub>), 5.12 (br m, 1H, CHCH<sub>3</sub>), 6.45 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.83-6.85 (m, 2H, CH benzene rings), 6.92-6.98 (m, 4H, CH benzene rings), 7.20-7.36 (m, 4H, CH benzene rings and C<sub>4</sub>-NH<sub>2</sub>), 7.76-7.78 (br m, 1H, NH), 8.85 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 23.3, 49.4, 55.9, 96.4, 115.5 (2C), 117.5 (2C), 121.0 (2C), 128.0 (2C), 140.4, 150.1, 155.9, 156.9, 157.0, 163.3, 163.6, 165.9, 166.2. MS (ESI),  $m/z$ : 404  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{21}H_{21}N_7O_2$ : C 62.52, H 5.25, N 24.30. Found: C 62.62, H 5.27, N 24.20.

***N*<sup>7</sup>-(4-(4-methoxyphenoxy)benzyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (26).**

Recryst. Solvent: methanol. Yield: 68.0 %. mp: 244-247 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 3.08 (br s, 3H, NCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 4.85 (s, 2H, N(CH<sub>3</sub>)CH<sub>2</sub>Ph), 6.50 (br s, 2H, C<sub>2</sub>-



*NH*<sub>2</sub>), 6.86-6.88 (d, 2H, *CH* benzene rings), 6.93-6.99 (m, 4H, *CH* benzene rings), 7.23 (br m, 2H, *CH* benzene rings), 7.41 (br s, 2H, C<sub>4</sub>-*NH*<sub>2</sub>), 8.94 (s, 1H, *CH* pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 35.0, 51.3, 55.9, 96.4, 115.5 (2C), 117.8 (2C), 121.0 (2C), 129.3 (2C), 133.2, 150.0, 156.0, 156.7, 157.4, 163.4, 163.5, 166.0, 166.1. MS (ESI), *m/z*: 404 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>21</sub>H<sub>21</sub>N<sub>7</sub>O<sub>2</sub>: C 62.52, H 5.25, N 24.30. Found: C 62.63, H 5.26, N 24.19.

***N*<sup>7</sup>-(1-(4-(4-methoxyphenoxy)phenyl)propyl)-*N*<sup>7</sup>-methylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (27).** Recryst. Solvent: acetonitrile/methanol. Yield: 60.7 %. mp: 186-190 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 0.85 (t, 3H, CH<sub>2</sub>CH<sub>3</sub>), 1.90-2.07 (m, 2H, CH<sub>2</sub>CH<sub>3</sub>), 2.79-2.82 (br s, 3H, NCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 6.10 (br m, 1H, CHCH<sub>3</sub>), 6.48 (br s, 2H, C<sub>2</sub>-*NH*<sub>2</sub>), 6.86-6.88 (m, 2H, *CH* benzene rings), 6.93-7.00 (m, 4H, *CH* benzene rings), 7.20-7.38 (br m, 4H, *CH* benzene rings and C<sub>4</sub>-*NH*<sub>2</sub>), 8.94 (s, 1H, *CH* pyrimidine ring). <sup>13</sup>C -NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 11.5, 23.5, 28.9, 55.9, 57.2, 96.3, 115.5 (2C), 117.5 (2C), 121.1 (2C), 129.1 (2C), 135.6, 149.9, 156.1, 156.5, 157.4, 163.4, 164.0, 166.0, 166.1. MS (ESI), *m/z*: 432 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>23</sub>H<sub>25</sub>N<sub>7</sub>O<sub>2</sub>: C 64.02, H 5.84, N 22.72. Found: C 64.12, H 5.86, N 22.60.

***N*<sup>7</sup>-(1-(4-(4-methoxyphenoxy)phenyl)-2-methylpropyl)-*N*<sup>7</sup>-methylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (28).** Recryst. Solvent: acetonitrile. Yield: 64.0 %. mp: 162-165 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) (mixture of two rotamers 50:50) δ ppm: 0.83 and 0.89 (two d, 6H, CH(CH<sub>3</sub>)<sub>2</sub>), 2.51-2.58 (m, 1H, CH(CH<sub>3</sub>)<sub>2</sub>), 2.81 and 2.89 (two br s, 3H, NCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 5.82-5.84 (m, 1H, CHCH(CH<sub>3</sub>)<sub>2</sub>), 6.46 (two br s, 2H, C<sub>2</sub>-*NH*<sub>2</sub>), 6.86-6.88 (m, 2H, *CH* benzene rings), 6.93-7.01 (m, 4H, *CH* benzene rings), 7.36-7.42 (m, 4H, 2 x *CH* benzene rings and C<sub>4</sub>-*NH*<sub>2</sub>), 8.91 and 8.95 (two s, 1H, *CH* pyrimidine ring). <sup>13</sup>C -NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 20.0 (2C), 21.0, 27.6, 28.9, 55.9, 96.2, 115.5 (2C), 117.4 (2C), 121.3 (2C),

130.1 (2C), 134.4, 149.7, 156.1, 156.2, 156.8, 157.5, 163.3, 163.7, 166.0. MS (ESI),  $m/z$ : 446 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>24</sub>H<sub>27</sub>N<sub>7</sub>O<sub>2</sub>: C 64.70, H 6.11, N 22.01. Found: C 64.81, H 6.13, N 21.89.

***N*<sup>7</sup>-((4-(4-methoxyphenoxy)phenyl)(phenyl)methyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (29)**. Recryst. Solvent: methanol. Yield: 58.8 %. mp: 248-250 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.89 (br s, 3H, NCH<sub>3</sub>), 3.93 (s, 3H, OCH<sub>3</sub>), 6.52 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.91-7.04 (m, 6H, CH benzene rings), 7.13-7.18 (m, 4H, CH benzene rings), 7.28-7.43 (m, 6H, 4 x CH benzene rings and C<sub>4</sub>-NH<sub>2</sub>), 8.97 (m, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 32.0, 55.9, 61.2, 96.8, 115.6 (2C), 117.6 (2C), 121.3 (2C), 127.6, 128.7 (2C), 128.9 (2C), 130.5 (2C), 134.4, 140.5, 149.7, 156.2, 156.7, 157.6, 163.4, 163.6, 166.0, 166.1. MS (ESI),  $m/z$ : 480 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>27</sub>H<sub>25</sub>N<sub>7</sub>O<sub>2</sub>: C 67.63, H 5.25, N 20.45. Found: C 67.74, H 5.27, N 20.34.

***N*<sup>7</sup>-(1-(4-(3-methoxyphenoxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (30)**. Recryst. Solvent: acetonitrile. Yield: 47.4 %. mp: 152-154 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.52-1.54 (d, 3H, CHCH<sub>3</sub>), 2.83 (br s, 3H, NCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.31 (br s, 1H, CHCH<sub>3</sub>), 6.49-6.54 (br m, 3H, CH benzene rings and C<sub>2</sub>-NH<sub>2</sub>), 6.58 (t, 1H, CH benzene ring), 6.69-6.72 (dd, 1H, CH benzene ring), 6.98-7.00 (d, 2H, CH benzene rings), 7.25-7.32 (m, 3H, CH benzene rings), 7.40 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.8, 29.2, 51.4, 55.7, 96.5, 100.0, 105.0, 109.5, 110.8, 119.1 (2C), 128.9, 130.9 (2C), 137.4, 155.7, 156.7, 158.4, 161.2, 163.3, 163.4, 166.0. MS (ESI),  $m/z$ : 418 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>22</sub>H<sub>23</sub>N<sub>7</sub>O<sub>2</sub>: C 63.30, H 5.55, N 23.49. Found: C 63.41, H 5.57, N 23.38.

***N*<sup>7</sup>-(1-(4-(3,4-dimethoxyphenoxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (31)**. Recryst. Solvent: acetonitrile/methanol. Yield: 69.2 %. mp: 185-188 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.50-1.52 (d, 3H, CHCH<sub>3</sub>), 2.82 (br s, 3H, NCH<sub>3</sub>), 3.72 (s, 3H, OCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 6.29 (br s, 1H, CHCH<sub>3</sub>), 6.48-6.53 (br m, 3H, CH benzene ring and C<sub>2</sub>-NH<sub>2</sub>), 6.74 (d, 1H, CH benzene ring), 6.89-6.95 (m, 3H, CH benzene rings), 7.25-7.27 (m, 2H, CH benzene rings), 7.40 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.9, 29.2, 51.3, 56.1, 56.4, 96.5, 105.3, 110.9, 113.0, 117.6 (2C), 128.7 (2C), 136.3, 145.8, 150.1, 150.3, 156.7, 157.2, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 448 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>23</sub>H<sub>25</sub>N<sub>7</sub>O<sub>3</sub>: C 61.73, H 5.63, N 21.91. Found: C 61.84, H 5.65, N 21.80.

***N*<sup>7</sup>-(1-(4-(3,5-dimethoxyphenoxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (32)**. Recryst. Solvent: methanol. Yield: 75.6 %. mp: 233-235 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.52-1.54 (d, 3H, CHCH<sub>3</sub>), 2.83 (br s, 3H, NCH<sub>3</sub>), 3.70 (s, 6H, 2 x OCH<sub>3</sub>), 6.13 (m, 2H, CH benzene rings), 6.28 (m, 2H, CHCH<sub>3</sub> and CH benzene ring), 6.48 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.99 (d, 2H, CH benzene rings), 7.29-7.40 (m, 4H, 2 x CH benzene rings and C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.8, 29.2, 51.4, 55.8 (2C), 95.9, 96.5, 97.5 (2C), 119.2 (2C), 128.9 (2C), 137.5, 155.5, 156.7, 159.0, 161.8 (2C), 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 448 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>23</sub>H<sub>25</sub>N<sub>7</sub>O<sub>3</sub>: C 61.73, H 5.63, N 21.91. Found: C 61.82, H 5.64, N 21.82.

***N*<sup>7</sup>-((4-(3,5-dimethoxyphenoxy)phenyl)(phenyl)methyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (33)**. Recryst. Solvent: acetonitrile. Yield: 58.2 %. mp: 152-154 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.91 (br s, 3H, NCH<sub>3</sub>), 3.71 (s, 6H, 2 x OCH<sub>3</sub>), 6.17 (d, 2H, CH benzene rings), 6.30 (t, 1H, CH benzene rings), 6.52 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>),

7.02-7.04 (d, 2H, CH benzene rings), 7.17-7.20 (m, 4H, CH benzene rings), 7.29-7.46 (m, 6H, CH benzene rings, CHPh and C<sub>4</sub>-NH<sub>2</sub>), 8.98 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 32.0, 55.8 (2C), 61.2, 96.0, 97.8 (2C), 119.1 (2C), 119.1, 127.6, 128.7 (2C), 129.0 (2C), 130.5 (2C), 135.5, 140.5, 155.9, 156.8, 158.8, 161.9 (2C), 162.0, 163.4, 163.6, 166.1. MS (ESI), *m/z*: 510 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>28</sub>H<sub>27</sub>N<sub>7</sub>O<sub>3</sub>: C 66.00, H 5.34, N 19.24. Found: C 66.10, H 5.35, N 19.14.

***N*<sup>7</sup>-(1-(4-(3,4,5-trimethoxyphenoxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (34).** Recryst. Solvent: acetonitrile. Yield: 67.1 %. mp: 172-176 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.51-1.53 (d, 3H, CHCH<sub>3</sub>), 2.82 (br s, 3H, NCH<sub>3</sub>), 3.64 (s, 3H, OCH<sub>3</sub>), 3.71 (s, 6H, 2 x OCH<sub>3</sub>), 6.30 (br s, 1H, CHCH<sub>3</sub>), 6.37 (s, 2H, CH benzene ring), 6.48 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.94-6.96 (d, 2H, CH benzene ring), 7.27-7.29 (d, 2H, CH benzene ring) 7.40 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.9, 29.2, 51.3, 56.4 (2C), 60.6, 96.5, 97.8 (2C), 117.9 (2C), 128.8 (2C), 134.4, 136.7, 152.6, 154.1, 156.6, 156.7 (2C), 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 478 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>24</sub>H<sub>27</sub>N<sub>7</sub>O<sub>4</sub>: C 60.37, H 5.70, N 20.53. Found: C 60.47, H 5.72, N 20.43.

***N*<sup>7</sup>-methyl-*N*<sup>7</sup>-(1-(4-(4-(trifluoromethoxy)phenoxy)phenyl)ethyl)pyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (35).** Recryst. Solvent: toluene. Yield: 58.3 %. mp: 140-143 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.53-1.55 (d, 3H, CHCH<sub>3</sub>), 2.85 (br s, 3H, NCH<sub>3</sub>), 6.31 (br s, 1H, CHCH<sub>3</sub>), 6.49 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 7.03-7.10 (m, 4H, CH benzene ring), 7.33-7.39 (m, 6H, CH benzene ring and C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.9, 29.2, 51.4, 96.5, 119.5 (2C), 120.0 (2C), 120.6 (q OCF<sub>3</sub>), 123.4 (2C), 129.1 (2C), 138.0, 144.0, 155.3, 156.3, 156.7, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 472 [M + H]<sup>+</sup>. Elemental

analysis calculated (%) for C<sub>22</sub>H<sub>20</sub>F<sub>3</sub>N<sub>7</sub>O<sub>2</sub>: C 56.05, H 4.28, N 20.80. Found: C 56.15, H 4.30, N 20.70.

***N*<sup>7</sup>-methyl-*N*<sup>7</sup>-(1-(4-(4-(methylthio)phenoxy)phenyl)ethyl)pyrimido[4,5-d]pyrimidine-2,4,7-triamine (36)**. Recryst. Solvent: toluene. Yield: 65.6 %. mp: 145-146 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.52-1.54 (d, 3H, CHCH<sub>3</sub>), 2.46 (s, 3H, SCH<sub>3</sub>), 2.83 (br s, 3H, NCH<sub>3</sub>), 6.30 (br m, 1H, CHCH<sub>3</sub>), 6.48 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.96-7.00 (m, 4H, CH benzene rings), 7.28-7.31 (m, 4H, CH benzene rings), 7.39 (br s, 2H C<sub>4</sub>-NH<sub>2</sub>), 8.95 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.2, 16.9, 29.2, 51.4, 96.5, 118.7 (2C), 119.9 (2C), 128.9 (2C), 128.9 (2C), 132.7, 137.2, 154.9, 156.0, 156.7, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 434 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>22</sub>H<sub>23</sub>N<sub>7</sub>OS: C 60.95, H 5.35, N 22.62, S 7.40. Found: C 61.06, H 5.36, N 22.52, S 7.37.

***N*<sup>7</sup>-(1-(4-(4-(benzyloxy)phenoxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (37)**. Recryst. Solvent: cyclohexane. Yield: 77.8 %. mp: 129-132 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.50-1.52 (d, 3H, CHCH<sub>3</sub>), 2.82 (br s, 3H, NCH<sub>3</sub>), 5.08 (s, 2H, OCH<sub>2</sub>Ph), 6.28 (br s, 1H, CHCH<sub>3</sub>), 6.48 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.88-6.90 (d, 2H, CH benzene rings), 6.98-7.05 (m, 4H, CH benzene rings), 7.26-7.28 (d, 2H, CH benzene rings), 7.34-7.47 (m, 7H, 5 x CH benzene rings and C<sub>4</sub>-NH<sub>2</sub>), 8.96 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.9, 29.2, 51.3, 70.1, 96.5, 116.5 (2C), 117.7 (2C), 121.0 (2C), 128.2 (2C), 128.3 (2C), 128.8, 128.9 (2C), 136.4, 137.5, 150.1, 155.1, 156.7, 157.2, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 494 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>28</sub>H<sub>27</sub>N<sub>7</sub>O<sub>2</sub>: C 68.14, H 5.51, N 19.87. Found: C 68.25, H 5.53, N 19.77.

***N*<sup>7</sup>-(1-(4-((4-methoxynaphthalen-1-yl)oxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (38)**. Recryst. Solvent: methanol. Yield: 71.9 %. mp: >300 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.49-1.51 (d, 3H, CHCH<sub>3</sub>), 2.82 (br s, 3H, NCH<sub>3</sub>), 3.98 (s, 3H, OCH<sub>3</sub>), 6.28 (br s, 1H, CHCH<sub>3</sub>), 6.46 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.89-6.96 (m, 3H, CH aromatic rings), 7.07-7.09 (d, 1H, CH naphthalene ring), 7.25-7.27 (d, 2H, CH aromatic rings), 7.38 (br s, 2H, C<sub>4</sub>-NH<sub>2</sub>), 7.53-7.58 (m, 2H, CH naphthalene ring), 7.89-7.91 (m, 1H, CH naphthalene ring), 8.19-8.21 (m, 1H, CH naphthalene ring), 8.95 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.9, 29.2, 51.3, 56.2, 96.4, 104.5, 116.3, 116.9 (2C), 121.9, 122.4, 126.3, 126.5, 127.4, 127.6, 128.8 (2C), 136.2, 144.9, 152.1, 156.7, 158.0, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 468 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>26</sub>H<sub>25</sub>N<sub>7</sub>O<sub>2</sub>: C 66.79, H 5.39, N 20.97. Found: C 66.90, H 5.41, N 20.85.

***N*<sup>7</sup>-(1-(4-((4-methoxybenzyl)oxy)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (39)**. Recryst. Solvent: methanol. Yield: 78.1 %. mp: 225-228 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.48-1.50 (d, 3H, CHCH<sub>3</sub>), 2.79 (br s, 3H, NCH<sub>3</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 4.99 (s, 2H, OCH<sub>2</sub>Ph), 6.26 (br s, 1H, CHCH<sub>3</sub>), 6.47 (br s, 2H, C<sub>2</sub>-NH<sub>2</sub>), 6.93-6.97 (m, 4H, CH benzene rings), 7.20-7.22 (d, 2H, CH benzene rings), 7.36-7.38 (m, 4H, 2 x CH benzene rings and C<sub>4</sub>-NH<sub>2</sub>), 8.95 (s, 1H, CH pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.8, 29.1, 51.2, 55.5, 69.4, 96.4, 114.3 (2C), 115.0 (2C), 128.4 (2C), 129.4, 129.9 (2C), 134.2, 156.7, 157.8, 159.4, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 432 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>23</sub>H<sub>25</sub>N<sub>7</sub>O<sub>2</sub>: C 64.02, H 5.84, N 22.72. Found: C 64.13, H 5.85, N 22.61.

***N*<sup>7</sup>-(1-(4-((4-methoxyphenyl)thio)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-d]pyrimidine-2,4,7-triamine (40)**. Recryst. Solvent: acetonitrile. Yield: 72.9 %. mp: 160-163 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.48-1.50 (d, 3H, CHCH<sub>3</sub>), 2.81 (br s, 3H, NCH<sub>3</sub>), 3.77 (s, 3H, OCH<sub>3</sub>),

6.24 (br s, 1H, *CHCH*<sub>3</sub>), 6.49 (br s, 2H, *C*<sub>2</sub>-*NH*<sub>2</sub>), 6.98-7.01 (d, 2H, *CH* benzene rings), 7.10-7.12 (d, 2H, *CH* benzene rings), 7.21-7.41 (m, 6H, 4 x *CH* benzene rings and *C*<sub>4</sub>-*NH*<sub>2</sub>), 8.94 (s, 1H, *CH* pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.8, 29.3, 51.5, 55.8, 96.5, 115.8 (2C), 123.8, 128.2 (2C), 128.6 (2C), 135.5 (2C), 136.4, 140.4, 156.7, 160.1, 163.3, 163.4, 166.0, 166.1. MS (ESI), *m/z*: 434 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>22</sub>H<sub>23</sub>N<sub>7</sub>OS: C 60.95, H 5.35, N 22.62, S 7.40. Found: C 61.06, H 5.36, N 22.51, S 7.37.

***N*<sup>7</sup>-(1-(4-((3,5-dimethoxyphenyl)thio)phenyl)ethyl)-*N*<sup>7</sup>-methylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (41).** Recryst. Solvent: toluene. Yield: 64.0 %. mp: 138-139 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.53-1.55 (d, 3H, *CHCH*<sub>3</sub>), 2.83 (br s, 3H, *NCH*<sub>3</sub>), 3.69 (s, 6H, 2 x *OCH*<sub>3</sub>), 6.30 (br s, 1H, *CHCH*<sub>3</sub>), 6.36 (m, 2H, *CH* benzene ring), 6.41 (m, 1H, *CH* benzene ring), 6.50 (br s, 2H, *C*<sub>2</sub>-*NH*<sub>2</sub>), 7.31-7.38 (m, 6H, 4 x *CH* benzene ring and *C*<sub>4</sub>-*NH*<sub>2</sub>), 8.96 (s, 1H, *CH* pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 16.7, 29.4, 51.6, 55.8 (2C), 96.5, 99.4, 107.9 (2C), 128.5 (2C), 132.3, 132.3 (2C), 138.0, 142.4, 156.7, 161.3 (2C), 163.4, 166.0, 166.1, 166.2. MS (ESI), *m/z*: 464 [M + H]<sup>+</sup>. Elemental analysis calculated (%) for C<sub>23</sub>H<sub>25</sub>N<sub>7</sub>O<sub>2</sub>S: C 59.59, H 5.44, N 21.15, S 6.92. Found: C 59.71, H 5.46, N 21.04, S 6.89.

***N*<sup>7</sup>-((4-((3,5-dimethoxyphenyl)thio)phenyl)(phenyl)methyl)-*N*<sup>7</sup>-methylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (42).** Recryst. Solvent: toluene. Yield: 76.2 %. mp: 139-141 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.90 (br s, 3H, *NCH*<sub>3</sub>), 3.70 (s, 6H, 2 x *OCH*<sub>3</sub>), 6.42 (m, 3H, *CH* benzene rings), 6.53 (br s, 2H, *C*<sub>2</sub>-*NH*<sub>2</sub>), 7.17-7.21 (m, 4H, *CH* benzene rings), 7.30-7.34 (m, 1H, *CH* benzene ring), 7.37-7.46 (m, 7H, *CH* benzene rings, *CHPh* and *C*<sub>4</sub>-*NH*<sub>2</sub>), 8.97 (s, 1H, *CH* pyrimidine ring). <sup>13</sup>C-NMR (100 MHz, *d*<sub>6</sub>-DMSO) δ 32.1, 55.8 (2C), 61.5, 96.9, 99.7, 108.3 (2C), 127.8, 128.9 (2C), 129.0 (2C), 130.0 (2C), 132.0 (2C), 133.0, 137.5, 140.1, 140.3, 156.8, 161.4 (2C), 163.4, 163.5, 166.1, 166.2. MS (ESI), *m/z*: 526 [M + H]<sup>+</sup>. Elemental analysis

calculated (%) for  $C_{28}H_{27}N_7O_2S$ : C 63.98, H 5.18, N 18.65, S 6.10. Found: C 64.09, H 5.20, N 18.54, S 6.07.

***N*-7-(1-(4-((4-methoxyphenyl)amino)phenyl)ethyl)-*N*-7-methylpyrimido[4,5-*d*]pyrimidine-2,4,7-triamine (43).** Recryst. Solvent: toluene. Yield: 52.1 %. mp: 146-149 °C.  $^1H$ -NMR (400 MHz;  $d_6$ -DMSO)  $\delta$  ppm: 1.46-1.48 (d, 3H,  $CHCH_3$ ), 2.79 (br s, 3H,  $NCH_3$ ), 3.71 (s, 3H,  $OCH_3$ ), 6.23 (br s, 1H,  $CHCH_3$ ), 6.46 (br s, 2H,  $C_2-NH_2$ ), 6.84-6.90 (m, 4H, CH benzene rings), 7.01-7.03 (d, 2H, CH benzene rings), 7.09-7.11 (m, 2H, CH benzene rings), 7.38 (br s, 2H,  $C_4-NH_2$ ), 7.83 (s, 1H,  $NHPh-OCH_3$ ), 8.95 (s, 1H, CH pyrimidine ring).  $^{13}C$ -NMR (100 MHz,  $d_6$ -DMSO)  $\delta$  16.7, 29.1, 51.3, 55.7, 96.3, 115.0 (2C), 115.2 (2C), 120.6 (2C), 128.2 (2C), 131.7, 136.7, 144.4, 154.1, 156.7, 163.3, 163.4, 166.0, 166.1. MS (ESI),  $m/z$ : 417  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{22}H_{24}N_8O$ : C 63.45, H 5.81, N 26.90. Found: C 63.56, H 5.83, N 26.79.

***N*-(4-(1-((5,7-diaminopyrimido[4,5-*d*]pyrimidin-2-yl)(methyl)amino)ethyl)phenyl)-4-methoxybenzamide (44).** Recryst. Solvent: acetonitrile/methanol. Yield: 74.0 %. mp: 196-198 °C.  $^1H$ -NMR (400 MHz;  $d_6$ -DMSO)  $\delta$  ppm: 1.51-1.53 (d, 3H,  $CHCH_3$ ), 2.81 (br s, 3H,  $NCH_3$ ), 3.83 (s, 3H,  $OCH_3$ ), 6.30 (br m, 1H,  $CHCH_3$ ), 6.46 (br s, 2H,  $C_2-NH_2$ ), 7.04-7.06 (d, 2H, CH benzene ring), 7.24-7.26 (d, 2H, CH benzene ring), 7.38 (br s, 2H,  $C_4-NH_2$ ), 7.71-7.73 (d, 2H, CH benzene ring), 7.94-7.96 (d, 2H, CH benzene ring), 8.96 (s, 1H, CH pyrimidine ring), 10.08 (s, 1H,  $NHCO$ ).  $^{13}C$ -NMR (100 MHz,  $d_6$ -DMSO)  $\delta$  16.7, 29.2, 51.5, 55.9, 96.4, 114.0 (2C), 120.7 (2C), 127.4, 127.5 (2C), 130.0 (2C), 137.1, 138.6, 156.7, 162.3, 163.3, 163.4, 165.3, 166.0, 166.1. MS (ESI),  $m/z$ : 445  $[M + H]^+$ . Elemental analysis calculated (%) for  $C_{23}H_{24}N_8O_2$ : C 62.15, H 5.44, N 25.21. Found: C 62.25, H 5.46, N 25.10.



**General Procedure for the Synthesis of the Intermediate Ketones 45f-q, s-u.** A mixture of the appropriate alkyl/phenyl 4'-fluorophenyl ketone (1 equiv, 3 mmol), the properly substituted phenol, 1-naphthol or phenylthiol (1 equiv, 3 mmol), and anhydrous potassium carbonate (1.2 equiv, 3.6 mmol) in anhydrous DMF (3 mL) was stirred at 175 °C for 5 h. After the completion of the reaction, the medium was quenched with water (50 mL) and the product was extracted with ethyl acetate (3 × 25 mL). The organic phase was washed with saturated sodium chloride (2 × 50 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by silica gel column chromatography eluting with a mixture of ethyl acetate/*n*-hexane to obtain the pure ketones **45f-q, s-u**.

**1-(4-(4-Methoxyphenoxy)phenyl)propan-1-one (45f).**<sup>62</sup> Yield: 83 %. mp: 50-51 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.14 (t, 3H, COCH<sub>2</sub>CH<sub>3</sub>), 2.87-2.89 (q, 2H, COCH<sub>2</sub>CH<sub>3</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 6.86 (m, 4H, CH benzene rings), 6.95 (m, 2H, CH benzene ring), 7.85 (m, 2H, CH benzene ring).

**1-(4-(4-Methoxyphenoxy)phenyl)-2-methylpropan-1-one (45g).** Oil. Yield: 68 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.13 (d, 6H, CH(CH<sub>3</sub>)<sub>2</sub>), 3.42-3.45 (m, 1H, CH(CH<sub>3</sub>)<sub>2</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 6.86 (m, 4H, CH benzene rings), 6.95 (m, 2H, CH benzene ring), 7.85 (m, 2H, CH benzene ring).

**(4-(4-Methoxyphenoxy)phenyl)(phenyl)methanone (45h).**<sup>63</sup> Yield: 86 %. mp: 107-108 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 3.76 (s, 3H, OCH<sub>3</sub>), 6.85-6.91 (m, 4H, CH benzene rings), 6.96-6.99 (m, 2H, CH benzene ring), 7.38-7.42 (m, 2H, CH ring), 7.48-7.50 (m, 1H, CH benzene ring), 7.68-7.74 (m, 4H, CH benzene rings).

**1-(4-(3-Methoxyphenoxy)phenyl)ethan-1-one (45i).**<sup>64</sup> Yield: 59 %. mp: 65-67 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.60 (s, 3H, COCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 6.64-6.68 (m, 2H, CH

benzene ring), 6.76-6.78 (m, 1H, *CH* benzene ring), 7.03-7.05 (m, 2H, *CH* benzene ring), 7.28-7.33 (m, 1H, *CH* benzene ring), 7.97 (m, 2H, *CH* benzene ring).

**1-(4-(3,4-Dimethoxyphenoxy)phenyl)ethan-1-one (45j).** Recryst. Solvent:cyclohexane. Yield: 86 %. mp: 99-101 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.50 (s, 3H, COCH<sub>3</sub>), 3.78 (s, 3H, OCH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 6.55-6.58 (m, 2H, *CH* benzene ring), 6.79-6.81 (d, 1H, *CH* benzene ring), 6.87-6.91 (m, 2H, *CH* benzene ring), 7.85-7.87 (m, 2H, *CH* benzene ring).

**1-(4-(3,5-Dimethoxyphenoxy)phenyl)ethan-1-one (45k).** Recryst. Solvent:cyclohexane. Yield: 79 %. mp: 85-87 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.51 (s, 3H, COCH<sub>3</sub>), 3.70 (s, 6H, 2 x OCH<sub>3</sub>), 6.15 (s, 2H, *CH* benzene ring), 6.23 (s, 1H, *CH* benzene ring), 6.96-6.98 (d, 2H, *CH* benzene ring), 7.86-7.88 (d, 2H, *CH* benzene ring).

**(4-(3,5-Dimethoxyphenoxy)phenyl)(phenyl)methanone (45l).** Oil. Yield: 92.0 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 3.78 (s, 6H, 2 x OCH<sub>3</sub>), 6.25 (d 2H, *CH* benzene ring), 6.30 (m, 1H, *CH* benzene ring), 7.07 (d, 2H, *CH* benzene ring), 7.46-7.50 (m, 2H, *CH* benzene ring), 7.56-7.60 (m, 1H, *CH* benzene ring), 7.77-7.79 (m, 4H, *CH* benzene rings).

**1-(4-(3,4,5-Trimethoxyphenoxy)phenyl)ethan-1-one (45m).** Recryst. Solvent:cyclohexane. Yield: 66%. mp: 94-96 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.60 (s, 3H, COCH<sub>3</sub>), 3.83 (s, 6H, 3,5-(OCH<sub>3</sub>)<sub>2</sub>), 3.88 (s, 3H, 4-OCH<sub>3</sub>), 6.34 (s, 2H, *CH* benzene ring), 7.02 (d, 2H, *CH* benzene ring), 7.96 (d, 2H, *CH* benzene ring).

**1-(4-(4-(Trifluoromethoxy)phenoxy)phenyl)ethan-1-one (45n).**<sup>65</sup> Oil. Yield: 95 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.51 (s, 3H, COCH<sub>3</sub>), 6.95 (d, 2H, *CH* benzene ring), 7.01 (d, 2H, *CH* benzene ring), 7.16-7.19 (d, 2H, *CH* benzene ring), 7.89 (d, 2H, *CH* benzene ring).

**1-(4-(4-(Methylthio)phenoxy)phenyl)ethan-1-one (45o).** Recryst. Solvent:cyclohexane. Yield: 87 %. mp: 83-84 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.50 (s, 3H, SCH<sub>3</sub>), 2.57 (s, 3H,

COCH<sub>3</sub>), 6.93-7.03 (m, 4H, CH benzene ring), 7.30 (m, 2., CH benzene ring), 7.92-7.95 (m, 2H, CH benzene ring).

**1-(4-(4-(Benzyloxy)phenoxy)phenyl)ethan-1-one (45p).**<sup>66</sup> Yield: 70%. mp: 101-103 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.49 (s, 3H, COCH<sub>3</sub>), 5.00 (s, 2H, CH<sub>2</sub>Ph), 6.86-6.89 (m, 2H, CH benzene ring), 6.94 (m, 4H, CH benzene rings), 7.27-7.39 (m, 5H, CH benzene rings), 7.83-7.85 (m, 2H, CH benzene ring).

**1-(4-((4-Methoxynaphthalen-1-yl)oxy)phenyl)ethan-1-one (45q).** Oil. Yield: 50 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.48 (s, 3H, COCH<sub>3</sub>), 3.96 (s, 3H, OCH<sub>3</sub>), 6.70-6.72 (d, 1H, CH naphthalene ring), 6.87-6.89 (m, 2H, CH benzene ring), 7.02-7.04 (d, 1H, CH naphthalene ring), 7.38-7.47 (m, 2H, CH naphthalene ring), 7.77-7.79 (d, 1H, CH naphthalene ring), 7.82 (m, 2H, CH benzene ring), 8.22-8.24 (d, 1H, CH naphthalene ring).

**1-(4-((4-Methoxyphenyl)thio)phenyl)ethan-1-one (45s).**<sup>67</sup> Oil. Yield: 54 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.47 (s, 3H, COCH<sub>3</sub>), 3.79 (s, 3H, OCH<sub>3</sub>), 6.88-6.90 (d, 2H, CH benzene ring), 7.01-7.04 (d, 2H, CH benzene ring), 7.41 (d, 2H, CH benzene ring), 7.72 (d, 2H, CH benzene ring).

**1-(4-((3,5-Dimethoxyphenyl)thio)phenyl)ethan-1-one (45t).** Recryst. Solvent:cyclohexane. Yield: 72 %. mp: 41-42 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.56 (s, 3H, COCH<sub>3</sub>), 3.77 (s, 6H, 2 x OCH<sub>3</sub>), 6.46 (m, 1H, CH benzene ring), 6.62 (m, 2H, CH benzene ring), 7.27 (d, 2H, CH benzene ring), 7.83 (d, 2H, CH benzene ring).

**(4-((3,5-Dimethoxyphenyl)thio)phenyl)(phenyl)methanone (45u).** Oil. Yield: 80 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 3.71 (s, 6H, 2 x OCH<sub>3</sub>), 6.39 (m, 1H, CH benzene ring), 6.58 (m, 2H, CH benzene ring), 7.23 (d, 2H, CH benzene ring), 7.38-7.42 (m, 2H, CH benzene ring), 7.48-7.52 (m, 1H, CH benzene ring), 7.65 (d, 2H, CH benzene ring), 7.69-7.71 (m, 2H, CH benzene ring).

**General Procedure for the Synthesis of the intermediate amines 46i-f'.** A 2 M methylamine solution in methanol or, alternatively, 7 M ammonia solution in methanol (3-6 equiv, 6-12 mmol) was added to a solution of titanium *isopropoxide* (1.3-2 equiv, 2.6-4 mmol) and the carbonyl compounds **45a-w** (1 equiv, 2 mmol) in THF (5 mL), and the reaction mixture was stirred under nitrogen atmosphere at room temperature. After 5-6 h, sodium borohydride (1.1 equiv, 2.2 mmol) was added portion wise at 0 °C, and the mixture was stirred for 2 h at room temperature. After the completion, the reaction was quenched with distilled water (2 mL) and acidified at 0 °C with 1 M hydrochloric acid until pH was 1-2. The resulting suspension was filtered on celite and washed with a mixture of water (50 mL) and ethyl acetate (50 mL). The filtrate and the washings were combined, extracted with ethyl acetate (3 × 30 mL), basified with 10 % w/w sodium hydroxide up to pH 10-12 and further extracted with ethyl acetate (3 × 30 mL). The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give an oily crude product. This was finally purified by silica gel column chromatography eluting with a mixture of chloroform/methanol/ammonia or ethyl acetate/*n*-hexane thus affording the pure amines **46i-f'**.

**1-(4-Methoxyphenyl)-*N*-methylethan-1-amine (46i).**<sup>68</sup> Oil. Yield: 87 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.26 (d, 3H, CHCH<sub>3</sub>), 2.22 (s, 3H, NHCH<sub>3</sub>), 3.52 (m, 1H, CHCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.81 (d, 2H, CH benzene ring), 7.15 (d, 2H, CH benzene ring).

**1-([1,1'-Biphenyl]-4-yl)-*N*-methylethan-1-amine (46j).** Oil. Yield: 55 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.42 (d, 3H, CHCH<sub>3</sub>), 2.37 (s, 3H, NHCH<sub>3</sub>), 3.71-3.73 (m, 1H, CHCH<sub>3</sub>), 7.35-7.41 (m, 3H, CH biphenyl ring), 7.44-7.47 (m, 2H, CH biphenyl ring), 7.58-7.62 (m, 4H, CH biphenyl ring).

***N*-methyl-1-(4-phenoxyphenyl)ethan-1-amine (46k).** Oil. Yield: 85 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.39 (d, 3H, CHCH<sub>3</sub>), 2.34 (s, 3H, NHCH<sub>3</sub>), 3.65-3.70 (m, 1H, CHCH<sub>3</sub>), 6.98-

7.04 (m, 4H, *CH* benzene rings), 7.09-7.13 (m, 1H, *CH* benzene ring), 7.29-7.32 (m, 2H, *CH* benzene ring), 7.35-7.37 (m, 2H, *CH* benzene ring).

**1-(4-(4-Methoxyphenoxy)phenyl)ethan-1-amine (46l).** Oil. Yield: 34 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.30 (d, 3H, CHCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 4.03 (m, 1H, CHCH<sub>3</sub>), 6.79-6.91 (m, 6H, *CH* benzene rings), 7.21 (m, 2H, *CH* benzene ring).

**1-(4-(4-Methoxyphenoxy)phenyl)-*N*-methylmethanamine (46m).** Oil. Yield: 78 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.38 (s, 3H, NHCH<sub>3</sub>), 3.63 (s, 2H, CH<sub>2</sub>NHCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.79-6.84 (m, 4H, *CH* benzene rings), 6.89-6.91 (d, 2H, *CH* benzene rings), 7.15-7.18 (m, 2H, *CH* benzene rings).

**1-(4-(4-Methoxyphenoxy)phenyl)-*N*-methylethan-1-amine (46n).** Oil. Yield: 73 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.27 (d, 3H, CHCH<sub>3</sub>), 2.24 (s, 3H, NHCH<sub>3</sub>), 3.55 (m, 1H, CHCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.79-6.84 (m, 4H, *CH* benzene rings), 6.89-6.92 (m, 2H, *CH* benzene ring), 7.14-7.16 (m, 2H, *CH* benzene ring).

**1-(4-(4-Methoxyphenoxy)phenyl)-*N*-methylpropan-1-amine (46o).** Oil. Yield: 83 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 0.72-0.75 (t, 3H, CHCH<sub>2</sub>CH<sub>3</sub>), 1.51-1.58 (m, 1H, CHCH<sub>2</sub>CH<sub>3</sub>), 1.63-1.71 (m, 1H, CHCH<sub>2</sub>CH<sub>3</sub>), 2.21 (s, 3H, NHCH<sub>3</sub>), 3.24-3.28 (m, 1H, CHCH<sub>2</sub>CH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 6.79-6.84 (m, 4H, *CH* benzene rings), 6.90-6.92 (m, 2H, *CH* benzene ring), 7.1 (m, 2H, *CH* benzene ring).

**1-(4-(4-Methoxyphenoxy)phenyl)-*N*,2-dimethylpropan-1-amine (46p).** Oil. Yield: 62 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 0.66-0.68 (d, 3H, CHCH<sub>3</sub>(CH<sub>3</sub>)), 0.88-0.90 (d, 3H, CHCH<sub>3</sub>(CH<sub>3</sub>)), 1.73-1.81 (m, 1H, CH(CH<sub>3</sub>)<sub>2</sub>), 2.16 (s, 3H, NHCH<sub>3</sub>), 3.08-3.10 (m, 1H, CHCH(CH<sub>3</sub>)<sub>2</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.79-6.83 (m, 4H, *CH* benzene rings), 6.89-6.94 (m, 2H, *CH* benzene ring), 7.07-7.10 (m, 2H, *CH* benzene ring).

**1-(4-(4-Methoxyphenoxy)phenyl)-*N*-methyl-1-phenylmethanamine (46q).** Oil. Yield: 60 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.33 (s, 3H, NHCH<sub>3</sub>), 3.72 (s, 3H, OCH<sub>3</sub>), 4.59 (m, 1H, CHPh), 6.77-6.80 (m, 4H, CH benzene rings), 6.86-6.89 (m, 2H, CH benzene ring), 7.14-7.16 (m, 1H, CH benzene ring), 7.22-7.25 (m, 4H, CH benzene rings), 7.29-7.31 (m, 2H, CH benzene ring).

**1-(4-(3-Methoxyphenoxy)phenyl)-*N*-methylethan-1-amine (46r).** Oil. Yield: 79 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.29 (d, 3H, CHCH<sub>3</sub>), 2.25 (s, 3H, NHCH<sub>3</sub>), 3.56-3.59 (m, 1H, CHCH<sub>3</sub>), 3.71 (s, 3H, OCH<sub>3</sub>), 6.50-6.52 (d, 2H, CH benzene ring), 6.56-6.58 (d, 1H, CH benzene ring), 6.90-6.92 (d, 2H, CH benzene ring), 7.12-7.20 (m, 3H, CH benzene rings).

**1-(4-(3,4-Dimethoxyphenoxy)phenyl)-*N*-methylethan-1-amine (46s).** Oil. Yield: 84 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.27 (d, 3H, CHCH<sub>3</sub>), 2.24 (s, 3H, NHCH<sub>3</sub>), 3.54-3.56 (m, 1H, CHCH<sub>3</sub>), 3.77 (s, 3H, OCH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 6.47-6.50 (d, 1H, CH benzene ring), 6.59 (s, 1H, CH benzene ring), 6.74-6.76 (d, 1H, CH benzene ring), 6.84-6.86 (d, 2H, CH benzene ring), 7.15-7.18 (m, 2H, CH benzene ring).

**1-(4-(3,5-Dimethoxyphenoxy)phenyl)-*N*-methylethan-1-amine (46t).** Oil. Yield: 72 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.8 (d, 3H, CHCH<sub>3</sub>), 2.25 (s, 3H, NHCH<sub>3</sub>), 3.54-3.59 (m, 1H, CHCH<sub>3</sub>), 3.68 (s, 6H, 2 x OCH<sub>3</sub>), 6.09 (m, 2H, CH benzene ring), 6.13 (m, 1H, CH benzene ring), 6.91-6.93 (d, 2H, CH benzene ring), 7.18-7.20 (d, 2H, CH benzene ring).

**1-(4-(3,5-Dimethoxyphenoxy)phenyl)-*N*-methyl-1-phenylmethanamine (46u).** Oil. Yield: 46 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.41 (s, 3H, NHCH<sub>3</sub>), 3.73 (s, 6H, 3,5-(OCH<sub>3</sub>)<sub>2</sub>), 4.68 (s, 1H, NHCHPh), 6.19 (m, 2H, CH benzene ring), 6.34 (m, 1H, CH benzene ring), 6.94-6.96 (d, 2H, CH benzene ring), 7.21-7.26 (m, 1H, CH benzene ring), 7.31-7.35 (m, 6H, CH phenyl rings).

***N*-Methyl-1-(4-(3,4,5-trimethoxyphenoxy)phenyl)ethan-1-amine (46v)**. Oil. Yield: 82 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.27-1.29 (d, 3H, CHCH<sub>3</sub>), 2.25 (s, 3H, NHCH<sub>3</sub>), 3.55-3.57 (m, 1H, CHCH<sub>3</sub>), 3.72 (s, 6H, 3,5-(OCH<sub>3</sub>)<sub>2</sub>), 3.76 (s, 3H, 4-OCH<sub>3</sub>), 6.20 (s, 2H, CH benzene ring), 6.88-6.90 (m, 2H, CH benzene ring), 7.18-7.20 (m, 2H, CH benzene ring).

***N*-Methyl-1-(4-(4-(trifluoromethoxy)phenoxy)phenyl)ethan-1-amine (46w)**. Oil. Yield: 77 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.28-1.30 (d, 3H, CHCH<sub>3</sub>), 2.25 (s, 3H, NHCH<sub>3</sub>), 3.57-3.59 (m, 1H, CHCH<sub>3</sub>), 6.89-6.93 (m, 4H, CH benzene rings), 7.09 (m, 2H, CH benzene ring), 7.21-7.23 (m, 2H, CH benzene ring).

***N*-Methyl-1-(4-(4-(methylthio)phenoxy)phenyl)ethan-1-amine (46x)**. Oil. Yield: 84 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.27-1.29 (d, 3H, CHCH<sub>3</sub>), 2.25 (s, 3H, SCH<sub>3</sub>), 2.40 (s, 3H, NHCH<sub>3</sub>), 3.54-3.58 (m, 1H, CHCH<sub>3</sub>), 6.87-6.89 (m, 4H, CH benzene rings), 7.18-7.20 (m, 4H, CH benzene rings).

**1-(4-(4-(Benzyloxy)phenoxy)phenyl)-*N*-methylethan-1-amine (46y)**. Recryst. Solvent: cyclohexane. Yield: 77 %. mp: 90-91 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.26-1.28 (d, 3H, CHCH<sub>3</sub>), 2.24 (s, 3H, NHCH<sub>3</sub>), 3.54-3.56 (m, 1H, CHCH<sub>3</sub>), 4.98 (s, 2H, OCH<sub>2</sub>Ph), 6.83-6.92 (m, 6H, CH benzene ring), 7.14-7.17 (m, 2H, CH benzene ring), 7.26-7.28 (m, 1H, CH benzene ring), 7.30-7.38 (m, 4H, CH benzene rings).

**1-(4-((4-Methoxynaphthalen-1-yl)oxy)phenyl)-*N*-methylethan-1-amine (46z)**. Oil. Yield: 84 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.26-1.28 (d, 3H, CHCH<sub>3</sub>), 2.24 (s, 3H, NHCH<sub>3</sub>), 3.52-3.57 (m, 1H, CHCH<sub>3</sub>), 3.93 (s, 3H, OCH<sub>3</sub>), 6.65-6.67 (d, 1H, CH naphthalene ring), 6.84-6.86 (d, 2H, CH naphthalene ring), 6.91-6.93 (d, 1H, CH naphthalene ring), 7.13-7.15 (m, 2H, CH benzene ring), 7.39-7.46 (m, 2H, CH benzene ring), 7.96-7.98 (d, 1H, CH naphthalene ring), 8.19-8.22 (d, 1H, CH naphthalene ring).

**1-(4-((4-Methoxybenzyl)oxy)phenyl)-*N*-methylethan-1-amine (46a')**. Recryst. Solvent: cyclohexane. Yield: 70 %. mp: 99-100 °C. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.25-1.27 (d, 3H, CHCH<sub>3</sub>), 2.23 (s, 3H, NHCH<sub>3</sub>), 3.50-3.55 (m, 1H, CHCH<sub>3</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 4.90 (s, 2H, OCH<sub>2</sub>Ph), 6.84-6.88 (m, 4H, CH benzene rings), 7.13-7.15 (m, 2H, CH benzene ring), 7.28-7.30 (m, 2H, CH benzene ring).

**1-(4-((4-Methoxyphenyl)thio)phenyl)-*N*-methylethan-1-amine (46b')**. Oil. Yield: 80 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.23-1.25 (d, 3H, CHCH<sub>3</sub>), 2.22 (s, 3H, NHCH<sub>3</sub>), 3.50-3.52 (m, 1H, CHCH<sub>3</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 6.81-6.83 (d, 2H, CH benzene ring), 7.06-7.12 (m, 4H, CH benzene rings), 7.32-7.34 (m, 2H, CH benzene ring).

**1-(4-((3,5-Dimethoxyphenyl)thio)phenyl)-*N*-methylethan-1-amine (46c')**. Oil. Yield: 78 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.26-1.28 (d, 3H, CHCH<sub>3</sub>), 2.24 (s, 3H, NHCH<sub>3</sub>), 3.54-3.59 (m, 1H, CHCH<sub>3</sub>), 3.66 (s, 6H, 3,5-(OCH<sub>3</sub>)<sub>2</sub>), 6.24 (m, 1H, CH benzene ring), 6.36 (m, 2H, CH benzene ring), 7.19-7.21 (d, 2H, CH benzene ring), 7.29-7.31 (d, 2H, CH benzene ring).

**1-(4-((3,5-Dimethoxyphenyl)thio)phenyl)-*N*-methyl-1-phenylmethanamine (46d')**. Oil. Yield: 30 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 2.34 (s, 3H, NHCH<sub>3</sub>), 3.64 (s, 6H, 2 x OCH<sub>3</sub>), 4.61 (s, 1H, NHCHPh), 6.23 (m, 1H, CH benzene ring), 6.35 (m, 2H, CH benzene ring), 7.14-7.16 (m, 1H, CH benzene ring), 7.23-7.26 (m, 8H, CH benzene rings).

**4-Methoxy-*N*-(4-(1-(methylamino)ethyl)phenyl)aniline (46e')**. Oil. Yield: 57 %. <sup>1</sup>H-NMR (400 MHz; CDCl<sub>3</sub>) δ ppm: 1.26-1.28 (d, 3H, CHCH<sub>3</sub>), 2.25 (s, 3H, NHCH<sub>3</sub>), 3.47-3.54 (m, 1H, CHCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 5.34 (s, 1H, PhNHPh-OCH<sub>3</sub>), 6.75-6.83 (m, 4H, CH benzene rings), 6.98-7.00 (d, 2H, CH benzene ring), 7.06-7.08 (d, 2H, CH benzene ring).

**4-Methoxy-*N*-(4-(1-(methylamino)ethyl)phenyl)benzamide (46f')**. Recryst. Solvent: methanol. Yield: 88 %. mp: 220-221 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.21-1.23 (d, 3H, CHCH<sub>3</sub>), 2.12 (s, 3H, NHCH<sub>3</sub>), 3.51-3.53 (m, 1H, CHCH<sub>3</sub>), 3.84 (s, 3H, OCH<sub>3</sub>), 7.05-7.07 (s,



2H, *CH* benzene ring), 7.25-7.27 (d, 2H, *CH* benzene ring), 7.67-7.69 (d, 2H, *CH* benzene ring), 7.94-7.96 (d, 2H, *CH* benzene ring), 10.02 (s, 1H, CONH).

**General Procedure for the Synthesis of the 2-Substituted 4-Aminopyrimidin-5-Carbonitriles 47a-f.** The 4-amino-2-bromopyrimidine-5-carbonitrile (1 equiv, 1 mmol) and triethylamine (1.6 equiv, 1.6 mmol) were added to a solution of the amines **46** (1 equiv, 1 mmol) in dry 2-methoxyethanol. After stirring at room temperature for 2.5 h, the reaction was stopped, and the solvent evaporated. The residue was diluted with ethyl acetate and washed two times with potassium hydrogen sulfate 0.1 N. The aqueous layer was counterextracted with ethyl acetate and the combined organic phases were dried over sodium sulfate, filtered and evaporated under reduced pressure. The crude residue was finally purified by silica gel column chromatography eluting with a mixture chloroform/*n*-hexane to afford the desired intermediate compounds **47** as white solids.

**4-Amino-2-((1-benzylpiperidin-4-yl)amino)pyrimidine-5-carbonitrile (47a).** Recryst. Solvent: acetonitrile/methanol. Yield: 83 %. mp: 205-208 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.44-1.53 (m, 2H, 2 x *CH* piperidine ring), 1.74 (m, 2H, 2 x *CH* piperidine ring), 1.93-1.99 (m, 2H, 2 x *CH* piperidine ring), 2.77-2.79 (m, 2H, 2 x *CH* piperidine ring), 3.43-3.45 (m, 2H, NCH<sub>2</sub>Ph), 3.72 (br m, 1H, NHC<sub>4</sub>-*H*-piperidine ring), 7.02-7.49 (m, 8H, *CH* benzene ring, NH<sub>2</sub> and NH), 8.13 and 8.23 (two s, 1H, *CH* pyrimidine ring of two tautomers).

**4-Amino-2-(2-methylpiperidin-1-yl)pyrimidine-5-carbonitrile (47b).** Recryst. Solvent: methanol. Yield: 82 %. mp: 157-161 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.07-1.09 (d, 3H, CH<sub>3</sub>), 1.27-1.31 (m, 1H, *CH* piperidine ring), 1.54-1.68 (m, 5H, 5 x *CH* piperidine ring), 2.84-2.92 (t, 1H, *CH* piperidine ring), 4.57-4.60 (m, 1H, *CH* piperidine ring), 5.02-5.04 (m, 1H, *CH* piperidine ring), 7.17 (br s, 2H, NH<sub>2</sub>), 8.24 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-(3,4-dihydroisoquinolin-2(1H)-yl)pyrimidine-5-carbonitrile (47c).** Recryst.

Solvent: acetonitrile. Yield: 88 %. mp: 186-189 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.83-2.86 (m, 2H, CH<sub>2</sub> tetrahydroisoquinoline ring), 3.97 (m, 2H, CH<sub>2</sub> tetrahydroisoquinoline ring), 4.87 (s, 2H, CH<sub>2</sub> tetrahydroisoquinoline ring), 7.19 (m, 4H, CH tetrahydroisoquinoline ring), 7.33 (br s, 2H, NH<sub>2</sub>), 8.31 (s, 1H, CH pyrimidine ring).

**4-Amino-2-(4-phenylpiperazin-1-yl)pyrimidine-5-carbonitrile (47d).** Recryst. Solvent:

methanol. Yield: 78 %. mp: 272-277 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 3.16-3.18 (m, 4H, 2 x CH<sub>2</sub> piperazine ring), 3.89 (m, 4H, 2 x CH<sub>2</sub> piperazine ring), 6.79-6.87 (m, 1H, CH benzene ring), 6.97-6.99 (m, 2H, CH benzene ring), 7.22-7.25 (m, 2H, CH benzene ring), 7.32 (br s, 2H, NH<sub>2</sub>), 8.30 (s, 1H, CH pyrimidine ring).

**4-Amino-2-((4-methoxybenzyl)amino)pyrimidine-5-carbonitrile (47e).** Recryst. Solvent:

methanol. Yield: 62 %. mp: 228-230 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) (mixture of two tautomers 60:40) δ ppm: 3.72 (s, 3H, OCH<sub>3</sub>), 4.37-4.42 (m, 2H, NHCH<sub>2</sub>Ph-OCH<sub>3</sub>), 6.85-6.87 (m, 2H, CH benzene ring), 7.12-7.24 (m, 4H, CH benzene ring and NH<sub>2</sub>), 7.81 and 7.97 (two t, 1H, NHCH<sub>2</sub>), 8.17 and 8.24 (two s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-phenylethyl)amino)pyrimidine-5-carbonitrile (47f).** Recryst. Solvent: toluene.

Yield: 94 %. mp: 130-132 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) (mixture of two tautomers 60:40) δ ppm: 1.41- (d, 3H, CHCH<sub>3</sub>), 5.12-5.16 (m, 1H, CHCH<sub>3</sub>), 7.20-7.36 (m, 7H, CH benzene ring and NH<sub>2</sub>), 7.81 and 8.05 (two d, 1H, NH), 8.16 (s, 1H, CH pyrimidine ring).

**4-Amino-2-(benzyl(methyl)amino)pyrimidine-5-carbonitrile (47g).** Recryst. Solvent:

acetonitrile. Yield: 91 %. mp: 187-187 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 3.04 (br s, 3H, NCH<sub>3</sub>), 4.84 (s, 2H, NCH<sub>2</sub>Ph), 7.21-7.34 (m, 7H, CH benzene ring and NH<sub>2</sub>), 8.29 (s, 1H, CH pyrimidine ring).

**4-Amino-2-(methyl(1-phenylethyl)amino)pyrimidine-5-carbonitrile (47h).** Recryst. Solvent: acetonitrile. Yield: 86 %. mp: 189-191 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.51 (d, 3H, CHCH<sub>3</sub>), 2.77 (s, 3H, NCH<sub>3</sub>), 6.21 (br m, 1H, CHCH<sub>3</sub>), 7.26-7.35 (m, 7H, CH benzene ring and NH<sub>2</sub>), 8.30 (s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-(4-methoxyphenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47i).** Recryst. Solvent: acetonitrile. Yield: 91 %. mp: 153-154 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.47 (d, 3H, CHCH<sub>3</sub>), 2.74 (s, 3H, NCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.16 (br m, 1H, CHCH<sub>3</sub>), 6.89-6.91 (d, 2H, CH benzene ring), 7.19-7.26 (m, 4H, CH benzene ring and NH<sub>2</sub>), 8.30 (s, 1H, CH pyrimidine ring).

**2-((1-([1,1'-Biphenyl]-4-yl)ethyl)(methyl)amino)-4-aminopyrimidine-5-carbonitrile (47j).** Recryst. Solvent: acetonitrile. Yield: 89 %. mp: 148-156 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.56 (d, 3H, CHCH<sub>3</sub>), 2.83 (s, 3H, NCH<sub>3</sub>), 6.25 (br m, 1H, CHCH<sub>3</sub>), 7.29-7.38 (m, 5H, CH biphenyl ring and NH<sub>2</sub>), 7.44-7.48 (m, 2H, CH biphenyl ring), 7.63-7.66 (m, 4H, CH biphenyl ring), 8.32 (s, 1H, CH pyrimidine ring).

**4-Amino-2-(methyl(1-(4-phenoxyphenyl)ethyl)amino)pyrimidine-5-carbonitrile (47k).** Recryst. Solvent: toluene. Yield: 85 %. mp: 147-150 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.50 (d, 3H, CHCH<sub>3</sub>), 2.79 (s, 3H, NCH<sub>3</sub>), 6.20 (br m, 1H, CHCH<sub>3</sub>), 6.97-7.01 (m, 4H, CH benzene rings), 7.13 (m, 1H, CH benzene ring), 7.28 (m, 4H, CH benzene rings and NH<sub>2</sub>), 7.36-7.40 (m, 2H, CH benzene rings), 8.30 (s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-(4-(4-methoxyphenoxy)phenyl)ethyl)amino)pyrimidine-5-carbonitrile (47l).** Recryst. Solvent: acetonitrile/methanol. Yield: 75 %. mp: 193-196 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) (mixture of two tautomers 60:40) δ ppm: 1.39-1.41 (d, 3H, CHCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 5.08-5.14 (two m, 1H, CHCH<sub>3</sub>), 6.84-6.86 (m, 2H, CH benzene ring), 6.93-6.99 (m, 4H, CH

benzene ring), 7.05 and 7.18 (two br s, 2H,  $NH_2$ ), 7.29-7.34 (m, 2H,  $CH$  benzene ring), 7.82 and 8.02 (two br m, 1H,  $NHCHCH_3$ ), 8.16 and 8.20 (two s, 1H,  $CH$  pyrimidine ring).

**4-Amino-2-((4-(4-methoxyphenoxy)benzyl)(methyl)amino)pyrimidine-5-carbonitrile (47m).**

Recryst. Solvent: acetonitrile. Yield: 80 %. mp: 188-189 °C.  $^1H$ -NMR (400 MHz;  $d_6$ -DMSO) (mixture of two rotamers 50:50)  $\delta$  ppm: 3.02 and 3.05 (two s, 3H,  $NCH_3$ ), 3.74 (s, 3H,  $OCH_3$ ), 4.77 and 4.81 (two s, 2H,  $NCH_2Ph$ ), 6.86-6.88 (d, 2H,  $CH$  benzene ring), 6.94-7.00 (m, 4H,  $CH$  benzene rings), 7.18-7.26 (m, 4H,  $CH$  benzene ring and  $NH_2$ ), 8.29 (s, 1H,  $CH$  pyrimidine ring).

**4-Amino-2-((1-(4-(4-methoxyphenoxy)phenyl)ethyl)(methyl)amino)pyrimidine-5-**

**carbonitrile (47n).** Recryst. Solvent: toluene. Yield: 81 %. mp: 145-149 °C.  $^1H$ -NMR (400 MHz;  $d_6$ -DMSO)  $\delta$  ppm: 1.49 (d, 3H,  $CHCH_3$ ), 2.77 (br s, 3H,  $NCH_3$ ), 3.75 (s, 3H,  $OCH_3$ ), 6.17 (br m, 1H,  $CHCH_3$ ), 6.88-6.90 (d, 2H,  $CH$  benzene ring), 6.94-7.00 (m, 4H,  $CH$  benzene rings), 7.26-7.31 (m, 4H,  $CH$  benzene ring and  $NH_2$ ), 8.32 (s, 1H,  $CH$  pyrimidine ring).

**4-Amino-2-((1-(4-(4-methoxyphenoxy)phenyl)propyl)(methyl)amino)pyrimidine-5-**

**carbonitrile (47o).** Recryst. Solvent: cyclohexane. Yield: 53 %. mp: 55-58 °C.  $^1H$ -NMR (400 MHz;  $d_6$ -DMSO) (mixture of two rotamers 50:50)  $\delta$  ppm: 0.82-0.86 (t, 3H,  $CHCH_2CH_3$ ), 1.91 and 2.06 (two br m, 2H,  $CHCH_2CH_3$ ), 2.75 and 2.81 (two br s, 3H,  $NCH_3$ ), 3.75 (s, 3H,  $OCH_3$ ), 5.93 and 6.01 (two br m, 1H,  $CHCH_2CH_3$ ), 6.86-6.88 (d, 2H,  $CH$  benzene ring), 6.94-7.00 (m, 4H,  $CH$  benzene rings), 7.26-7.34 (m, 4H,  $CH$  benzene ring and  $NH_2$ ), 8.30 (s, 1H,  $CH$  pyrimidine ring).

**4-Amino-2-((1-(4-(4-methoxyphenoxy)phenyl)-2-methylpropyl)(methyl)amino)pyrimidine-**

**5-carbonitrile (47p).** Recryst. Solvent: cyclohexane. Yield: 37 %. mp: 58-65 °C.  $^1H$ -NMR (400 MHz;  $d_6$ -DMSO) (mixture of two rotamers 50:50)  $\delta$  ppm: 0.80 and 0.89 (two m, 6H,  $CHCH(CH_3)_2$ ), 1.09 and 1.24 (two br m, 1H,  $CHCH(CH_3)_2$ ), 2.79 and 2.86 (two br s, 3H,  $NCH_3$ ), 3.92 (s, 3H,  $OCH_3$ ), 5.58 and 5.73 (two br m, 1H,  $CHCH(CH_3)_2$ ), 6.86-6.88 (m, 2H,  $CH$

benzene ring), 6.94-7.01 (m, 4H, *CH* benzene rings), 7.22 (br m, 2H, *NH*<sub>2</sub>), 7.34 and 7.43 (two m, 2H, *CH* benzene ring), 8.24 and 8.32 (two s, 1H, *CH* pyrimidine ring).

**4-Amino-2-(((4-(4-methoxyphenoxy)phenyl)(phenyl)methyl)(methyl)amino)pyrimidine-5-carbonitrile (47q).** Recryst. Solvent: cyclohexane. Yield: 45 %. mp: 60-66 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.84 (s, 3H, *NCH*<sub>3</sub>), 3.75 (s, 3H, *OCH*<sub>3</sub>), 6.91-6.93 (m, 2H, *CH* benzene ring), 6.96-6.99 (m, 2H, *CH* benzene ring), 7.01-7.04 (m, 2H, *CH* benzene ring), 7.14-7.16 (m, 4H, *CH* benzene rings and *NH*<sub>2</sub>), 7.31-7.40 (m, 6H, *CH* benzene rings and *CHPh*), 8.32 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-((1-(4-(3-methoxyphenoxy)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47r).** Recryst. Solvent: cyclohexane. Yield: 86 %. mp: 45-41 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.50-1.52 (d, 3H, *CHCH*<sub>3</sub>), 2.79 (s, 3H, *NCH*<sub>3</sub>), 3.73 (s, 3H, *OCH*<sub>3</sub>), 6.20 (br m, 1H, *CHCH*<sub>3</sub>), 6.51-6.58 (m, 2H, *CH* benzene ring), 6.70-6.73 (m, 1H, *CH* benzene ring), 6.98-7.00 (m, 2H, *CH* benzene ring), 7.25-7.29 (m, 5H, *CH* benzene rings and *NH*<sub>2</sub>), 8.31 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-((1-(4-(3,4-dimethoxyphenoxy)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47s).** Recryst. Solvent: cyclohexane. Yield: 90 %. mp: 61-63 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.48-1.50 (d, 3H, *CHCH*<sub>3</sub>), 2.77 (br s, 3H, *NCH*<sub>3</sub>), 3.72 (s, 3H, *OCH*<sub>3</sub>), 3.74 (s, 3H, *OCH*<sub>3</sub>), 6.17 (br m, 1H, *CHCH*<sub>3</sub>), 6.50-6.53 (m, 1H, *CH* benzene ring), 6.74 (s, 1H, *CH* benzene ring), 6.89-6.95 (m, 3H, *CH* benzene rings), 7.25 (m, 4H, *CH* benzene rings and *NH*<sub>2</sub>), 8.30 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-((1-(4-(3,5-dimethoxyphenoxy)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47t).** Recryst. Solvent: cyclohexane. Yield: 65 %. mp: 60-63 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.51 (d, 3H, *CHCH*<sub>3</sub>), 2.79 (br s, 3H, *NCH*<sub>3</sub>), 3.70 (s, 6H, 3,5 (*OCH*<sub>3</sub>)<sub>2</sub>),

6.12-6.29 (m, 4H, *CH* benzene ring and *CHCH*<sub>3</sub>), 6.99-7.01 (m, 2H, *CH* benzene ring), 7.28 (m, 4H, *CH* benzene ring and *NH*<sub>2</sub>), 8.30 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-(((4-(3,5-dimethoxyphenoxy)phenyl)(phenyl)methyl)(methyl)amino)pyrimidine-5-carbonitrile (47u).** Recryst. Solvent: cyclohexane. Yield: 54 %. mp: 54-58 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.85 (s, 3H, *NCH*<sub>3</sub>), 3.71 (s, 6H, 3,5 (*OCH*<sub>3</sub>)<sub>2</sub>), 6.16 (s, 2H, *CH* benzene ring), 6.30 (t, 1H, *CH* benzene ring), 7.01-7.03 (m, 2H, *CH* benzene ring), 7.16 (m, 4H, *CH* benzene rings and *NH*<sub>2</sub>), 7.32-7.40 (m, 6H, *CH* benzene rings and *CHPh*), 8.31 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-(methyl(1-(4-(3,4,5-trimethoxyphenoxy)phenyl)ethyl)amino)pyrimidine-5-carbonitrile (47v).** Recryst. Solvent: cyclohexane. Yield: 56 %. mp: 65-67 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.50 (d, 3H, *CHCH*<sub>3</sub>), 2.77 (s, 3H, *NCH*<sub>3</sub>), 3.64 (s, 3H, 4-*OCH*<sub>3</sub>), 3.71 (s, 6H, 3,5 (*OCH*<sub>3</sub>)<sub>2</sub>), 6.17 (br m, 1H, *CHCH*<sub>3</sub>), 6.37 (s, 2H, *CH* benzene ring), 6.94-6.96 (m, 2H, *CH* benzene ring), 7.27 (m, 4H, *CH* benzene ring and *NH*<sub>2</sub>), 8.30 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-(methyl(1-(4-(4-(trifluoromethoxy)phenoxy)phenyl)ethyl)amino)pyrimidine-5-carbonitrile (47w).** Recryst. Solvent: cyclohexane. Yield: 65 %. mp: 97-98 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.51-1.53 (d, 3H, *CHCH*<sub>3</sub>), 2.80 (br s, 3H, *NCH*<sub>3</sub>), 6.21 (br m, 1H, *CHCH*<sub>3</sub>), 7.03-7.05 (d, 2H, *CH* benzene ring), 7.08-7.10 (d, 2H, *CH* benzene ring), 7.31 (m, 4H, *CH* benzene ring and *NH*<sub>2</sub>), 7.37-7.39 (d, 2H, *CH* benzene ring), 8.31 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-(methyl(1-(4-(4-(methylthio)phenoxy)phenyl)ethyl)amino)pyrimidine-5-carbonitrile (47x).** Recryst. Solvent: cyclohexane. Yield: 84 %. mp: 48-50 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.49-1.51 (d, 3H, *CHCH*<sub>3</sub>), 2.45 (s, 3H, *SCH*<sub>3</sub>), 2.78 (br s, 3H, *NCH*<sub>3</sub>), 6.19 (br m, 1H, *CHCH*<sub>3</sub>), 6.95-6.98 (m, 4H, *CH* benzene rings), 7.27-7.30 (m, 6H, *CH* benzene rings and *NH*<sub>2</sub>), 8.30 (s, 1H, *CH* pyrimidine ring).

**4-Amino-2-((1-(4-(4-(benzyloxy)phenoxy)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47y).** Recryst. Solvent: cyclohexane/toluene. Yield: 51 %. mp: 109-110 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.49 (d, 3H, CHCH<sub>3</sub>), 2.77 (br s, 3H, NCH<sub>3</sub>), 5.08 (s, 2H, OCH<sub>2</sub>Ph), 6.18 (br m, 1H, CHCH<sub>3</sub>), 6.88-6.90 (d, 2H, CH benzene ring), 6.97-7.05 (m, 4H, CH benzene rings), 7.26 (m, 4H, CH benzene ring and NH<sub>2</sub>), 7.32-7.47 (m, 5H, CH benzene ring), 8.30 (s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-(4-((4-methoxynaphthalen-1-yl)oxy)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47z).** Recryst. Solvent: cyclohexane. Yield: 49 %. mp: 82-83 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.54 (d, 3H, CHCH<sub>3</sub>), 2.84 (br s, 3H, NCH<sub>3</sub>), 4.04 (s, 3H, OCH<sub>3</sub>), 6.22 (br m, 1H, CHCH<sub>3</sub>), 6.95-6.97 (d, 2H, CH naphthalene ring), 7.00-7.02 (m, 1H, CH naphthalene ring), 7.13-7.15 (m, 1H, CH naphthalene ring), 7.30 (br m, 4H, CH benzene ring and NH<sub>2</sub>), 7.59-7.64 (m, 2H, CH benzene ring), 7.93-7.95 (m, 1H, CH naphthalene ring), 8.24-8.27 (m, 1H, CH naphthalene ring), 8.35 (s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-(4-((4-methoxybenzyl)oxy)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47a').** Recryst. Solvent: acetonitrile/methanol. Yield: 52 %. mp: 193-194 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.46-1.48 (d, 3H, CHCH<sub>3</sub>), 2.74 (br s, 3H, NCH<sub>3</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 4.99 (s, 2H, OCH<sub>2</sub>Ph), 6.15 (br m, 1H, CHCH<sub>3</sub>), 6.93-6.97 (m, 4H, CH benzene rings), 7.18 (m, 4H, CH benzene ring and NH<sub>2</sub>), 7.36-7.38 (m, 2H, CH benzene ring), 8.30 (s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-(4-((4-methoxyphenyl)thio)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47b').** Recryst. Solvent: acetonitrile. Yield: 91 %. mp: 167-168 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.46-1.48 (d, 3H, CHCH<sub>3</sub>), 2.76 (br s, 3H, NCH<sub>3</sub>), 3.78 (s, 3H, OCH<sub>3</sub>), 6.13 (br m, 1H, CHCH<sub>3</sub>), 6.98-7.02 (d, 2H, CH benzene ring), 7.10-7.12 (m, 2H, CH benzene

ring), 7.26 (m, 4H, CH benzene ring and NH<sub>2</sub>), 7.38-7.42 (m, 2H, CH benzene ring), 8.29 (s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-(4-((3,5-dimethoxyphenyl)thio)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47c')**. Recryst. Solvent: cyclohexane. Yield: 81 %. mp: 48-50 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.52 (d, 3H, CHCH<sub>3</sub>), 2.79 (br s, 3H, NCH<sub>3</sub>), 3.79 (s, 6H, 3,5-(OCH<sub>3</sub>)<sub>2</sub>), 6.19 (br m, 1H, CHCH<sub>3</sub>), 6.37 (s, 2H, CH benzene ring), 6.41 (s, 1H, CH benzene ring), 7.29-7.31 (br m, 4H, CH benzene ring and NH<sub>2</sub>), 7.35-7.38 (m, 2H, CH benzene ring), 8.30 (s, 1H, CH pyrimidine ring).

**4-Amino-2-(((4-((3,5-dimethoxyphenyl)thio)phenyl)(phenyl)methyl)(methyl)amino)pyrimidine-5-carbonitrile (47d')**. Recryst. Solvent: cyclohexane. Yield: 63 %. mp: 84-87 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 2.85 (br s, 3H, NCH<sub>3</sub>), 3.71 (s, 6H, 3,5-(OCH<sub>3</sub>)<sub>2</sub>), 6.43 (m, 3H, CH benzene ring), 7.17-7.19 (m, 4H, CH benzene ring and NH<sub>2</sub>), 7.31-7.41 (m, 8H, CH benzene rings and CHCH<sub>3</sub>), 8.32 (br s, 1H, CH pyrimidine ring).

**4-Amino-2-((1-(4-((4-methoxyphenyl)amino)phenyl)ethyl)(methyl)amino)pyrimidine-5-carbonitrile (47e')**. Recryst. Solvent: cyclohexane. Yield: 65 %. mp: 52-55 °C. <sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.45 (d, 3H, CHCH<sub>3</sub>), 2.74-2.76 (br s, 3H, NCH<sub>3</sub>), 3.71 (s, 3H, 4-OCH<sub>3</sub>), 6.12 (br m, 1H, CHCH<sub>3</sub>), 6.84-7.89 (m, 4H, CH benzene rings), 7.00-7.07 (m, 4H, CH benzene ring and NH<sub>2</sub>), 7.24 (m, 2H, CH benzene ring), 7.85 (s, 1H, NH), 8.29 (s, 1H, CH pyrimidine ring).

**N-(4-(1-((4-Amino-5-cyanopyrimidin-2-yl)(methyl)amino)ethyl)phenyl)-4-methoxybenzamide (47f')**. Recryst. Solvent: acetonitrile. Yield: 50 %. mp: 164-165 °C.

<sup>1</sup>H-NMR (400 MHz; *d*<sub>6</sub>-DMSO) δ ppm: 1.50 (d, 3H, CHCH<sub>3</sub>), 2.67 (br s, 3H, NCH<sub>3</sub>), 3.84 (s, 3H, 4-OCH<sub>3</sub>), 6.18-6.20 (br m, 1H, CHCH<sub>3</sub>), 7.05-7.07 (d, 2H, CH benzene ring), 7.25 (m, 4H,



CH benzene ring and NH<sub>2</sub>), 7.73-7.75 (d, 2H, CH benzene ring), 7.94-7.96 (d, 2H, CH benzene ring), 8.31 (s, 1H, CH pyrimidine ring), 10.11 (s, 1H, NHCO).

### ***In vitro* Antischistosomal effects of SmSirt2 inhibitors:**

#### **Parasite material and ethics statement**

A Puerto Rican strain of *Schistosoma mansoni* is maintained in the laboratory using albino *Biomphalaria glabrata* snails as intermediate host and *Mesocricetus auratus* (golden hamsters) as definitive host. Cercaria were released from infested snails and harvested on ice as described previously.<sup>69</sup> Schistosomula were prepared *in vitro* by mechanical transformation.<sup>69</sup> 8 weeks post infestation, *S. mansoni* adult worms were recovered from the hamster hepatic system by whole body perfusion with saline solution pumped through a perfusing needle placed in the left ventricle of the heart.<sup>70</sup> All animal experimentation was conducted in accordance with the European Convention for the Protection of Vertebrate Animals used for Experimental and other Scientific Purposes (ETS No 123, revised Appendix A) and was approved by the committee for ethics in animal experimentation of the Nord-Pas de Calais region (Authorization No. APAFIS#8289-2016122015127050V3) and the Pasteur Institute of Lille (Agreement No. B59-350009).

**Schistosomula viability.** 500 schistosomula were incubated at 37°C under a humid atmosphere containing 5% CO<sub>2</sub> during 72 H in a 24-well plate containing 1 mL of complete medium (M199 medium (Invitrogen) supplemented with penicillin (50 U/mL), streptomycin (50 µg/mL), gentamycin (15 µg/mL) and rifampicin (60 µg/mL) and 10% fetal calf serum (Gibco). Parasite death was evaluated by visual examination under a microscope 72 hours after the beginning of treatment using three major criteria: absence of motility, tegument defects, granular appearance. For each condition, we observed a minimum of 300 larvae in order to determine the ratio of dead

larvae to total larvae. Moreover, for each condition, two different assays were performed and two independent batches of schistosomula (biological replicates) were used. *SmSirt2* inhibitors were dissolved in DMSO and two different concentrations (10 and 20  $\mu\text{M}$ ) were used (single dose at D0).

**Adult worm pairing stability and egg laying.** Ten pairs of *S. mansoni* adult worms were maintained in culture for 72 h in a 5%  $\text{CO}_2$  atmosphere at 37°C in a 6 well-plate containing 4 mL of complete medium in the presence of *SmSirt2* inhibitors at 10 and 20  $\mu\text{M}$  final concentration. Every day, the number of paired couples was evaluated by visual examination. At the end of the experiment, medium containing eggs was harvested and the total number of eggs was determined after centrifugation by microscopy. Two different assays were performed for each condition and repeated with two independent biological replicates.

**Cell proliferation assay.** HL-60 cells (grown in RPMI 1640 supplemented with 10% fetal bovine serum) were incubated in 96 well-tissue culture plates (density of 5000 per well) with *SmSirt2* inhibitors at 10  $\mu\text{M}$  final concentration or DMSO vehicle as control, in a total volume of 100  $\mu\text{L}$  for 72 hours at 37 °C; three replicates per concentration were used. Growth inhibition was determined using the CellTiter 96<sup>®</sup> AQueous Non-Radioactive Cell Proliferation Assay according to the manufacturer's instructions. Data was plotted as absorbance units against compound concentration using GraphPad Prism 7.0.

### Calculation of molecular properties

Data in Table S6 was calculated using DataWarrior (Version 5.0, <http://www.openmolecules.org/datawarrior/>).<sup>71</sup>

## ANCILLARY INFORMATION

**Acknowledgments.** This work was supported by Deutsche Forschungsgemeinschaft (DFG, GRK1976, to D.M., N.W. and M.J., testing on human Sirtuins Ju295/14-1), Italian PRIN 2015 (prot. 20152TE5PK to A.M.), AIRC 2016 (n. 19162 to A.M.), Progetto Ateneo Sapienza 2017 (to D.R.), the A-ParaDDisE program funded under the European Union's Seventh Framework Programme (grant agreement no. 602080 to M.J., C.R., R.P., A.M.) This study was supported by the grant ANR-10-LABX-0030-INRT, a French State fund managed by the Agence Nationale de la Recherche under the frame program Investissements d'Avenir ANR-10-IDEX-0002-02 to C.R., M.M. and E.R-M. J.L. and R.J.P. are supported by institutional funds from the Centre National de la Recherche Scientifique (CNRS), the Institut National de la Sante et de la Recherche, Medicale (INSERM), the Institut Pasteur de Lille and the Universite de Lille. We thank GSK for kindly donating the Kineto Boxes compounds (aka TCKAS) for biological testing and the COST action CM1406 (Epigenetic Chemical Biology) for support.

## ABBREVIATIONS USED

AMC, 7-amino-4-methylcoumarin; NA, nicotinamide; Z-Lys-OH, benzyloxycarbonyl lysine.

## SUPPORTING INFORMATION AVAILABILITY

Additional data for **1** and **3a-g** conversion by *SmSirt2* and *hSirt*,  $IC_{50}$  of NA with **1** and **3a-g** as substrates, source of intermediate compounds **45a-e**, **r**, **v**, **w** and **46a-h**. microscopy images of the effects of **7** and its analogues on schistosomula viability, **molecular properties of selected compounds** and HPLC data.

Molecular formula strings for the synthesized compounds are also available.

## REFERENCES

- (1) Gray, D. J.; Ross, A. G.; Li, Y. S.; McManus, D. P. Diagnosis and Management of

- Schistosomiasis. *Brit. Med. J.* **2011**, *342*, 1–12.
- (2) Cioli, D.; Pica-Mattoccia, L.; Basso, A.; Guidi, A. Schistosomiasis Control: Praziquantel Forever? *Mol. Biochem. Parasitol.* **2014**, *195*, 23–29.
- (3) Fenwick, A.; Webster, J. P.; Bosque-Oliva, E.; Blair, L.; Fleming, F. M.; Zhang, Y.; Garba, A.; Stothard, J. R.; Gabrielli, A. F.; Clements, A. C. A.; et al. The Schistosomiasis Control Initiative (SCI): Rationale, Development and Implementation from 2002-2008. *Parasitology* **2009**, *136*, 1719–1730.
- (4) Webster, J. P.; Molyneux, D. H.; Hotez, P. J.; Fenwick, A. The Contribution of Mass Drug Administration to Global Health: Past, Present and Future. *Philos. Trans. R. Soc. B Biol. Sci.* **2014**, *369*, 20130434.
- (5) Cleland, C. R.; Tukahebwa, E. M.; Fenwick, A.; Blair, L. Mass Drug Administration with Praziquantel Reduces the Prevalence of *Schistosoma Mansoni* and Improves Liver Morbidity in Untreated Preschool Children. *Trans. R. Soc. Trop. Med. Hyg.* **2014**, *108*, 575–581.
- (6) Vale, N.; Gouveia, M. J.; Rinaldi, G.; Brindley, P. J.; Gärtner, F.; Da Costa, J. M. C. Praziquantel for Schistosomiasis: Single-Drug Metabolism Revisited, Mode of Action, and Resistance. *Antimicrob. Agents Chemother.* **2017**, *61*, e02582-16.
- (7) Da Silva, V. B. R.; Campos, B. R. K. L.; De Oliveira, J. F.; Decout, J. L.; Alves de Lima, M. D. C. Medicinal Chemistry of Antischistosomal Drugs: Praziquantel and Oxamniquine. *Bioorganic Med. Chem.* **2017**, *25*, 3259–3277.
- (8) Siqueira, L. D. P.; Fontes, D. A. F.; Aguilera, C. S. B.; Timóteo, T. R. R.; Ângelos, M. A.; Silva, L. C. P. B. B.; De Melo, C. G.; Rolim, L. A.; Da Silva, R. M. F.; Neto, P. J. R. Schistosomiasis: Drugs Used and Treatment Strategies. *Acta Trop.* **2017**, *176*, 179–187.
- (9) Pica-Mattoccia, L.; Doenhoff, M. J.; Valle, C.; Basso, A.; Troiani, A. R.; Liberti, P.;

- Festucci, A.; Guidi, A.; Cioli, D. Genetic Analysis of Decreased Praziquantel Sensitivity in a Laboratory Strain of *Schistosoma Mansoni*. *Acta Trop.* **2009**, *111*, 82–85.
- (10) Pierce, R. J.; Dubois-Abdeselem, F.; Caby, S.; Trolet, J.; Lancelot, J.; Oger, F.; Bertheaume, N.; Roger, E. Chromatin Regulation in Schistosomes and Histone Modifying Enzymes as Drug Targets. *Mem. Inst. Oswaldo Cruz* **2011**, *106*, 794–801.
- (11) Noël, F.; Mendonça-Silva, D. L.; Thibaut, J. P. B.; Lopes, D. V. S. Characterization of Two Classes of Benzodiazepine Binding Sites in *Schistosoma Mansoni*. *Parasitology* **2007**, *134*, 1003–1012.
- (12) Berriman, M.; Haas, B. J.; Loverde, P. T.; Wilson, R. A.; Dillon, G. P.; Cerqueira, G. C.; Mashiyama, S. T.; Al-Lazikani, B.; Andrade, L. F.; Ashton, P. D.; et al. The Genome of the Blood Fluke *Schistosoma Mansoni*. *Nature* **2009**, *460*, 352–358.
- (13) Dissous, C.; Grevelding, C. G. Piggy-Backing the Concept of Cancer Drugs for Schistosomiasis Treatment: A Tangible Perspective? *Trends Parasitol.* **2011**, *27*, 59–66.
- (14) Pierce, R.; Dubois-Abdeselem, F.; Lancelot, J.; Andrade, L.; Oliveira, G. Targeting Schistosome Histone Modifying Enzymes for Drug Development. *Curr. Pharm. Des.* **2012**, *18*, 3567–3578.
- (15) Hailu, G.; Robaa, D.; Forgione, M.; Sippl, W.; Rotili, D.; Mai, A. Lysine Deacetylase Inhibitors in Parasites: Past, Present, and Future Perspectives. *J. Med. Chem.* **2017**, *60*, 4780–4804.
- (16) Stolfi, D. A.; Marek, M.; Lancelot, J.; Hauser, A. T.; Walter, A.; Leproult, E.; Melesina, J.; Rumpf, T.; Wurtz, J. M.; Cavarelli, J.; et al. Molecular Basis for the Antiparasitic Activity of a Mercaptoacetamide Derivative That Inhibits Histone Deacetylase 8 (HDAC8) from the Human Pathogen *Schistosoma Mansoni*. *J. Mol. Biol.* **2014**, *426*, 3442–3453.
- (17) Simoben, C. V.; Robaa, D.; Chakrabarti, A.; Schmidtkunz, K.; Marek, M.; Lancelot, J.;

- Kannan, S.; Melesina, J.; Shaik, T. B.; Pierce, R. J.; et al. A Novel Class of Schistosoma Mansoni Histone Deacetylase 8 (HDAC8) Inhibitors Identified by Structure-Based Virtual Screening and in Vitro Testing. *Molecules* **2018**, *23*, 1–14.
- (18) Heimburg, T.; Chakrabarti, A.; Lancelot, J.; Marek, M.; Melesina, J.; Hauser, A. T.; Shaik, T. B.; Duclaud, S.; Robaa, D.; Erdmann, F.; et al. Structure-Based Design and Synthesis of Novel Inhibitors Targeting HDAC8 from Schistosoma Mansoni for the Treatment of Schistosomiasis. *J. Med. Chem.* **2016**, *59*, 2423–2435.
- (19) Peck, B.; Chen, C. Y.; Ho, K. K.; Di Fruscia, P.; Myatt, S. S.; Coombes, R. C.; Fuchter, M. J.; Hsiao, C. D.; Lam, E. W. F. SIRT Inhibitors Induce Cell Death and p53 Acetylation through Targeting Both SIRT1 and SIRT2. *Mol. Cancer Ther.* **2010**, *9*, 844–855.
- (20) Chakrabarty, S. P.; Saikumari, Y. K.; Bopanna, M. P.; Balaram, H. Biochemical Characterization of Plasmodium Falciparum Sir2, a NAD<sup>+</sup>-Dependent Deacetylase. *Mol. Biochem. Parasitol.* **2008**, *158*, 139–151.
- (21) Tonkin, C. J.; Carret, C. K.; Duraisingh, M. T.; Voss, T. S.; Ralph, S. A.; Hommel, M.; Duffy, M. F.; Da Silva, L. M.; Scherf, A.; Ivens, A.; et al. Sir2 Paralogues Cooperate to Regulate Virulence Genes and Antigenic Variation in Plasmodium Falciparum. *PLoS Biol.* **2009**, *7*, 0771–0788.
- (22) Moretti, N. S.; Augusto, L. D. S.; Clemente, T. M.; Antunes, R. P. P.; Yoshida, N.; Torrecilhas, A. C.; Cano, M. I. N.; Schenkman, S. Characterization of Trypanosoma Cruzi Sirtuins as Possible Drug Targets for Chagas Disease. *Antimicrob. Agents Chemother.* **2015**, *59*, 4669–4679.
- (23) Ritagliati, C.; Alonso, V. L.; Manarin, R.; Cribb, P.; Serra, E. C. Overexpression of Cytoplasmic TcSIR2RP1 and Mitochondrial TcSIR2RP3 Impacts on Trypanosoma Cruzi Growth and Cell Invasion. *PLoS Negl. Trop. Dis.* **2015**, *9*, 1–22.

- (24) Yahiaoui, B.; Taibi, A.; Ouaiissi, A. A Leishmania Major Protein with Extensive Homology to Silent Information Regulator 2 of *Saccharomyces Cerevisiae*. *Gene* **1996**, *169*, 115–118.
- (25) Vergnes, B.; Sereno, D.; Madjidian-Sereno, N.; Lemesre, J. L.; Ouaiissi, A. Cytoplasmic SIR2 Homologue Overexpression Promotes Survival of Leishmania Parasites by Preventing Programmed Cell Death. *Gene* **2002**, *296*, 139–150.
- (26) Tavares, J.; Ouaiissi, A.; Silva, A. M.; Lin, P. K. T.; Roy, N.; Cordeiro-da-Silva, A. Anti-Leishmanial Activity of the Bisnaphthalimidopropyl Derivatives. *Parasitol. Int.* **2012**, *61*, 360–363.
- (27) Lancelot, J.; Caby, S.; Dubois-Abdesselem, F.; Vanderstraete, M.; Trolet, J.; Oliveira, G.; Bracher, F.; Jung, M.; Pierce, R. J. Schistosoma Mansoni Sirtuins: Characterization and Potential as Chemotherapeutic Targets. *PLoS Negl. Trop. Dis.* **2013**, *7*, 1–13.
- (28) Grozinger, C. M.; Chao, E. D.; Blackwell, H. E.; Moazed, D.; Schreiber, S. L. Identification of a Class of Small Molecule Inhibitors of the Sirtuin Family of NAD-Dependent Deacetylases by Phenotypic Screening. *J. Biol. Chem.* **2001**, *276*, 38837–38843.
- (29) Lara, E.; Mai, A.; Calvanese, V.; Altucci, L.; Lopez-Nieva, P.; Martinez-Chantar, M. L.; Varela-Rey, M.; Rotili, D.; Nebbioso, A.; Ropero, S.; et al. Salermide, a Sirtuin Inhibitor with a Strong Cancer-Specific Proapoptotic Effect. *Oncogene* **2009**, *28*, 781–791.
- (30) Rotili, D.; Tarantino, D.; Nebbioso, A.; Paolini, C.; Huidobro, C.; Lara, E.; Mellini, P.; Lenoci, A.; Pezzi, R.; Botta, G.; et al. Discovery of Salermide-Related Sirtuin Inhibitors: Binding Mode Studies and Antiproliferative Effects in Cancer Cells Including Cancer Stem Cells. *J. Med. Chem.* **2012**, *55* (24), 10937–10947.
- (31) Wang, J.; Kim, T. H.; Ahn, M. Y.; Lee, J.; Jung, J. H.; Choi, W. S.; Lee, B. M.; Yoon, K.

- S.; Yoon, S.; Kim, H. S. Sirtinol, a Class III HDAC Inhibitor, Induces Apoptotic and Autophagic Cell Death in MCF-7 Human Breast Cancer Cells. *Int. J. Oncol.* **2012**, *41*, 1101–1109.
- (32) Pasco, M. Y.; Rotili, D.; Altucci, L.; Farina, F.; Rouleau, G. A.; Mai, A.; Néri, C. Characterization of Sirtuin Inhibitors in Nematodes Expressing a Muscular Dystrophy Protein Reveals Muscle Cell and Behavioral Protection by Specific Sirtinol Analogues. *J. Med. Chem.* **2010**, *53*, 1407–1411.
- (33) Du, J.; Zhou, Y.; Su, X.; Yu, J. J.; Khan, S.; Jiang, H.; Kim, J.; Woo, J.; Kim, J. H.; Choi, B. H.; et al. Sirt5 Is a NAD-Dependent Protein Lysine Demalonylase and Desuccinylase. *Science (80-. )*. **2011**, *334*, 806–809.
- (34) Feldman, J. L.; Dittenhafer-Reed, K. E.; Kudo, N.; Thelen, J. N.; Ito, A.; Yoshida, M.; Denu, J. M. Kinetic and Structural Basis for Acyl-Group Selectivity and NAD + Dependence in Sirtuin-Catalyzed Deacylation. *Biochemistry* **2015**, *54*, 3037–3050.
- (35) Jiang, H.; Khan, S.; Wang, Y.; Charron, G.; He, B.; Sebastian, C.; Du, J.; Kim, R.; Ge, E.; Mostoslavsky, R.; et al. SIRT6 Regulates TNF- $\alpha$  Secretion through Hydrolysis of Long-Chain Fatty Acyl Lysine. *Nature* **2013**, *496*, 110–113.
- (36) Olsen, C. A. An Update on Lysine Deacylases Targeting the Expanding “acylome.” *ChemMedChem* **2014**, *9*, 434–437.
- (37) Bao, X.; Wang, Y.; Li, X.; Li, X. M.; Liu, Z.; Yang, T.; Wong, C. F.; Zhang, J.; Hao, Q.; Li, X. D. Identification of “Erasers” for Lysine Crotonylated Histone Marks Using a Chemical Proteomics Approach. *Elife* **2014**, *3*, 1–18.
- (38) Tan, M.; Peng, C.; Anderson, K. A.; Chhoy, P.; Xie, Z.; Dai, L.; Park, J.; Chen, Y.; Huang, H.; Zhang, Y.; et al. Lysine Glutarylation Is a Protein Posttranslational Modification Regulated by SIRT5. *Cell Metab.* **2014**, *19*, 605–617.



- (39) Anderson, K. A.; Huynh, F. K.; Fisher-Wellman, K.; Stuart, J. D.; Peterson, B. S.; Douros, J. D.; Wagner, G. R.; Thompson, J. W.; Madsen, A. S.; Green, M. F.; et al. SIRT4 Is a Lysine Deacylase That Controls Leucine Metabolism and Insulin Secretion. *Cell Metab.* **2017**, *25*, 838–855.
- (40) Lin, H.; Su, X.; He, B. Protein Lysine Acylation and Cysteine Succination by Intermediates of Energy Metabolism. *Chem. Biol.* **2012**, *7*, 947–960.
- (41) Feldman, J. L.; Baeza, J.; Denu, J. M. Activation of the Protein Deacetylase SIRT6 by Long-Chain Fatty Acids and Widespread Deacylation by Mammalian Sirtuins. *J. Biol. Chem.* **2013**, *288*, 31350–31356.
- (42) Zhu A. Y., Zhou Y., Kahn S., Deitsch K. W., Hao Q., L. H. Plasmodium Falciparum Sir2A Preferentially Hydrolyzes Medium and Long Chain Fatty Acyl Lysine. *Chem. Biol.* **2012**, *7*, 155–159.
- (43) Schiedel, M.; Marek, M.; Lancelot, J.; Karaman, B.; Almlöf, I.; Schultz, J.; Sippl, W.; Pierce, R. J.; Romier, C.; Jung, M. Fluorescence-Based Screening Assays for the NAD<sup>+</sup>-Dependent Histone Deacetylase smSirt2 from Schistosoma Mansoni. *J. Biomol. Screen.* **2015**, *20*, 112–121.
- (44) Peña, I.; Pilar Manzano, M.; Cantizani, J.; Kessler, A.; Alonso-Padilla, J.; Bardera, A. I.; Alvarez, E.; Colmenarejo, G.; Cotillo, I.; Roquero, I.; et al. New Compound Sets Identified from High Throughput Phenotypic Screening against Three Kinetoplastid Parasites: An Open Resource. *Sci. Rep.* **2015**, *5*, 8771.
- (45) Roquis, D.; Lepesant, J. M. J.; Picard, M. A. L.; Freitag, M.; Parrinello, H.; Groth, M.; Emans, R.; Cosseau, C.; Grunau, C. The Epigenome of Schistosoma Mansoni Provides Insight about How Cercariae Poise Transcription until Infection. *PLoS Negl. Trop. Dis.* **2015**, *9*, 1–22.

- (46) Heltweg, B.; Trapp, J.; Jung, M. In Vitro Assays for the Determination of Histone Deacetylase Activity. *Methods* **2005**, *36*, 332–337.
- (47) Sauve, A. A.; Youn, D. Y. Sirtuins : NAD<sup>+</sup>-Dependent Deacetylase Mechanism and Regulation. *Curr. Opin. Chem. Biol.* **2012**, *16*, 535–543.
- (48) Lancelot, J.; Cabezas-Cruz, A.; Caby, S.; Marek, M.; Schultz, J.; Romier, C.; Sippl, W.; Jung, M.; Pierce, R. J. Schistosome Sirtuins as Drug Targets. *Future Med. Chem.* **2015**, *7*, 765–782.
- (49) North, B. J.; Marshall, B. L.; Borra, M. T.; Denu, J. M.; Verdin, E. The Human Sir2 Ortholog, SIRT2, Is an NAD<sup>+</sup>-Dependent Tubulin Deacetylase. *Mol. Cell* **2003**, *11* (2), 437–444.
- (50) Borra, M. T.; Langer, M. R.; Slama, J. T.; Denu, J. M. Substrate Specificity and Kinetic Mechanism of the Sir2 Family of NAD<sup>+</sup>-Dependent Histone/protein Deacetylases. *Biochemistry* **2004**, *43*, 9877–9887.
- (51) Rumpf, T.; Schiedel, M.; Karaman, B.; Roessler, C.; North, B. J.; Lehotzky, A.; Oláh, J.; Ladwein, K. I.; Schmidtkunz, K.; Gajer, M.; et al. Selective Sirt2 Inhibition by Ligand-Induced Rearrangement of the Active Site. *Nat. Commun.* **2015**, *6*, 6263.
- (52) Hoffmann, G.; Breitenbücher, F.; Schuler, M.; Ehrenhofer-Murray, A. E. A Novel Sirtuin 2 (SIRT2) Inhibitor with p53-Dependent pro-Apoptotic Activity in Non-Small Cell Lung Cancer. *J. Biol. Chem.* **2014**, *289*, 5208–5216.
- (53) Swyter, S.; Schiedel, M.; Monaldi, D.; Sippl, W.; Lehotzky, A.; Rumpf, T.; Ovádi, J.; Jung, M. New Chemical Tools for Probing Activity and Inhibition of the NAD<sup>+</sup>-Dependent Lysine Deacetylase Sirtuin 2. *Phil. Trans. R. Soc.* **2018**, *373*, 20170083.
- (54) Panic, G.; Vargas, M.; Scandale, I.; Keiser, J. Activity Profile of an FDA-Approved Compound Library against *Schistosoma Mansoni*. *PLoS Negl. Trop. Dis.* **2015**, *9*, 1–15.

- (55) Abdulla, M. H.; Ruelas, D. S.; Wolff, B.; Snedecor, J.; Lim, K. C.; Xu, F.; Renslo, A. R.; Williams, J.; McKerrow, J. H.; Caffrey, C. R. Drug Discovery for Schistosomiasis: Hit and Lead Compounds Identified in a Library of Known Drugs by Medium-Throughput Phenotypic Screening. *PLoS Negl. Trop. Dis.* **2009**, *3*, e478.
- (56) Schiedel, M.; Rumpf, T.; Karaman, B.; Lehotzky, A.; Oláh, J.; Gerhardt, S.; Ovádi, J.; Sippl, W.; Einsle, O.; Jung, M. Aminothiazoles as Potent and Selective Sirt2 Inhibitors: A Structure-Activity Relationship Study. *J. Med. Chem.* **2016**, *59*, 1599–1612.
- (57) Green, B. T.; Lee, S. T.; Welch, K. D.; Panter, K. E. Plant Alkaloids That Cause Developmental Defects through the Disruption of Cholinergic Neurotransmission. *Birth Defects Res. Part C - Embryo Today Rev.* **2013**, *99*, 235–246.
- (58) Barradas, S.; Hern, G.; Urbano, A.; Carre, M. C. Total Synthesis of Natural P - Quinol Cochinchinenone. *Org. Lett.* **2012**, *14*, 5952–5955.
- (59) Biscoe, M. R.; Fors, B. P.; Buchwald, S. L. A New Class of Easily Activated Palladium Precatalysts for Facile C - N Cross-Coupling Reactions and the Low Temperature Oxidative Addition of Aryl Chlorides. *J. Am. Chem. Soc.* **2008**, *130*, 6686–6687.
- (60) Vicker, N.; Xiangdong, S.; Ganeshpillai, D.; Purhoit, A.; Reed, M. J.; Potter, B. WO2005042513A1. 2005.
- (61) Spiegelman, N. A.; Price, I. R.; Jing, H.; Wang, M.; Yang, M.; Cao, J.; Hong, J. Y.; Zhang, X.; Aramsangtienchai, P.; Sadhukhan, S.; et al. Direct Comparison of SIRT2 Inhibitors: Potency, Specificity, Activity-Dependent Inhibition, and On-Target Anticancer Activities. *ChemMedChem* **2018**, 1–6.
- (62) Hamid, M. H. S. A.; Allen, C. L.; Gareth, W.; Maytum, H. C.; Maxwell, A. C.; Watson, A. J. A.; Williams, J. M. J. Ruthenium-Catalyzed Remote Electronic Activation of Aromatic. *2013*, *5*, 734–745.

- (63) Lovering, J. R.; Ridd, J. H.; Parker, D. G.; Rose, J. B. Polymerisation and Related Reactions Involving Nucleophilic Aromatic Substitution. Part 2.' The Rates of Reaction of Substituted 4-Halogenobenzophenones with the Salts of Substituted Hydroquinones. **1988**, No. 9, 1735–1738.
- (64) Zhao, Y.; Wang, Y.; Sun, H.; Li, L.; Zhang, H. Ullmann Reaction in Tetraethyl Orthosilicate : A Novel Synthesis of Triarylaminines and Diaryl Ethers. **2007**, 3186–3188.
- (65) Yang, Y.; Yu, Y.; Li, X.; Li, J.; Wu, Y.; Yu, J.; Ge, J.; Huang, Z.; Jiang, L.; Rao, Y.; et al. Target Elucidation by Co-Crystal Structures of NADH-Ubiquinone Oxidoreductase of Plasmodium Falciparum ( Pf NDH2 ) with Small Molecule to Eliminate Drug-Resistant Malaria. *J Med Chem* **2017**, 5 (60), 1994–2005.
- (66) Breyoltz, H.-J.; Schafers, M.; Wagner, S.; Holtke, C.; Faust, A.; Rabeneck, H.; Levkau, B.; Schober, O.; Kopka, K. C-5-Disubstituted Barbiturates as Potential Molecular Probes for Noninvasive Matrix Metalloproteinase Imaging. *J Med Chem* **2005**, 3450–3459.
- (67) Barradas, S.; Hern, G.; Urbano, A.; Carre, M. C. Total Synthesis of Natural p - Quinol Cochinchinenone. *Org. Lett.* **2012**, 14 (23), 5952–5955.
- (68) Wakchaure, V. N.; Kaib, P. S. J.; Leutzsch, M.; List, B. Disulfonimide-Catalyzed Asymmetric Reduction of *N*-Alkyl Imines. *Angew. Chem. Int. Ed. Engl.* **2015**, 11852–11856.
- (69) Ramalho-Pinto, F. J.; Gazzinelli, G.; Howells, R. E.; Mota-Santos, T. A.; Figueiredo, E. A.; Pellegrino, J. Schistosoma Mansoni: Defined System for Stepwise Transformation of Cercaria to Schistosomule in Vitro. *Exp. Parasitol.* **1974**, 36 (3), 360–372.
- (70) Smithers, S. R.; Terry, R. J. The Infection of Laboratory Hosts with Cercariae of Schistosoma Mansoni and the Recovery of the Adult Worms. *Parasitology* 1965, 55 (4), 695–700.

- (71) Thomas Sander, Joel Freyss, Modest von Korff, Christian Rufener. DataWarrior: An Open-Source Program For Chemistry Aware Data Visualization And Analysis. *J Chem Inf Model* **2015**, *55*, 460-473,

## Table of Contents graphic

