

# The Reaction of Sulfur Dioxide Radical Cation with Hydrogen and its Relevance in Solar Geoengineering Models

Mauro Satta\*, Antonella Cartoni\*, Daniele Catone, Mattea Carmen Castrovilli,  
Paola Bolognesi, Nicola Zema, Lorenzo Avaldi

## Abstract

*SO<sub>2</sub> has been proposed in solar geoengineering as a precursor of H<sub>2</sub>SO<sub>4</sub> aerosol, a cooling agent active in the stratosphere to contrast climate change. Atmospheric ionization sources can ionize SO<sub>2</sub> into excited states of SO<sub>2</sub><sup>+</sup>, quickly reacting with trace gases in the stratosphere. In this work we explore the reaction of H<sub>2</sub>(D<sub>2</sub>) with SO<sub>2</sub><sup>+</sup> excited by tunable synchrotron radiation, leading to HSO<sub>2</sub><sup>+</sup>+H (DSO<sub>2</sub><sup>+</sup>+D), where H contributes to O<sub>3</sub> depletion. Density Functional Theory and Variational Transition State Theory have been used to investigate the dynamics of the title reaction, which is barrierless and exothermic. The present results suggest that solar geoengineering models should test the reactivity of SO<sub>2</sub><sup>+</sup> with major trace gases in the stratosphere, such as H<sub>2</sub> and H<sub>2</sub>O, since these are relevant channels for the OH<sup>•</sup> formation during the nighttime when there is not OH<sup>•</sup> production by sunlight. OH<sup>•</sup> oxides SO<sub>2</sub>, starting the chem-*

*ical reactions leading to H<sub>2</sub>SO<sub>4</sub> aerosol.*

## Introduction

Technologies that aim to cool the planet by partially reflecting sunlight away from Earth back into space are the core of the solar geoengineering<sup>[1–5]</sup>. These technologies, which aim to manipulate the atmosphere to fight climate change, have triggered an active discussion<sup>[6–9]</sup> even at the level of the highest international institutions, such as the United Nations Environment Assembly<sup>[10,11]</sup>. The goal of solar geoengineering is to artificially reduce global temperatures by sunlight-reflecting particles generated in the atmosphere mainly by sulfur dioxide, which would be injected in the stratosphere to contrast global climate warming<sup>[8]</sup>. The idea is that this strategy should mimic the effects of the natural emission of sulfur dioxide by volcanic eruption. Nevertheless, several worries have been raised about the long-term consequences of these technologies on the ozone-layer<sup>[7]</sup>, on the unbalances of the water cycle, i.e. droughts and/or floods<sup>[12]</sup>, or even on the biodiversity threats<sup>[13]</sup>. Up to now, almost only theoretical models have been used to predict the right amount of the cooling agent necessary to produce a climate cooling without a severe irreversible alteration of the global climate<sup>[6]</sup>. The present models take into account several neutral molecules, their thermal reactions and photoinduced dissociation<sup>[14]</sup>. However, ion-molecule reactions may also play a relevant role in the chemical transformation of several species since,

---


Dr. M. Satta  
ISMN (CNR) c/o Dipartimento di Chimica Sapienza Università di Roma  
P.le Aldo Moro 5, Roma, Italy  
E-mail:mauro.satta@cnr.it  
Dr. A. Cartoni  
Dipartimento di Chimica, Sapienza Università di Roma,  
P.le Aldo Moro 5, Roma, Italy  
E-mail:antonella.cartoni@uniroma1.it  
Dr. D. Catone, Dr. N. Zema  
CNR-ISM, Area della Ricerca di Tor Vergata,  
Via del Fosso del Cavaliere, Roma, Italy  
Dr. M. C. Castrovilli, Dr. P. Bolognesi, Dr. L. Avaldi  
CNR-ISM, Area della Ricerca di Roma 1,  
Via Salaria Km 29,300, Monterotondo Scalo (RM), Italy

especially in the stratosphere, ionizing radiation are not negligible and much effort has been paid to determine the amount and nature of these radiation produced by cosmic rays and space weather events in the atmosphere<sup>[15–20]</sup>. Typical average values of the neutral/ion ratio during daytime at mid-latitudes in the stratosphere is about  $10^{12}$ , and it is independent by the type of molecular species because cosmic rays ionize all neutral molecules in the stratosphere with a similar efficiency<sup>[19]</sup>. This low ion/neutral ratio can be counterbalanced by the fact that ion-molecule reactions are generally much faster than the neutral-neutral reactions (i.e. the neutral chemistry involved in the conversion of  $SO_2$  into  $H_2SO_4$  takes approximately 30 days<sup>[21]</sup>), and can heavily alter the total budget of important molecular species in the atmosphere. Hence, reactions involving the radical cation  $SO_2^+$ , which can be produced in the stratosphere by ionizing radiation, should be considered in the chemistry models used when evaluating the climate impact of solar geoengineering involving sulfur dioxide as aerosol precursor. The role of the ionizing radiation on the  $SO_2$  geoengineering release in the stratosphere should be deeply considered also in view of the fact that during natural volcanic emission of sulfur dioxide large amount of particulates are also emitted. These can largely absorb the ionizing radiation, and thus reduce the natural production of  $SO_2^+$ . This radiation “shield” effect obviously will not act when only gaseous  $SO_2$  is injected by balloons or aircrafts to artificially produce aerosol particulates. This aspect should be considered when comparison between sulfur dioxide volcanic and geoengineering emission are discussed as similar processes.

Moreover, the non-thermal conditions, under which the ion-molecule reactions in the stratosphere can occur, should be also taken into account. The ionizing radiation produces highly excited ions, which slowly thermalize with the surrounding molecular species in equilibrium at temperatures between 200K and 250K depending on the altitude of the stratosphere<sup>[22–25]</sup>. These non-thermal effects are generally amplified for reactions between hot molecular ions and rigid neutral species with high vibrational frequencies<sup>[26,27]</sup>. Hence, we have explored the reaction between the radical cation  $SO_2^+$  and the  $H_2$

molecule (eq 1), a major trace gas in the stratosphere (about 0.5 ppmv<sup>[28]</sup>) with almost constant concentration with the altitude.

This work follows recent studies on ion-molecule reactions<sup>[29–32]</sup>, and on the dynamic of the Hydrogen Atom Transfer (HAT) reaction of sulfur dioxide radical cation  $SO_2^+$  with methane and water leading to  $HSO_2^+$  ions<sup>[33]</sup> and  $CH_3$  or  $OH$ , which is a relevant oxidant in the environmental chemistry<sup>[34]</sup>. The ionic product of these reactions,  $HSO_2^+$  has been detected in the terrestrial stratosphere with other sulfur oxides protonated species<sup>[35]</sup> and reacts with water via proton transfer<sup>[33]</sup>. The title reaction (eq 1) has been studied experimentally using tunable synchrotron radiation to produce  $SO_2^+$  in the excited ro-vibrational levels of the ionic ground state<sup>[33]</sup>:



In the stratosphere one of main chemical process involving  $H$  (eq 1) is the reaction with ozone leading to molecular oxygen and hydroxyl radical:



Eq 2 has a rate coefficient of  $2.89 \cdot 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ <sup>[14]</sup>, and it is both a chemical route for the destruction of ozone and a source of  $OH$ . Eq 1, together with the efficient reaction of  $SO_2^+$  with water<sup>[33]</sup>, is a nighttime alternative pathway to the sunlight formation of  $OH$ <sup>[14]</sup>. The latter can oxidize  $SO_2$ , and trigger the chemical reactions network leading to the formation of  $H_2SO_4$  aerosol in the stratosphere, which in turn could modify the expected size distribution of the particulates with consequences on the global expected cooling effects. It is noteworthy that another largely studied oxidant species in the troposphere are the Criegee Intermediates (CI)<sup>[36]</sup>. However, in the stratosphere the role of the CI in the oxidation chemistry is not so relevant due to the hereby low concentration of organic compounds, and to the low pressures that hinder the stabilization of the CI<sup>[37]</sup>.

The experiments performed in this work have been supported by a theoretical exploration of the reaction dynamics on the potential energy surface with Density Functional Theory (DFT) and Variational

Transition State Theory (VTST)<sup>[38,39]</sup>. The spin and charge effects, as well as ionization energy (IE), polarizability, binding energy, radical stability and ro-vibrational degrees of freedom along the reaction path, have been also considered to explain the dynamics behavior of the reaction. The same study has been repeated with the  $D_2$  molecule, and a comparison with the reaction of  $SO_2^+$  with water and methane, as well as other previous ion-molecule reaction studies of  $SO_2^+$  and  $CH_4^+$  with hydrogen, has been carried out<sup>[33,40,41,26]</sup>. All along the manuscript the radical symbol in  $SO_2^+$  has been omitted for the sake of simplicity.

## Results and Discussion

### Synchrotron Experiments

The measurements were carried out at the “Circular Polarization” beamline (CiPo) of ELETTRA synchrotron (Trieste, Italy)<sup>[42]</sup> with the setup operated at the room temperature and described in the experimental section. The  $SO_2^+$  ions were obtained at low pressure (about  $10^{-6}$  mbar) from the ionization of an effusive molecular beam of  $SO_2$  by monochromatic synchrotron radiation tuned in the photon energy range 12.4-15.0 eV. The  $SO_2^+$  thus can be produced in the ground  $X^2A_1$  state (IE=12.349 eV) or in the two excited electronic states: the  $^2B_2$  (12.988 eV) and  $^2A_2$  (13.338 eV) ones, very close in energy<sup>[43,44]</sup>. However, as already well established<sup>[33]</sup>, the electronically excited states of  $SO_2^+$  ion decay to excited ro-vibrational states of the ground electronic state on a nanosecond timescale before the reaction with the neutral ( $H_2/D_2$ ) occurs (timescale fraction of ms). In this way the reacting  $SO_2^+$  ions are “hot”, with an internal energy increasing as the photoionizing energy is increased, and it is not in thermal equilibrium with the environment at room temperature. Moreover, the mass spectra acquired in the photon energy range 12.4-15.0 show that the  $SO_2^+$  ions do not dissociate, as also demonstrated by photoelectron-photoion coincidence (PEPICO) measurements<sup>[33,45]</sup>, and no reaction with the residual water in the octupole (base pressure about  $10^{-7}$  mbar) was observed. The  $SO_2^+$  ions were guided in the reaction zone (octupole) at




Figure 1: Mass spectra recorded at the photon energy 14.0 eV, nominal CE=0.0 eV and  $P = 5.1 \cdot 10^{-5}$  and  $8.0 \cdot 10^{-5}$  mbar for the reaction with  $H_2$  (black line) and  $D_2$  (red line), respectively.

the nominal Collision Energy (CE) of 0.0 eV with an estimated energy spread of 0.1 eV. Neutral gas  $H_2/D_2$  was introduced in the octupole at the room temperature via a controlled needle valve at the pressure of about  $10^{-5}$  mbar. The experiments were performed at different photon energy, pressure and CE by changing one parameter at the time. The ion yields of the ionic reagent and products were recorded and their ratio was analyzed. The mass spectra acquired at the photon energy 14.0 eV for the ion-molecule reaction between  $SO_2^+$  and  $H_2/D_2$  are shown in Figure 1.

The mass spectra were acquired at the pressure of  $5.1 \cdot 10^{-5}$  and  $8.0 \cdot 10^{-5}$  mbar for neutral  $H_2$  and  $D_2$ , respectively. In the case of the  $H_2$  as reagent (black line in Figure 1), a peak at  $m/z$  65 ( $HSO_2^+$ ) is observed with a signal of about 11% relative to the most intense peak  $SO_2^+$   $m/z$  64. In the case of reaction with deuterium (red line in Figure 1) a peak at  $m/z$  66 ( $DSO_2^+$ ) is observed with an intensity of about 8% of the most intense peak  $SO_2^+$  at  $m/z$  64 (see experimental section).

In Figure 2a the mass spectra of the  $SO_2^+ + H_2$  reaction, acquired at the photon energy of 14.0 eV and as a function of the  $H_2$  pressure are shown. The peaks intensity of the ions at  $m/z$  65 is normalized




Figure 2: a) The mass spectrum in the region of  $62 < m/z < 68$  acquired at the photon energy 14.0 eV, nominal CE=0.0 eV and pressures of  $H_2$ :  $1.1 \cdot 10^{-5}$  mbar (black line),  $2.6 \cdot 10^{-5}$  mbar (red line),  $5.1 \cdot 10^{-5}$  (green line);  $7.0 \cdot 10^{-5}$  (blue) and  $8.3 \cdot 10^{-5}$  (light blue) mbar. The intensities are normalized to the one of the reagent ion.  $SO_2$  pressure in the ion source was  $4.4 \cdot 10^{-6}$  mbar; b)  $HSO_2^+/SO_2^+$  ratio as a function of the  $H_2$  pressure.

with respect to the  $SO_2^+$  peak at  $m/z = 64$ . As expected an increase of  $m/z = 65$  signal is observed relative to all other peaks when the pressure varies from  $1.1 \cdot 10^{-5}$  to  $8.3 \cdot 10^{-5}$  mbar. The trend of the ion intensity ratio ( $HSO_2^+/SO_2^+$ ) vs nominal pressure of  $H_2$  (Figure 2b) demonstrates that the  $HSO_2^+$  is produced by reaction (1). Afterwards, the yields of  $SO_2^+$  and  $H(D)SO_2^+$  ions were acquired at fixed pressures of the neutral  $H_2/D_2$  and at the nominal CE=0.0 eV varying the photon energy from 12.4 to 15.0 eV.




Figure 3:  $H(D)SO_2^+/SO_2^+$  ratio as a function of photon energy in the reaction of  $SO_2^+$  with  $H_2$  (black line) and  $D_2$  (red line) at the fixed pressure of neutrals of  $8.3 \cdot 10^{-5}$  mbar and nominal CE=0.0 eV

The ratio  $H(D)SO_2^+/SO_2^+$  for both reactions at the nominal fixed pressure of about  $8.3 \cdot 10^{-5}$  mbar as a function of photon energy is reported in Figure 3.

The results clearly show that as the internal energy of  $SO_2^+$  increases the ratio  $H(D)SO_2^+/SO_2^+$  decreases. The observed isotopic effect demonstrates the main leading role of hydrogen in the reaction coordinate. Furthermore, the ratio  $HSO_2^+/SO_2^+$  decreases at increasing nominal CE from 0.0 to 1.2 eV at fixed photon energy (Figure S1 in SI).

Thermochemical literature data of reactants and products of reaction (1) show that the HAT reaction is exothermic by 75.0 kJ/mol<sup>[46]</sup>. The literature data on experimental rate coefficients are referred to different experimental mass spectrometric techniques: in Selected Ion Flow Tube Mass Spectrometry (SIFT-MS) experiments the values of  $4.2 \cdot 10^{-12} \pm 20\%$  and  $5.0 \cdot 10^{-12} \pm 20\%$   $\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$  were obtained<sup>[40,41]</sup> at 300 K, while a value of  $1.7 \cdot 10^{-11} \pm 40\%$   $\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$  was measured from the Fourier Transform Ion Cyclotron Resonance (FT-ICR) experiments at 298 K<sup>[47]</sup>. Although the values are quite different for SIFT and FT-ICR experiments, they are all lower than the Langevin rate coefficients<sup>[48,49]</sup> of  $1.50 \cdot 10^{-9}$  or  $1.53 \cdot 10^{-9}$   $\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$ . Actually,

	$H_2$	$H_2O$	$CH_4$
BDE (kJ/mol)	435.58	496.7	438.9
IE (eV)	15.42593	12.621	12.61
$\mu$ (D)	//	1.8550	//
Polarizability ( $\text{\AA}^3$ )	0.787	1.501	2.448

Table 1: Bond Dissociation Energy (BDE), Ionization Energy (IE), dipole moment and polarizability for  $H_2$ ,  $H_2O$  and  $CH_4$  molecules.<sup>[52]</sup>

many H-atom abstraction reactions involving  $H_2$  are very slow<sup>[47]</sup>. In the specific case of the reaction involving  $SO_2^+ + H_2$  a barrier of 19.9 kJ/mol has been found by early theoretical and experimental studies that account for the low rate coefficient at 300 K<sup>[40]</sup>. This evidence can not be only a consequence of the  $H-H$  binding energy since the HAT reactions of  $SO_2^+$  with  $CH_4$  and  $H_2O$  are faster even if the  $C-H$ ,  $O-H$ ,  $H-H$  molecular Bond Dissociation Energies (BDE) are comparable (Table 1)<sup>[50,51]</sup>.

### Minimum Energy Path (MEP)

In order to explore the main factors affecting the reaction dynamics of the HAT from  $H_2$  to  $SO_2^+$  a theoretical study has been carried out by combining DFT and VTST approaches<sup>[53]</sup>. DFT was used to investigate the MEP of the reaction and to identify possible transition states and minima along the reactive coordinate, while VTST has been used to search for the “bottleneck” of the reaction, namely the Variational Transition State (**VTS**) configuration.

In Figure 4 the MEP of the reaction is shown. The reaction is barrierless and exothermic by 75.5 kJ/mol at 300 K with a minimum energy reaction complex configuration  $[OSOH \cdots H]^+$  **Min** at 85.0 kJ/mol in cis conformation (See Table 2 for details on the geometric parameters of the molecular adducts whose atomic labels are defined in Figure 5).

The calculated exothermicity is in agreement with experimental value of 75.0 kJ/mol<sup>[46]</sup>, and no transition state configuration was found above the energy of the reactants. The barrierless nature of the reaction has been also confirmed by the higher level of calculations (see computational methods and SI). Furthermore, the VTST approach identifies a **VTS** con-




Figure 4: Minimum Energy Path for the  $SO_2^+ + H_2$  reaction. The minimum energy of the complex  $[OSOH \cdots H]^+$  is fixed at 0.0 kJ/mol. See the main text for further details.

figuration  $[O_dSO_cH_b \cdots H_a]^+$  along the MEP very close to **Min**, with similar geometry, charge and spin distribution (See Table 2 and Figure 5).

In both **VTS** and **Min** configurations the main bonds involved in the reaction,  $O_c - H_b$  (about 1.0 Å) and  $H_a - H_b$  (about 1.6 Å) are almost formed and fully stretched, respectively. In these configurations the charge ( $H_bSO_2^+$ : 0.90  $e$ ) and the spin ( $H_a$ : 0.85  $\hbar$ ) are almost fully localized as in the two final product moieties (see Figure 5). This dynamical picture seems very similar to that observed in the reaction of  $SO_2^+$  with  $H_2O$  (see Figure 5 of Carton et al.<sup>[33]</sup>). However, through a careful analysis of the reaction it can be seen that the **VTS** structure comes before the **Min** adduct in the

	Reagents	VTS	Min	Products
$R_{H_a-H_b}$	0.75	1.59	1.61	//
$R_{O_c-H_b}$	//	1.02	1.01	0.99
$R_{S-O_c}$	1.46	1.56	1.56	1.56
$R_{S-O_d}$	1.46	1.44	1.44	1.44
$\theta_{O_cSO_d}$	130.4	115.1	115.1	114.9
$\theta_{SO_cH_b}$	//	122.4	122.4	121.2
$\theta_{O_cH_bH_a}$	//	171.2	171.2	//

Table 2: Geometrical parameters for the main molecular species relevant to the title reaction. Distances in Å and angles in degrees.




Figure 5: Mulliken Atomic Charge (Top panel) and Spin population (Bottom panel) along the MEP in the region of the reaction. See further details in the main text.

reaction coordinate (Figure 4), and the dynamical evolution of the charge and spin distribution (Figure 5) in the complex along the reaction path is very different from that of the reaction of  $SO_2^+$  with water<sup>[33]</sup>. As for the partial charges and spin in the reaction complex  $[O_dSO_c \cdots H_bH_a]^+$  in the region of approaching reagents, the charge and the spin remain mainly localized on  $SO_2^+$ . Moving forward with the reaction up to  $R_{OH} = 1.5$  Å the charge increases on  $H_a$  from 0.0 to  $+0.3 e$  and decreases on  $SO_2^+$  from 1.0 to  $0.7 e$  and on  $H_b$  from 0.1 to  $0.0 e$

(see top panel of Figure 5). The spin remains almost constant on  $H_b$ , but it decreases for  $SO_2^+$  from 1.0 to  $0.5 h$ , and increases for  $H_a$  from 0.0 to  $0.4 h$  (see bottom panel of Figure 5). On the other hand, in the region of the **Min** the charge decreases on  $H_a$  and increase on  $H_b$ , while the spin goes to zero for  $SO_2^+$ , and increases for  $H_a$  up to  $1.0 h$ . It is interesting to note that, in the reaction of  $SO_2^+$  with water, the chemical rearrangement occurs at the beginning of the reaction with a fast proton-coupled electron transfer<sup>[33]</sup>. The higher IE 15.4 eV of  $H_2$  relative to IE 12.6 eV of  $H_2O$ , the lowest  $H_2$  polarizability (see Table 1) and the higher relative stability of OH ( $\Delta H_f^\circ = +38.99$  kJ/mol) with respect to H ( $\Delta H_f^\circ = +218.0$  kJ/mol) should make the reaction dynamics  $SO_2^+ + H_2$  more constrained and slower<sup>[52]</sup>. The “bottleneck” of the reaction’s trajectories, occurring at the **VTS** complex, reduces the reactivity below the collision limit of  $1.53 \cdot 10^{-9}$  cm<sup>3</sup>molecule<sup>-1</sup>s<sup>-1</sup>.

#### Theoretical rate coefficients

The VTST allows the calculation of the kinetic coefficients ( $k$ ) trough the following equation (see theoretical section for more details):

$$k(T) = \sigma \frac{k_B T}{h} \left[ \frac{Q_{\mathbf{VTS}}(T)}{Q_{SO_2^+}(T) Q_{H_2}(T)} \right] \quad (3)$$

which holds in thermal equilibrium conditions (Canonical ensemble). In eq 3  $\sigma$  is the symmetry factor,  $Q_{\mathbf{VTS}}(T)$  is the total molecular partition function of the reaction complex divided for the  $Q_{vib}^{irc}(T)$ , the molecular vibrational partition function of the frequency associated to the internal reaction coordinate (*irc*) over which the MEP is computed.  $Q_{SO_2^+}(T)$  and  $Q_{H_2}(T)$  are the total molecular partition functions of the reactants - each can be factorized in the usual electronic, vibrational, rotational and translational factors:  $Q(T) = Q_{el}(T) Q_{vib}(T) Q_{rot}(T) Q_{tr}(T)$ .

A value of  $2.9 \cdot 10^{-11}$  cm<sup>3</sup>molecule<sup>-1</sup>s<sup>-1</sup> was obtained at 300 K by eq 3 for reaction  $SO_2^+ + H_2$ , close to that obtained experimentally by Anicich<sup>[47]</sup>. The calculated  $k$  for the reaction with  $D_2$  is  $1.20 \cdot 10^{-11}$  cm<sup>3</sup> molecule<sup>-1</sup>s<sup>-1</sup>, smaller than that with  $H_2$ , and with an isotope effect of 2.42 at 300 K. The rate coef-




Figure 6: Rate coefficients for the reactions of  $SO_2^+$  with  $H_2$  and  $D_2$  calculated both under thermal equilibrium conditions and following the non-thermal model. See further details in the main text.

efficient  $k$  for the reaction with  $H_2$  calculated from eq 3, increases monotonically with the temperature as shown by the blue dashed line in Figure 6. The same trend is observed for the reaction with  $D_2$  (green dashed line of Figure 6). The thermal rate for  $H_2$  changes its slope at about 800 K, while this is not the case for  $D_2$ . This different trend with temperature leads to the crossing of the two rates at about 1200 K, where the rate coefficient of  $D_2$  becomes higher than the rate of  $H_2$ . This behavior is due to the different increase of the  $Q(T)$  of  $H_2$  relative to that of  $D_2$  caused by a different population with temperature of the respective calculated vibrational levels:  $\nu_{stretching}^{H_2} = 4381 \text{ cm}^{-1}$ ,  $\nu_{stretching}^{D_2} = 3100 \text{ cm}^{-1}$ .

It is proper to point out here that the experimental data are acquired as a function of the photon energy which simultaneously ionize the neutral and excite the ro-vibration levels of the ion, meanwhile the calculated rate coefficient (eq 3) is a function of the temperature. In order to have the rate coefficient as a function of the photon energy, we proceed as follow. A complete conversion of electronic excitation of the ion into ro-vibrational excitation by internal relaxation is assumed<sup>[33]</sup>, and an internal temperature for  $SO_2^+$  is defined with a dependence from the internal

ro-vibrational excitation energy  $E_{SO_2^+}^{int}$ :

$$E_{SO_2^+}^{int} = 3/2k_B T_{SO_2^+} + E_{exc}^{vib}(T_{SO_2^+}) \quad (4)$$

where the first term is the thermal rotational energy for a non-linear molecule, and the second term is the vibrational excitation energy<sup>[54]</sup>:

$$E_{exc}^{vib}(T_{SO_2^+}) = \sum_{i=1}^{3N-6} \frac{h\nu_i}{e^{\frac{h\nu_i}{k_B T_{SO_2^+}} - 1}} \quad (5)$$

where  $\nu_i$  is the  $i^{th}$  vibrational frequency of the ion. Hence, in the top axis of abscissas of Figure 6 we have reported the photon energy ( $h\nu = IE(SO_2) + E_{SO_2^+}^{int}$ ), while on the bottom axis of abscissas it is reported the temperature of the excited ion. It is evident that the trend of the thermal rates (blue and green lines of Figure 6) does not match the experimental observation, where a decrease of the ratio ( $H(D)SO_2^+/SO_2^+$ ) is recorded with the increase of internal energy of  $SO_2^+$  when the photon energy is greater then 12.4 eV (Figure 3). The mismatch between the experimental product/reactant ratio and the theoretical thermal rates led us to carefully analyze the compliance of the experimental conditions with the hypotheses underneath the VTS model. In particular we observe that:

1. The  $SO_2^+$  ion has an internal energy content given by the absorption of the VUV photon, ranging from room temperature, when the incident photon is at the ionization threshold, up to several thousands of Kelvin degrees. On the other hand, the  $H_2$  neutral reactant has an energy content given by its thermal state, which corresponds to the room temperature at which the experiment is performed. Due to low pressure conditions, the thermalization by collisions between the “heated up” ion  $SO_2^+$  and the neutral  $H_2$  can be neglected.
2. The reactive complex formed by  $SO_2^+$  and  $H_2$  is not in thermal equilibrium with the other molecules present in the reaction chamber (low pressure conditions), and it is characterized by non-thermal equilibrium within its internal vibrational degrees of freedom.



Since the IVR (Internal Vibrational Redistribution) within the **VTS** complex can be considered slower than the fast barrierless reaction, we can reasonably assume that the **VTS** complex is non-ergodic in the present experimental conditions<sup>[55]</sup>. The energy flux within the **VTS** complex is controlled by the IVR processes occurring between the reaction coordinate stretching  $H-H$  and the internal degrees of freedom of  $HSO_2$ . We hypothesize that during the reaction the energy flux is from the  $HSO_2$  subset of the **VTS** complex towards the reaction coordinate  $H-H$ , with an increase of the energy flux as the energy content of  $SO_2^+$  increases. The reasons beyond this assumptions are due to the fact that the IVR rate is: *i*) proportional to the density of the final states, and *ii*) is inversely proportional to the exponential of the energy difference between the initial and final states of the IVR (see eq. 5-2 of reference<sup>[56]</sup>). Hence, as the  $SO_2^+$  is excited, the **VTS** complex reaches higher energy states which efficiently couple with the high energy states of the reaction coordinate  $H-H$ , populated during the downhill along the MEP.

The VTST approach, which assumes a local-equilibrium, namely a fast and full molecular energy redistribution in the reaction system, is generally valid when the reactants couple their degrees of freedom within the reactive complex. If the reaction follows a statistic energy redistribution, faster than reaction time, a local thermal equilibrium is always reached<sup>[57,58]</sup>. The hydrogen molecule, which has a very high vibrational stretching frequency due to its light mass, is an optimal neutral reagent to check how much IVR is efficient; i.e. if its high energy vibration can efficiently couple with the other vibrations within the complex. It is known that the VTST is generally valid, but it could fail when a high frequency vibration does not couple efficiently to other low frequencies in a reaction complex as, for instance, in the case of the reaction of H with OH leading to a non-equilibrium effects<sup>[59-61]</sup>.

Following the hypothesis discussed above, we have reformulated the model to calculate the rate coefficient. Eq 3 has been therefore adapted to the conditions of the experiments. We assume that the molecular partition function of  $H_2$  is fixed at environmental temperature of 300 K, the partition function of the

	$a$	$b$	$c$	$d$
$H_2$	238.390	10.160	0.361	10391.0
$D_2$	190.611	21.726	-0.307	10103.7

Table 3: Parameters for  $T_{eff}$  at environmental temperature fixed at  $T=300K$ :  $a$  is in K,  $b$  is in  $K^{1-c}$ ,  $c$  and  $d$  are dimensionless.

**VTS** is factorized in two terms: an effective temperature dependent ro-vibrational part  $Q_{VTS}^{ro-vib}(T_{eff})$  and a translational factor depending on the environmental temperature  $Q_{VTS}^{tr}(300K)$ . The effective temperature ( $T_{eff}$ ) is a function of the internal temperature of  $SO_2^+$  ( $T_{SO_2^+}$ ), and takes into account the part of the ro-vibrational energy which is transferred to the  $H-H$  coordinate during the reaction. The translational partition function of  $SO_2^+$  is fixed at the environmental temperature of 300K. Eq 3 is modified according to the above considerations, and the non-thermal rate coefficient  $k^{NT}$  becomes:

$$k^{NT}(T_{SO_2^+}, 300K) = \quad (6)$$

$$= \sigma \frac{k_B 300K}{h} \left[ \frac{Q_{VTS}^{ro-vib}(T_{eff}) Q_{VTS}^{tr}(300K)}{Q_{SO_2^+}^{ro-vib}(T_{SO_2^+}) Q_{SO_2^+}^{tr}(300K) Q_{H_2}(300K)} \right]$$

We have parameterized the effective temperature in order to reproduce the experimental trend of the data in Figure 3 using the environmental temperature fixed at  $T=300K$ :

$$T_{eff}(T_{SO_2^+}) = a + b \cdot T_{SO_2^+}^c - \frac{d}{T_{SO_2^+}} \quad (7)$$

The parameters used are reported in Table 3, while in Figure S2 of SI the  $T_{eff}(T_{SO_2^+})$  for both  $H_2$  and  $D_2$  are reported.

The non-thermal conditions produce an energy flow in the **VTS** complex which can be analyzed in term of the energy in the reaction coordinate  $H-H$  and in all the other ro-vibrational degrees of freedom, whose vibrational frequencies are reported in Figure S3 and S4 of SI. In Figure 7 the internal energy of the **VTS** complex  $E_{VTS}^{int}$  (See eq S1 of SI), and of the reaction coordinate  $E_{RC}^{int}$  (See eq S3 of SI) for both the reaction with  $H_2$  and  $D_2$  are reported as






Figure 7: Energy redistribution for the reactions of  $SO_2^+$  with  $H_2$  and  $D_2$  calculated as a function of the temperature of the ionic reagent. See further details in the main text.

a function of the temperature  $T_{SO_2^+}$ , and its associated energy  $E_{SO_2^+}^{int}$ , which is also reported in the same figure. The energy is clearly preferentially stored in the reaction coordinate, especially when the  $T_{SO_2^+}$  increases. This can be rationalized in term of the IVR rate, which is more efficient when transferring energy from the **VTS** complex to the reaction coordinate then vice-versa. This is also in agreement with the experimental product/reagent ratio, which decreases as the energy of the  $SO_2^+$  increases. This counter-intuitive trend can be explained considering that the excitation photon energy acquired by the **VTS** complex is not used to increase the number of trajectories in the phase space of the reaction which lead to the products, but this excitation energy is almost entirely directed into the reaction coordinate, which at the end of the reaction will determine a high kinetic energy content of the  $H/D$  atom product.

By plotting the rate coefficient of eq 6 as a function of the temperature (see black line of Figure 6), it is evident that the behavior of the rate coefficient for reaction with  $H_2$  is completely different from that obtained with thermal equilibrium conditions using eq 3, (blue line in Figure 6). The thermal approach



produces an increase of the rate coefficient with temperature, whereas the non-thermal rate exhibits a decrease with temperature in agreement with the experimental data (black line in Figure 3). The same considerations are valid for the reaction with  $D_2$ . Moreover, at 300 K, when all the vibrational modes are at the same temperature, a value of  $1.66 \cdot 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  is calculated for  $SO_2^+ + H_2$  reaction, in agreement with the experimental rate constant  $1.7 \cdot 10^{-11} \pm 40\% \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  [47]. It is noteworthy to mention that the same discrepancy between experiments and theory was also observed in the reaction of  $CH_4^+$  with  $H_2$  at temperatures higher than 100 K [26,27,62].

The fitting parameters for thermal rates for reactions of  $SO_2^+$  with  $H_2O$ ,  $CH_4$ ,  $H_2$  and  $D_2$  are reported in table S1 of SI in the temperature range 200-5000K. These thermal rates have been calculated by means of eq. 3, whose values between relevant stratospheric temperatures (200-250K) are reported in table S2 of SI. The fitting parameters for non-thermal rates of the reaction of  $SO_2^+$  with  $H_2$  and  $D_2$  in the environment temperature  $T$  fixed at 300K are calculated for  $T_{SO_2^+}$  in the range 300-5000K (eq. 6) and given in table S3 of SI. While the fitting parameters for non-thermal rates of the reaction of “hot”  $SO_2^+$  ( $T_{SO_2^+} = 5000\text{K}$ ) with  $H_2$  and  $D_2$  are reported in table S4 of SI for the environment temperature  $T$  in the range 200-250 K, and whose values are also shown in table S5. These last rate coefficients are calculated following eq. S5-S7 of SI. All kinetic data of these relevant chemical reactions have been reported and should be tested in the solar geoengineering models.

### $SO_2^+$ reactivity in the $O_3$ depletion and $OH$ formation

The potential relevance in the stratospheric environment of the reactions involving  $SO_2^+$  with neutral species such as  $H_2O$ ,  $CH_4$  and  $H_2$  is strictly connected with their fast rate coefficients leading to  $HSO_2^+$  and relevant radical species such as  $OH$ ,  $CH_3$  and  $H$ . The hydrogen atom quickly reacts with  $O_3$  producing  $O_2$  and  $OH$  (see eq 2), with the effect of destroying  $O_3$  and increasing the amount of hy-



droxyl radical. This is the most important oxidant in the stratosphere<sup>[34]</sup>, which oxides the  $SO_2$  to  $HSO_3$ , triggering the process which leads to  $H_2SO_4$  aerosol formation<sup>[14,63]</sup>. The main source of  $OH$  during day-time is the reactions of water (eq 9) and molecular hydrogen (eq 10) with  $O(^1D)$ <sup>[14]</sup>, produced by sunlight photodissociation of ozone (eq 8), while  $OH$  density is strongly reduced during nighttime<sup>[64]</sup>:



On the basis of the chemical reactions eqs 8-10 the oxidation of  $SO_2$  should only occur in the day-time, whereas clear evidences<sup>[65]</sup> exist that a significant amount of  $SO_2$  is oxidized during the night. Indeed, alternative oxidation pathways leading to sulfuric acid have been investigated, and ionic mechanisms seem to be operative during the night due to ionizing radiation<sup>[18]</sup>. Hence, the  $OH$  produced by the reactions of  $SO_2^+$  with water<sup>[33]</sup> ( $SO_2^+ + H_2O \rightarrow HSO_2^+ + OH$ ) and  $H_2$  (through eq 2) should be considered as potential relevant ionic processes of  $H_2SO_4$  aerosol formation during nighttime, and should be tested in the atmospheric models.

Common rate coefficients of ion-molecule reactions involving  $SO_2^+$  ( $10^{-11} - 10^{-9} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) are several orders of magnitude higher than the rate coefficient of the oxidation of  $SO_2$  by  $OH$ , which in standard stratospheric pressure and temperature conditions can be reduced to a pseudo second order reaction, with a rate coefficient<sup>[63]</sup> of about  $5 \cdot 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ . Obviously, the relative efficiency of these neutral-neutral and ion-neutral reactions does not only depend on the relative rate coefficients, but also on the relative concentration of the reagents. These concentrations change with the altitude for  $H_2O$ ,  $CH_4$  and  $H_2$ , as well as with the day-time cycle for the  $OH$ <sup>[64]</sup>; whereas the relative abundance of  $SO_2^+/SO_2$  ( $\sim 10^{-12}$ ) is almost constant<sup>[19]</sup>. Solar geoengineering schemes require that  $SO_2$  is at a

pressure of 1 bar when injected by balloons or aircrafts<sup>[66]</sup>, this implies that the  $SO_2^+$  density could locally reach a maximum value of about  $10^7 \text{ ions/cm}^3$ , which is a relevant concentration for the chemistry in the stratosphere. On the basis of the discussion above, the ion-neutral reactions involving  $SO_2^+$  and leading to  $OH$  may be considered as a relevant source of  $OH$  which can oxide  $SO_2$  during the night, when  $OH$  can not be produced via neutral-neutral reactions triggered by sunlight. Alternative routes for the oxidation of  $SO_2^+$  and  $HSO_2^+$  by  $O_3$  are:



Both reactions in eqs. 11 and 12 are exothermic by 102.9 and 93.4 kJ/mol, respectively<sup>[52,46]</sup>. The experimental rate coefficients for these reactions have not been so far determined, probably due to their low reaction efficiency. This indicates that these oxidation reactions could be negligible with respect to oxidation of  $SO_2$  by  $OH$  produced by reaction of  $SO_2^+$  with water and molecular hydrogen during nighttime.

It is also noteworthy to say that  $HSO_2^+$  fast reacts<sup>[30,33]</sup> with water leading to  $H_3O^+$  and  $SO_2$ . The latter, once re-ionized, can act as a catalyst in the formation of hydronium ion, a dominant ionic species in the stratosphere<sup>[67]</sup>.

## Conclusions

In this work the reaction of the radical cation  $SO_2^+$  with  $H_2/D_2$  has been explored at low pressures, mainly focusing on non thermal effects in the redistribution of the internal energy within the reactive complex. The reaction is a barrierless exothermic process, guided by a molecular complex whose substantial charge and spin changes along the reaction coordinate are the factors that slow down the reactivity of the system. The experimental results show that the ratio  $H(D)SO_2^+/SO_2^+$  decreases when the  $SO_2^+$  ion is excited in the ro-vibrational levels of its ground electronic state. DFT and VTST theories have been used to study the dynamic of the reaction along the MEP,

where a **VTS** reaction complex  $[OSOH \cdots H]^+$  has been identified as the “bottleneck” which controls the reactivity of the system. VTST in canonical approximation allows the calculation of the rate coefficient at different excitation energies of  $SO_2^+$ . The value at 300 K is  $2.9 \cdot 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  in reasonable agreement with the value of  $1.7 \cdot 10^{-11} \pm 40\% \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  measured with a FT-ICR instrument. At higher excitation energy of  $SO_2^+$  the standard VTST rate coefficient  $k$  increases within the frame of the thermal equilibrium model, while a decrease of the ratio  $H(D)SO_2^+/SO_2^+$  is observed in the experiments. This opposite trend between theory and experiment has been attributed to a fast reaction dynamics with a light atom, where the thermal equilibrium approximation holds no more. This favors the IVR processes which transfer ro-vibrational energy from the **VTS** reaction complex towards the reaction coordinate  $H-H$ .

The non-thermal assumptions have been taken into account in the calculation of partition functions of  $H_2$  and **VTS**. The  $H_2$  temperature has been kept at 300 K, the initial temperature in the experimental apparatus, while the **VTS** has been evaluated at an effective temperature parameterized over the experimental product/reagent ratio. The rate coefficients obtained with this non-thermal model decrease with the increase of the excitation energy of  $SO_2^+$ , consistently with the experimental data.

The ionization of  $SO_2$  and its reactions with neutral molecules such as  $H_2$  and  $H_2O$  in the stratosphere should be considered as relevant channels in the  $OH$  production during nighttime, when the common neutral-neutral reactions producing  $OH$  via sunlight are not at work. Hence, in the night the above ion-neutral reactions may trigger the oxidation of  $SO_2$ , which subsequently leads to  $H_2SO_4$  aerosol formation. This may explain the decrease of  $SO_2$  observed during the nighttime by several studies. Moreover, the reaction of  $SO_2^+$  with  $H_2$  is relevant for the ozone depletion process. The results of this work indicate that the ion chemistry should be considered and tested in the atmospheric models for solar geoengineering strategies because it can alter the chemistry in the stratosphere.

## Experimental Section

The description of the CiPo beamline at ELETTRA has been reported previously<sup>[68–70,32,71,72]</sup>. Briefly, the beamline is equipped with an electromagnetic elliptical undulator/wiggler and a Normal Incidence Monochromator (NIM) to cover the vacuum ultraviolet (VUV) 8–40 eV energy range. The aluminum grating of the NIM operates in the energy range 8–17 eV with a resolving power of about 1000. The photon energy was calibrated against the autoionization features observed in the Ar total photoionization between the 3p spin orbit components<sup>[73]</sup>. The  $SO_2$  is introduced in the ionization source at the pressure of about  $5.0 \cdot 10^{-6}$  mbar and ionized with the VUV radiation. The ions produced at a photon energy higher than the ionization threshold (about 12.4 eV) are internally excited. The  $SO_2^+$  ions are guided into the octupole (reactive zone) at a CE=0.0 eV, which is determined by measuring the  $SO_2^+$  yield as a function of the retarding field. The estimated energy spread is about 100 meV. The neutral reagent  $H_2/D_2$  gas is introduced in the reactive zone by a needle valve at pressure of about  $10^{-5}$  mbar and room temperature. The ion-molecule reaction of  $SO_2^+$  with  $H_2/D_2$  is followed by recording the yields of ionic reagent ( $m/z$  64) and ionic product ( $m/z$  65 or 66), corrected for the natural isotopic abundance<sup>[52]</sup> contribution of  $^{33}S$ ,  $^{17}O$ ,  $^{34}S$ ,  $^{18}O$ , as a function of the photon energy from 12.4 to 15.0 eV in steps of 100 meV and with the acquisition time of 30s/point. The pressure and CE were kept fixed during the energy scan. The mass spectrum of all the ionic species has been acquired in the mass over charge range  $10 < m/z < 70$  (acquisition time of 3s/point) at the photon energy  $h\nu=14.0$  eV, and CE=0.0 eV. The presence of possible contaminants in the apparatus, as water, has been checked and excluded with and without the neutral ( $H_2/D_2$ ) inside the octupole. Sometime a small amount of  $SO^+$  due to the ionization of  $SO_2$ , without the neutral in the reaction zone, has been observed. However, it is well known the  $SO^+$  does not affect the title reaction since it does not react with  $H_2$  as also verified in our experiments<sup>[47]</sup>. The reaction efficiencies were evaluated by calculating the ratio  $HSO_2^+/SO_2^+$  and using statistical propagation error formula to es-

timate the error bars. Data analysis has been performed using OriginPro8 program.

**Materials.** All the samples were used at room temperature. Sulfur dioxide was purchased from Sigma-Aldrich with a purity > 99.98% whereas hydrogen  $H_2$  and deuterium  $D_2$  are from SIAD with purity > 99.99%.

## Computational Methods

The energetic and dynamical description of the title reaction has been based on the Density Functional Theory formalism, which has been used to explore the Potential Energy landscape of the hydrogen transfer from  $H_2$  to the  $SO_2^+$  ion. The hybrid exchange-correlation functional Becke, three-parameter, Lee-Yang-Parr<sup>[74-77]</sup> has been used with the split-valence double-zeta Pople with polarization and diffuse functions: 6-31++g\*\* basis set<sup>[78,79]</sup>. The region of reaction of the PES has been computed by full optimizations by scanning the  $R_{O-H}$  and  $R_{H-H}$  coordinates, where the hydrogen atom is involved in the exchange with the O atom of the  $SO_2$  ion. The scanning coordinate  $R_{O-H}$  started at a 0.9 Å and ended at 4.0 Å, the coordinate  $R_{H-H}$  was scanned by starting at 0.77 Å up to 5.0 Å, both coordinates were scanned with a variable step whose minimum value as been taken as 0.00375 Å: this quantum calculation has generated a Minimum Energy Path (MEP) which is needed to compute the reaction rate coefficients. The variational transition state theory (VTST) model, specifically developed for applications with barrierless reactions<sup>[59]</sup> has been adopted here to model the present reaction. All these calculations have been corrected by the zero point energies, with the underlying harmonic vibrational frequencies scaled by the coefficient 0.986<sup>[80]</sup>. The charge and spin population are based on the Mulliken analysis of the electron density<sup>[81]</sup>. The MEP has been used to compute the total molecular partition functions  $[Q(T)]$  of the reactive complex  $[OSO \cdots H_2]^+$  in the range of temperatures 200-5000K. Within the Variational Transition State Theory (VTST)<sup>[39]</sup> these partition functions are used to localize the kinetic bottleneck of the reactive flux of trajectories moving along the MEP.

Even if B3LYP is not an high level approach

to compute reaction energy paths, nevertheless it has been demonstrated to correctly reproduce with the VTST the experimental rate coefficients for ionic reactions involving Hydrogen Atom Transfer (HAT)<sup>[33,82]</sup>. In particular B3LYP minimum energy paths are in agreement with those calculated at an higher level of theory (CCSD(T)/aug-cc-pVTZ//MP2/aug-cc-pVTZ) for the HAT reaction studied in reference<sup>[82]</sup>. B3LYP has also been used to compute VTST rate coefficients for barrierless HAT reactions for the modeling of polycyclic aromatic hydrocarbon (PAH) formation in the Interstellar Medium<sup>[53,83,84]</sup>. The DFT has been used as a faster computational method for calculating the rates because the VTST approach we have followed requires a very dense grid (typically with a step  $\leq 0.01$  Å) of the minimum energy path, with frequency analysis at every point of the minimum energy path around which the VTS is located. It would therefore, given the several thousands of points needed to be evaluated, be very computationally demanding the use of an high-level ab initio method. Anyhow, the reason beyond the reliability of B3LYP-VTS rate coefficients for barrierless reactions is based on the fact that for these reactions the VTST does not use energy values but only geometrical data (molecular rotational partition functions) and vibrational frequencies (molecular vibrational partition functions), which are generally well reproduced by B3LYP even if the reactive potential energy path is not of high level. In any case the barrierless nature of this reaction has been confirmed by exploring the MEP with an higher level of ab-initio methodology, namely by computing the MEP at B2PLYPD/aug-cc-pvtz whose geometries has been used to compute an energy profile at CCSD/aug-cc-pvtz level of theory (see fig. S6 of SI). All the quantum chemical calculations were performed with the Gaussian09 package<sup>[85]</sup>.

## Acknowledgments

The work has been supported by the MIUR FIRB RBFR10SQZI project. The authors thank Fabio Zuccaro for his excellent technical assistance.

**Keywords:** ion-molecule reactions, rate coeffi-

cient, reaction mechanisms, synchrotron radiation, VTST.

## References

- [1] *Climate Intervention: Reflecting Sunlight to Cool Earth*, Washington, DC: The National Academies Press, **2015**.
- [2] K. Caldeira, G. Bala, L. Cao, *Annu. Rev. of Earth and Pl. Sci.* **2013**, *41*, 231–256.
- [3] C. Marchetti, *Clim. Change* **1977**, *1*, 59–68.
- [4] D. W. Keith, *Annu. Rev. Energ. Environ.* **2000**, *25*, 245–284.
- [5] A. Robock, L. Oman, G. L. Stenchikov, *J. Geophys. Res.* **2008**, *113*, D16101.
- [6] P. Irvine, K. Emanuel, J. He, L. W. Horowitz, G. Vecchi, D. Keith, *Nat. Clim. Change* **2019**, *9*, 295–299.
- [7] P. Heckendorn, D. Weisenstein, S. Fueglistaler, B. P. Luo, E. Rozanov, M. Schraner, L. W. Thomason, T. Peter, *Environ. Res. Lett.* **2009**, *4*, 045108.
- [8] P. J. Crutzen, *Clim. Change* **2006**, *77*, 211–219.
- [9] U. Niemeier, S. Tilmes, *Science* **2017**, *357*, 246.
- [10] J. Tollefson, *Nature* **2019**, *567*, 156.
- [11] M. G. Lawrence, S. Schafer, H. Muri, V. Scott, N. E. Oschlies, A. Vaughan, H. Boucher, O. Schmidt, J. Haywood, J. Scheffran, *Nat. Comm.* **2018**, *9*, 3734.
- [12] . C. Jones, J. M. Haywood, N. Dunstone, K. Emanuel, M. K. Hawcroft, K. I. Hodges, A. Jones, *Nat. Comm.* **2017**, *8*, 1382.
- [13] C. H. Trisos, G. Amatulli, J. Gurevitch, A. Robock, L. Xia, B. Zambri, *Nat. Ecol. & Evol.* **2018**, *2*, 475–482.
- [14] J. B. Burkholder, S. P. Sander, J. R. Abbatt, J. amd Barker, R. E. Huie, C. E. Kolb, M. J. Kurylo, V. L. Orkin, D. M. Wilmouth, , P. H. Wine, *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 18*, Jet propulsion laboratory, pasadena technical report, **2015**.
- [15] K. L. Aplin, A. A. Briggs, R. G. Harrison, G. J. Marlton, *Space Weather* **2017**, *15*, 663–672.
- [16] G. A. Bazilevskaya, I. G. Usoskin, E. O. Fluckiger, R. G. Harrison, L. Â. Desorgher, R. Butikofer, M. B. Krainev, V. S. Makhmutov, Y. I. Stozhkov, A. K. Svirzhevskaya, N. S. Svirzhevsky, G. A. Kovaltsov, *Space Sci. Rev.* **2008**, *137*, 149–173.
- [17] Y. I. Stozhkov, N. S. Svirzhevsky, G. A. Bazilevskaya, A. N. Kvashnin, V. S. Makhmutov, A. K. Svirzhevskaya, *Adv. Space Res.* **2009**, *44*, 124–1137.
- [18] M. B. Enghoff, N. Bork, S. Hattori, C. Meusinger, M. Nakagawa, J. O. P. Pedersen, S. Danielache, Y. Ueno, M. S. Johnson, N. Yoshida, H. Svensmark, *Atmospheric Chemistry and Physics* **2012**, *12*, 5319–5327.
- [19] N. S. Shuman, D. E. Hunton, A. A. Viggiano, *Chemical Reviews* **2015**, *115*, 4542–4570.
- [20] H. Svensmark, M. B. Enghoff, N. J. Shaviv, J. Svensmark, *Nat. Comm.* **2017**, *8*, 2199.
- [21] Z. Dai, D. K. Weisenstein, D. W. Keith, *Geophys. Res. Lett.* **2018**, *45*, 1030.
- [22] V. Kharchenko, A. Dalgarno, *J. Geophys. Res.* **2004**, *109*, D18311.
- [23] K. Takahashi, N. Taniguchi, Y. Sato, Y. Matsumi, *J. Geophys. Res.* **2002**, *107*, 4290.
- [24] J. O. Wise, D. R. Smith, N. B. Wheeler, M. Ahmadian, R. M. Nadile, *J. Spacecr. Rockets* **2001**, *38*, 297.

- [25] M. Goss-Custard, J. J. Remedios, A. Lambert, E. W. Taylor, C. D. Rodgers, M. Lopez-Puertas, G. Zaragoza, M. R. Gunson, M. R. Suttie, J. E. Harries, J. M. Russell, *J. Geophys. Res.* **1996**, *101*, 9907.
- [26] O. Asvany, I. Savic, S. Schlemmer, D. Gerlich, *Chem. Phys.* **2004**, *298*, 97–105.
- [27] B. Wang, H. Hou, *J. Phys. Chem. A* **2005**, *109*, 8537–8547.
- [28] T. Rockmann, T. S. Rhee, A. Engel, *Atmos. Chem. Phys. Discuss.* **2003**, *3*, 3745–3768.
- [29] G. de Petris, A. Cartoni, A. Troiani, V. Barone, P. Cimino, G. Angelini, O. Ursini, *Chem. Eur. J.* **2010**, *16*, 6234–6242.
- [30] G. de Petris, A. Cartoni, A. Troiani, G. Angelini, O. Ursini, *Phys. Chem. Chem. Phys.* **2009**, *11*, 9976–9978.
- [31] G. de Petris, A. Cartoni, M. Rosi, V. Barone, C. Puzzarini, A. Troiani, *ChemPhysChem* **2011**, *12*, 112–115.
- [32] D. Catone, M. Satta, A. Cartoni, M. C. Castro-villi, P. Bolognesi, L. Avaldi, *Front. Chem.* **2019**, *7*, 1–10.
- [33] A. Cartoni, D. Catone, P. Bolognesi, M. Satta, P. Markus, L. Avaldi, *Chem. Eur. J.* **2017**, *23*, 6772–6780.
- [34] R. G. Prinn, J. Huang, R. F. Weiss, D. M. Cunnold, P. J. Fraser, P. G. Simmonds, A. McCulloch, C. Harth, P. Salameh, S. O’Doherty, R. H. J. Wang, L. Porter, B. R. Miller, *Science* **2001**, *292*, 1882–1888.
- [35] D. M. Murphy, D. S. Thomson, M. J. Mahoney, *Science* **1998**, *282*, 1664–1669.
- [36] R. L. M. Iii, T. Berndt, M. Sipila, P. Paasonen, T. Petaja, S. Kim, T. Kurten, F. Stratmann, V.-M. Kerminen, M. Kulmala, *Nature* **2012**, *488*, 193–196.
- [37] B. Long, J. L. Bao, D. G. Truhlar, *J. Am. Chem. Soc.* **2016**, *138*, 14409–14422.
- [38] D. G. Truhlar, B. C. Garrett, *Annu. Rev. Phys. Chem.* **1984**, *35*, 159–189.
- [39] J. L. Bao, D. G. Truhlar, *Chem. Soc. Rev.* **2017**, *46*, 7548–7596.
- [40] D. Fairley, G. Scott, D. Milligan, R. MacLagan, M. McEwan, *Int. J. Mass Spectrom.* **1998**, *172*, 79–87.
- [41] G. B. Scott, D. A. Fairley, C. G. Freeman, M. J. McEwan, P. Spanel, D. Smith, *J. Chem. Phys.* **1997**, *106*, 3982.
- [42] A. Derossi, F. Lama, M. Piacentini, T. Prosperi, N. Zema, *Rev. Sci. Instrum.* **1995**, *66*, 1718–1720.
- [43] W. Z. Li, M. B. Huang, B. Z. Chen, *J. Chem. Phys.* **2004**, *120*, 4677–4682.
- [44] D. M. P. Holland, M. A. MacDonald, M. A. Hayes, P. Baltzer, L. Karlsson, M. Lundqvist, B. Wannberg, W. von Niessen, *Chem. Phys.* **1994**, *188*, 317–337.
- [45] B. Brehm, J. Eland, R. Frey, A. Kustler, *Int. J. Mass Spectrom. Ion Phys.* **1973**, *12*, 197–211.
- [46] S. Lias, J. Bartmess, J. Liebman, J. Holmes, R. Levin, W. Mallard, *J. Phys. Ref. Data* **1988**, *17*, Suppl. No.1.
- [47] V. G. Anicich, *J. Phys. Chem. Ref. Data* **1993**, *22*, 1469–1569.
- [48] T. Su, W. J. Chesnavich, *J. Chem. Phys.* **1982**, *76*, 5183–5185.
- [49] A. M. Kosmas, *J. Physique Lett.* **1984**, *46*, 799–804.
- [50] S. J. Blanksby, G. B. Ellison, *Acc. Chem. Res.* **2003**, *36*, 255–263.
- [51] B. Ruscic, D. Feller, D. A. Dixon, K. A. Peterson, L. B. Harding, R. L. Asher, A. F. Wagner, *J. Phys. Chem. A* **2001**, *105*, 1–4.

- [52] *Computational Chemistry Comparison and Benchmark Database NIST Standard Reference Database Number 101 Release 19*, Editor: Russell d. johnson iii <http://cccbdb.nist.gov/> technical report, **April 2018**.
- [53] M. Satta, F. Gianturco, F. Carelli, R. Wester, *Astrophys. J.* **2015**, 799, 228.
- [54] D. Egorov, L. Schwob, M. Lalande, R. Hoekstra, T. Schlatholter, *Phys. Chem. Chem. Phys.* **2016**, 18, 26213–26223.
- [55] T. I. Solling, T. S. Kuhlman, A. B. Stephansen, L. B. Klein, K. B. Moller, *ChemPhysChem* **2014**, 15, 249–259.
- [56] *Chemical Reactions in Clusters*, (Ed.: E. R. Bernstein), Oxford University Press, Oxford, New York, **1996**, p. 158.
- [57] A. A. Viggiano, R. A. Morris, J. S. Paschkewitz, J. F. Paulson, *J. Am. Chem. Soc.* **1992**, 114, 10477–10482.
- [58] J. Liu, S. L. Anderson, *Int. J. Mass Spectrom.* **2005**, 241, 173–184.
- [59] A. Fernandez-Ramos, J. A. Miller, S. J. Klippenstein, D. G. Truhlar, *Chem. Rev.* **2006**, 106, 4518.
- [60] J. A. Miller, B. C. Garrett, *Int. J. Chem. Kinet.* **1997**, 29, 275–287.
- [61] J. A. Miller, S. J. Klippenstein, *Int. J. Chem. Kinet.* **1999**, 31, 753–756.
- [62] W. Federer, H. Villingen, P. Tosi, D. Bassi, E. Fergusson, W. Lindinger, *Astrophysics - State of the Art and Future Directions*, Eds.; Reidel: Boston, **1985**.
- [63] D. K. Weisentein, G. K. Yue, M. K. W. Ko, N. R. J. M. Sze, C. J. Scott, *J. Geophys. Res.* **1997**, 13, 019–035.
- [64] M. Carlotti, P. A. R. Ade, B. Carli, M. Chipperfield, P. A. Hamilton, F. Mencaraglia, I. Nolt, M. Ridolfi, *Journal of Atmospheric and Solar Terrestrial Physics* **2001**, 63, 1509–1518.
- [65] J. R. Green, M. N. Fiddler, J. S. Holloway, D. L. Fibiger, E. E. McDuffie, P. Campuzano-Jost, J. C. Schroder, J. L. Jimenez, A. J. Weinheimer, J. Aquino, D. D. Montzka, S. R. Hall, K. Ullmann, V. Shah, L. Jaegle, J. A. Thornton, S. Bililign, S. S. Brown, *Journal of Geophysical Research: Atmospheres* **2019**, 124, 6630–6649.
- [66] J. P. Smith, J. A. Dykema, D. W. Keith, *Earth and Space Science* **2018**, 5, 150–162.
- [67] A. A. Viggiano, F. Arnold in *Handbook of ATMOSPHERIC ELECTRODYNAMICS Volume I* (Ed.: H. Volland), Taylor and Francis, **1995**, Chapter 1.
- [68] A. Cartoni, A. R. Casavola, P. Bolognesi, S. Borocci, L. Avaldi, *J. Phys. Chem. A* **2015**, 119, 3704–3709.
- [69] A. Cartoni, P. Bolognesi, E. Fainelli, L. Avaldi, *J. Chem. Phys.* **2014**, 140, 184307.
- [70] A. Cartoni, A. R. Casavola, P. Bolognesi, M. C. Castrovilli, D. Catone, J. Chiarinelli, R. Richter, L. Avaldi, *J. Phys. Chem. A* **2018**, 122, 4031–4041.
- [71] M. C. Castrovilli, P. Bolognesi, A. Cartoni, D. Catone, O. P., A. R. Casavola, S. Turchini, N. Zema, L. Avaldi, *J. Am. Soc. Mass Spectrom.* **2014**, 25, 351–367.
- [72] M. Satta, P. Bolognesi, A. Cartoni, A. R. Casavola, D. Catone, P. Markus, L. Avaldi, *J. Chem. Phys.* **2015**, 143, 244312.
- [73] G. V. Marr, J. B. West, *At. Data Nucl. Data Tables* **1976**, 18, 497–508.
- [74] A. D. Becke, *J. Chem. Phys.* **1993**, 98, 5648–5652.
- [75] C. Lee, W. Yang, R. G. Parr, *Phys. Rev. B* **1988**, 37, 785–789.
- [76] S. H. Vosko, L. Wilk, M. Nusair, *Can. J. Phys.* **1980**, 58, 1200–1211.



- [77] P. J. Stephens, F. J. Devlin, C. F. Chabalowski, M. J. Frisch, *J. Phys. Chem.* **1994**, *98*, 11623–11627.
- [78] P. Hariharan, J. A. Pople, *Theor. Chim. Acta* **1973**, *28*, 213.
- [79] M. M. Francl, W. J. Pietro, W. J. Hehre, J. S. Binkley, M. S. Gordon, D. J. DeFrees, J. A. Pople, *J. Chem. Phys.* **1982**, *77*, 3654.
- [80] M. Alecu, J. Zheng, Y. Zhao, D. G. Truhlar, *J. Chem. Comp. Theory Comput.* **2010**, *6*, 2872.
- [81] R. S. Mulliken, *J. Chem Phys.* **1955**, *23*, 1833–1840.
- [82] F. A. Gianturco, E. Yurtsever, M. Satta, R. Wester, *J. Phys. Chem. A* **2019**, *123*, 9905–9918.
- [83] F. A. Gianturco, E. Yurtsever, M. Satta, Wester, *Astrophysical Journal* **2017**, *850*, year.
- [84] F. Gianturco, M. Satta, M. Mendolicchio, F. Palazzetti, A. Piserchia, V. Barone, R. Wester, *Astrophysical Journal* **2016**, *830*, 2.
- [85] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Men-  
nucci, H. P. Hratchian, J. V. Ortiz, A. F. Iz-  
maylov, J. L. Sonnenberg, D. Williams-Young,  
F. Ding, F. Lipparini, F. Egidi, J. Goings,  
B. Peng, A. Petrone, T. Henderson, D. Ranas-  
inghe, V. G. Zakrzewski, J. Gao, N. Rega,  
G. Zheng, W. Liang, M. Hada, M. Ehara,  
K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida,  
T. Nakajima, Y. Honda, O. Kitao, H. Nakai,  
T. Vreven, K. Throssell, J. A. Montgomery,  
Jr., J. E. Peralta, F. Ogliaro, M. Bearpark,  
J. J. Heyd, E. Brothers, K. N. Kudin, V. N.  
Staroverov, T. Keith, R. Kobayashi, J. Nor-  
mand, K. Raghavachari, A. Rendell, J. C.  
Burant, S. S. Iyengar, J. Tomasi, M. Cossi,  
J. M. Millam, M. Klene, C. Adamo, R. Cammi,  
J. W. Ochterski, R. L. Martin, K. Morokuma,  
O. Farkas, J. B. Foresman, , D. J. Fox, *Gaussian*  
*09, Revision A.02*, Gaussian, inc., wallingford ct  
technical report, **2016**.