

Analysis of Whole Genome Sequences and Homology Modelling of a 3C Like Peptidase and a Non-Structural Protein of the Novel Coronavirus COVID-19 Shows Protein Ligand Interaction with an Aza-Peptide and a Noncovalent Lead Inhibitor with Possible Antiviral Properties

Arun K. Shanker¹ *, Divya Bhanu^{1,2} and Anjani Alluri³

¹ICAR - Central Research Institute for Dryland Agriculture

Santoshnagar, Hyderabad – 500059, India

Corresponding author email: arunshank@gmail.com

² Centre for Plant Molecular Biology, Osmania University, Hyderabad, India

³ Advanced Post Graduate Centre, Acharya N.G.Ranga Agricultural University, Guntur, India

Abstract

The family of viruses belonging to Coronaviridae mainly consist of virulent pathogens that have a zoonotic property, Severe Respiratory Syndrome (SARS-CoV) and Middle East Respiratory Syndrome (MERS-CoV) of this family have emerged before and now the Novel COVID-19 has emerged in China. Characterization of spike glycoproteins, polyproteins and other viral proteins from viruses are important for vaccine development. Homology modelling of these proteins with known templates offers the opportunity to discover ligand binding sites and possible antiviral properties of these protein ligand complexes. Any information emerging from these protein models can be used for vaccine development. In this study we did a complete bioinformatic analysis, sequence alignment, comparison of multiple sequences and homology modelling of the Novel COVID-19 whole genome sequences, the spike protein and the polyproteins for homology with known proteins, we also analysed receptor binding sites in these models for possible vaccine development. Our results showed that the tertiary structure of the polyprotein isolate COVID-19_HKU-SZ-001_2020 had 98.94 percent identity with SARS-Coronavirus NSP12 bound to NSP7 and NSP8 co-factors. Our results indicate that a part of the viral genome (residues 254 to 13480 in Frame 2 with 4409 amino acids) of the Novel COVID-19 virus isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) when modelled with template 2a5i of the PDB database had 96 percent identity with a 3C like peptidase of SARS-CoV which has ability to bind with Aza-Peptide Epoxide (APE) which is known for irreversible inhibition of SARS-CoV main peptidase. The part of the genome when modelled with template 3e9s of the PDB database had 82 percent identity with a papain-like protease/deubiquitinase which when complexed with ligand GRL0617 acts as inhibitor which can block SARS-CoV replication. It is possible that these viral inhibitors can be used for vaccine development for the Novel COVID-19.

Introduction

More than a decade has passed since the emergence human Coronavirus that caused Severe Respiratory Syndrome (SARS-CoV) and it is about 7 years since the emergence of another type of Coronavirus - Middle East Respiratory Syndrome (MERS-CoV) and now the Novel

34 Coronavirus COVID-19 has emerged in China. This repeated onslaught of these viruses goes
35 to show that it can assume pandemic proportions at any time and at any place.

36 The family of viruses belonging to Coronaviridae mainly consist of virulent pathogens that
37 have a zoonotic property and this large family of corona viruses, have been known to be
38 circulating in animals including camels, cats and bats. It has been seen in the past that Severe
39 Acute Respiratory Syndrome associated coronavirus (SARS-CoV) and Middle East
40 Respiratory Syndrome-associated coronavirus (MERS-CoV) belonging to this family of
41 viruses can be transmitted from animals to humans and can cause respiratory diseases. Human
42 to human transmission on this virus has been a concern and due to this search for antiviral
43 compounds and vaccine development for this family of virus becomes the need of the hour.

44 The SARS was first seen in 2002 in Guangdong province of China, and later spread globally
45 and has caused close to about 8096 cases (WHO 2004, de Wit et al., 2016). In 2012, a novel
46 betacoronavirus, designated Middle East respiratory syndrome coronavirus or MERS-CoV
47 associated with severe respiratory disease in humans, emerged in the Arabian Peninsula (de
48 Wit et al., 2013).

49 The World Health Organization (WHO), China Country Office was informed of cases of
50 pneumonia of unknown aetiology in Wuhan City, Hubei Province, on 31 December 2019
51 (WHO 2020). A novel coronavirus currently termed COVID-19 was officially announced as
52 the causative agent by Chinese authorities on 7 January 2020. As on 3 Feb 2020 China's
53 National Health Commission reported that there are 20,438 confirmed cases in China,
54 including 15 in Hong Kong and eight in Macao. The self-governing island of Taiwan reported
55 10 cases. The World Health Organization reported 319 confirmed cases in 23 countries outside
56 China (WHO Situation Report 21 2020). This novel corona virus has been designated as novel
57 coronavirus COVID-19.




Fig.1 Countries, territories or areas with reported confirmed cases of COVID-19 , 3 February 2020 Source WHO (https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200211-sitrep-22-ncov.pdf?sfvrsn=fb6d49b1_2).

Coronaviruses are RNA viruses and have large genomes structures and due to this they can have high error in replication as compared to host genomes. It is also known that various CoVs can do effective recombination of their genomes after infecting host cells (Luo et al 2018). This recombination can be a factor for their evolution to novel types which may have new animals as their intermediate hosts. These factors give the CoVs high adaptive ability and the capability to jump across species and have a relatively large host range.

Characterization of Spike glycoproteins from viruses are important for vaccine development. Any information coming from the protein model can be used for vaccine development. *In Silico* Epitope, polyprotein and spike protein-based peptide vaccine designing for infectious viruses is a way that can hasten the process of vaccine development. Spike (S) protein, polyprotein and other viral proteins of the novel coronavirus COVID-19 as a target for the development of vaccines and therapeutics for the prevention and treatment of infection is an important

approach. In the case of SARS-CoV, these proteins can mediate binding of the virus with its receptor and promotes the fusion between the viral and host cell membranes and virus entry into the host cell, hence peptides, antibodies, organic compounds and short interfering RNAs that interact with the spike protein can have a potential role in vaccine development (Du et al 2009).

Here in this study we did a complete bioinformatic analysis, sequence alignment, comparison of multiple sequences of the Novel COVID-19 whole genome sequences, the Spike protein and the polyproteins for homology with known spike proteins and also analysed receptor binding sites for possible vaccine development.

Materials and Methods

Six complete viral genome sequences, seven polyproteins (RdRp region) and seven glycoproteins available on NCBI portal on 4 Feb 2020 were taken for analysis. The sequence details and GenBank accession numbers are listed in Table 1. Amongst the seven polyproteins, five are of Wuhan pneumonia virus isolate COVID-19 and two sequences are of Wuhan pneumonia virus isolate SI200040-SP. The seven Glycoproteins are of the same isolate, Wuhan pneumonia virus isolate COVID-19.

Table 1 List of available Wuhan seafood market pneumonia virus isolate sequences at NCBI

Genbank Accession Number	Title	Description
MN988713.1	Wuhan seafood market pneumonia virus isolate COVID-19 /USA-IL1/2020	Complete genome
MN938384.1	Wuhan seafood market pneumonia virus isolate COVID-19 _HKU-SZ-002a_2020	Complete genome

MN975262.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-005b_2020	Complete genome
MN985325.1	Wuhan seafood market pneumonia virus isolate COVID- 19 /USA-WA1/2020	Complete genome
NC_045512.2	Wuhan seafood market pneumonia virus isolate Wuhan- Hu-1	Complete genome
MN908947.3	Wuhan seafood market pneumonia virus isolate Wuhan- Hu-1	Complete genome
MN938385.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-001_2020 ORF1ab polyprotein, RdRp region, (orf1ab) gene, partial cds	Polyprotein, RdRp region
MN938386.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-004_2020 ORF1ab polyprotein, RdRp region, (orf1ab) gene, partial cds	Polyprotein, RdRp region
MN975263.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-007a_2020 ORF1ab polyprotein, RdRp region, (orf1ab) gene, partial cds	Polyprotein, RdRp region
MN975264.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-007b_2020 ORF1ab polyprotein, RdRp region, (orf1ab) gene, partial cds	Polyprotein, RdRp region

MN975265.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-007c_2020 ORFlab polyprotein, RdRp region, (orf1ab) gene, partial cds	Polyprotein, RdRp region
MN970003.1	Wuhan seafood market pneumonia virus isolate SI200040-SP orflab polyprotein, RdRP region, (orf1ab) gene, partial cds	Polyprotein, RdRp region
MN970004.1	Wuhan seafood market pneumonia virus isolate SI200121-SP orflab polyprotein, RdRP region, (orf1ab) gene, partial cds	Polyprotein, RdRp region
MN938387.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-001_2020 surface glycoprotein (S) gene, partial cds	Glycoprotein
MN938388.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-002b_2020 surface glycoprotein (S) gene, partial cds	Glycoprotein
MN938389.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-004_2020 surface glycoprotein (S) gene, partial cds	Glycoprotein
MN938390.1	Wuhan seafood market pneumonia virus isolate COVID- 19 _HKU-SZ-005_2020 surface glycoprotein (S) gene, partial cds	Glycoprotein
MN975266.1	Wuhan seafood market pneumonia virus isolate COVID-	Glycoprotein

	19_HKU-SZ-007a_2020 surface glycoprotein (S) gene, partial cds	
MN975267.1	Wuhan seafood market pneumonia virus isolate COVID-19_HKU-SZ-007b_2020 surface glycoprotein (S) gene, partial cds	Glycoprotein
MN975268.1	Wuhan seafood market pneumonia virus isolate COVID-19_HKU-SZ-007c_2020 surface glycoprotein (S) gene, partial cds	Glycoprotein

92

93 The available polyproteins (RdRp region) and glycoprotein sequences were retrieved from
94 Genbank, NCBI (Benson et al., 2000). These sequences were translated to amino acid
95 sequences using sorted six frame translation with Bioedit (Hall et al., 2011). Multiple sequence
96 alignment of the translated protein sequences was performed and phylogenetic tree was
97 constructed using Mega-X (Kumar et al., 2018). The alignment shows that amongst the seven
98 polyproteins, five sequences were identical being from the same isolate and two other
99 sequences of the other isolate are identical. Similar analysis of the seven glycoproteins was
100 done, all the seven glycoprotein sequences were found to be identical. Therefore, further
101 analysis was carried out for three sequences.

102 1. MN938385.1 Wuhan seafood market pneumonia virus isolate COVID-19_HKU-SZ-
103 001_2020 ORF1ab polyprotein, RdRp region, (orf1ab) gene, partial cds: 0 to 284: Frame 3 95
104 aa

105 2. MN970003.1 Wuhan seafood market pneumonia virus isolate SI200040-SP orf1ab
106 polyprotein, RdRP region, (orf1ab) gene, partial cds: 2 to 289: Frame 2 96 aa

107 3. MN938387.1 Wuhan seafood market pneumonia virus isolate COVID-19_HKU-SZ-
108 001_2020 surface glycoprotein (S) gene, partial cds: 1 to 105: Frame 1 35 aa

Expasy proteomics server (Gasteiger et al., 2003) was used to study the protein sequence and structural details. These peptides were studied for their physio-chemical properties using the tool Protparam (Gasteiger et al., 2005). The secondary structure analysis was done using Chou and Fasman algorithm with CFSSP (Kumar, 2013). To generate the 3D structure from the fasta sequence, homology modelling was performed and the templates were identified. The model was built using the template with highest identity. The structural assessment was also performed to validate the model built. Swiss-model (Schwede et al., 2003) was used to build and validate the 3D model.


Complete genome sequence of the Wuhan seafood market pneumonia virus isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) which has 29903 bp ss-RNA linear was translated sorted 6 frame with minimum ORF of 20 with any start codon and the resultant protein sequence was used for homology modelling, homology models were done with large chunks of proteins 21503 to 25381 in Frame 2 with 1293 amino acids, 13450 to 21552 in Frame 1 with 2701 amino acids and 254 to 13480 in Frame 2 with 4409 amino acids.

SWISS-MODEL server was used for homology modelling (Waterhouse et al 2018) where computation was on ProMod3 engine which is based on Open Structure (Biasini et al 2013). Structural information is extracted from the template, sequence alignment is used to define insertions and deletions.

Protein ligand interaction profile with hydrogen bonding, hydrophobic interactions, salt bridges and π -Stacking was done with PLIP server (Salentin et al., 2015)

132

133 Results and Discussion



134

135 The phylogenetic tree of the seven polyproteins is shown in Fig.2. It is seen that two
136 polyproteins were distinctly different from the rest.

137 **Fig.2** Phylogenetic tree of the seven polyproteins of Wuhan seafood market pneumonia virus
138 isolate

139 The tertiary structure analysis of the isolate COVID-19 _HKU-SZ-001_2020 ORF1ab
140 polyprotein is given in Table 2. It is seen that the polyprotein has a 98.94 percent identity with
141 PDB structure 6nur.1.A and a 19.74 percent identity with a ABC-type uncharacterized transport
142 system periplasmic component-like protein.

143 The Phylogenetic tree of the seven glycoproteins of the Wuhan seafood market pneumonia
144 virus isolate is shown in Fig.3, it is seen that the glycoproteins are similar in all the isolates.



Fig.3 Phylogenetic tree of the seven polyproteins of Wuhan seafood market pneumonia virus isolate

Table 2 Tertiary Structure of Wuhan seafood market pneumonia virus isolate COVID-19 _HKU-SZ-001_2020 ORF1ab polyprotein alignment with templates

PDB Template	Gene	Identity
6nur.1.A	NSP12	98.947
1khv.1.A	RNA-directed RNA polymerase	8.97
1khv.2.A	RNA-directed RNA polymerase	8.97
5z6v.1.A	ABC-type uncharacterized transport system periplasmic component-like protein	19.74
6k1y.1.A	ABC-type uncharacterized transport system periplasmic component-like protein	19.74
2ckw.1.A	RNA-directed RNA polymerase	10.53

part of the viral genome where in the RNA viruses its function is to catalyze the synthesis of the RNA strand complementary to a given RNA template.

Table 3 Physico-chemical properties of polyproteins of Novel Coronavirus 2019 n-CoV

Accession Number	MN938385.1	MN938386.1	MN975263.1	MN975264.1	MN975265.1	MN970003.1	MN970004.1
Reading Frame	3	3	3	3	3	2	2
Number of amino acids	95	95	95	95	95	96	96
Molecular weight	10640.22	10640.22	10640.22	10640.22	10640.22	11239.26	11239.26
Theoretical pI	9.87	9.87	9.87	9.87	9.87	8.9	8.9
Formula	C ₄₇₂ H ₇₅₂ N ₁₃₄ O ₁₃₈ S ₄	C ₄₇₂ H ₇₅₂ N ₁₃₄ O ₁₃₈ S ₄	C ₄₇₂ H ₇₅₂ N ₁₃₄ O ₁₃₈ S ₄	C ₄₇₂ H ₇₅₂ N ₁₃₄ O ₁₃₈ S ₄	C ₄₇₂ H ₇₅₂ N ₁₃₄ O ₁₃₈ S ₄	C ₅₁₆ H ₇₈₆ N ₁₃₂ O ₁₃₂ S ₉	C ₅₁₆ H ₇₈₆ N ₁₃₂ O ₁₃₂ S ₉
Total number of atoms	1500	1500	1500	1500	1500	1575	1575
Extinction coefficients	12950	12950	12950	12950	12950	24200	24200
Instability index	20.51	20.51	20.51	20.51	20.51	29.66	29.66
Aliphatic index	80.11	80.11	80.11	80.11	80.11	89.27	89.27
Grand average of hydropathicity (GRAVY)	-0.264	-0.264	-0.264	-0.264	-0.264	0.161	0.161
Estimated half-life	1.9 hours (mammalian reticulocytes, in vitro).	1.9 hours (mammalian reticulocytes, in vitro).	1.9 hours (mammalian reticulocytes, in vitro).	1.9 hours (mammalian reticulocytes, in vitro).	1.9 hours (mammalian reticulocytes, in vitro).	1.3 hours (mammalian reticulocytes, in vitro).	1.3 hours (mammalian reticulocytes, in vitro).
	>20 hours (yeast, in vivo).	>20 hours (yeast, in vivo).	>20 hours (yeast, in vivo).	>20 hours (yeast, in vivo).	>20 hours (yeast, in vivo).	3 min (yeast, in vivo).	3 min (yeast, in vivo).
	>10 hours (<i>Escherichia coli</i> , in vivo).	>10 hours (<i>Escherichia coli</i> , in vivo).					

The isolates SI200040-SP orflab polyprotein and the isolate SI200121-SP orflab polyprotein had 2 reading frames as compared to the rest of the isolates which had 3 reading frames. The presence of multiple reading frames suggests the possibility of overlapping genes as seen in many virus and prokaryotes and mitochondrial genomes. This could affect how the proteins are made. The number of amino acid residues in all the polyproteins were the same except one isolate SI200040-SP which had one amino acid more than the other polyproteins. The extinction coefficients of the two isolates SI200040-SP orflab polyprotein and the isolate SI200121-SP orflab polyprotein was much higher compared to the rest of the polyproteins.

180 The extinction coefficient is important when studying protein-protein and protein-ligand
181 interactions. The instability index of these two isolates was also high when compared to the
182 others indicating the that these two isolates are instable. Regulation of gene expression by
183 polyprotein processing is known in viruses and this is seen in many viruses that are human
184 pathogens (Yost et al 2013).

185 The isolates here like many other viruses may be using replication strategy which could involve
186 the translation of a large polyprotein with subsequent cleavage by viral proteases. The two
187 isolates SI200040-SP orflab polyprotein and the isolate SI200121-SP orflab polyprotein also
188 showed shorter half-lives as compared to the other isolates indicating that they are susceptible
189 to enzymatic degradation.


190

191 The homology model developed from the residues 254 to 13480 in Frame 2 with 4409 amino
192 acids from the Complete genome sequence of the Wuhan seafood market pneumonia virus
193 isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) which has 29903 bp with
194 linear ss-RNA linear showed interesting template alignments, in all the model aligned with
195 50 templates from the PDB database with most of them being replicase polyprotein lab which
196 is a SARS-CoV papain-like protease (Daczkowski 2017). The maximum similarity of 97.3
197 percent was with template structure of a Nsp9 protein from SARS-coronavirus indicating that
198 this novel coronavirus has high degree of similarity with the SARS-coronavirus and this can
199 be used for gaining insights into vaccine development.

200 The homology models of the 4409 amino acid residues of the whole genome of the Wuhan
201 seafood market pneumonia virus isolate Wuhan-Hu-1 with the ligand association with
202 templates 2a5i and 3e9s are shown in Fig. 5 and Fig. 6 respectively. The regions of their
203 alignment are shown in Fig.7 and Fig.8 respectively. The model has similarity with the 3Clike
204 proteinase and a papain-like protease/deubiquitinase protein which are known antiviral drug

205 targets. Ligand binding and their action is on viral replication and inactivation can be useful in
206 stopping the viral replication (Baez-Santos et al 2015).


207



208

209 **Fig. 5** Homology model with ligand binding of protein from 4409 amino acids 254 to 13480
210 in Frame 2 of the Complete genome sequence of the Wuhan seafood market pneumonia virus
211 isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) which has 29903 bp linear
212 ss-RNA with 2a5i of the PDB database as template.

213



214

215

216 **Fig. 6** Homology model with ligand binding of protein from 4409 amino acids 254 to 13480
 217 in Frame 2 of the Complete genome sequence of the Wuhan seafood market pneumonia virus
 218 isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) which has 29903 bp linear
 219 ss-RNA with 3e9s of the PDB database as template.

220

221

Seqres SGFRKMAFFSGKVEGCMVQVTCGTTTTLNGLWDDTVYCPRHVICTAEDMLNPNVEDLLIRKSNHSFLVQAGNVQLRVIGH 80
 2a5i.1. (AB) SGFRKMAFFSGKVEGCMVQVTCGTTTTLNGLWDDTVYCPRHVICTAEDMLNPNVEDLLIRKSNHSFLVQAGNVQLRVIGH 80

Seqres SMQNCCLRLKVDTSNPKTFKYKPVRIQPGQTFPSVLACYNGSPSGVYQCAMRPNHTIKGSFLNGSCGSGVGFNIIDYDCVSFC 160
 2a5i.1. (AB) SMQNCCLRLKVDTSNPKTFKYKPVRIQPGQTFPSVLACYNGSPSGVYQCAMRPNHTIKGSFLNGSCGSGVGFNIIDYDCVSFC 160

Seqres YHHHMELEPTGVHAGTDLEGKFYGFVDRQTAAAGTDTTITLNVLANLYAAVINGDRWFLNRFTTTLNDNFNLVAMKYNVE 240
 2a5i.1. (AB) YHHHMELEPTGVHAGTDLEGKFYGFVDRQTAAAGTDTTITLNVLANLYAAVINGDRWFLNRFTTTLNDNFNLVAMKYNVE 240

Seqres PLTQDHYVDILGPLSAQTGIAVLDMCAALKELLQNGMNGRTILGSTILEDEFTPFDDVVRQCSCGVTFQ 306
 2a5i.1. (AB) PLTQDHYVDILGPLSAQTGIAVLDMCAALKELLQNGMNGRTILGSTILEDEFTPFDDVVRQCSCGVTFQ 306

Fig.7 Alignment of the region of the Novel Coronavirus COVID-19 protein with template 2a4i of PDB database

Seqres ASMEVKTIKVFTTYDNTNLHTQLVDMSTYGGQFGPTYLDGADVTKIKPHYNHEGKTFEVLPSDDTLRSEAFEYVHTLDESFLGR 85
 3e9s.1.A ASMEVKTIKVFTTYDNTNLHTQLVDMSTYGGQFGPTYLDGADVTKIKPHYNHEGKTFEVLPSDDTLRSEAFEYVHTLDESFLGR 85

Seqres YMSALNHTKKWFFQVGGLTSLKWDNNCYLSSVLLALQOLEVKNAPALQEAAYRARGDAANFCALILAYSNNKTVGELGDVRE 170
 3e9s.1.A YMSALNHTKKWFFQVGGLTSLKWDNNCYLSSVLLALQOLEVKNAPALQEAAYRARGDAANFCALILAYSNNKTVGELGDVRE 170


Seqres TMTHLLQHANLESARKVLNVVCKHCGQKTTTLTGVEAVMYMGTLSYDNLKTVSIPCVCGRDATQYLVQQESSFVMMSPAPAEYK 255
 3e9s.1.A TMTHLLQHANLESARKVLNVVCKHCGQKTTTLTGVEAVMYMGTLSYDNLKTVSIPCVCGRDATQYLVQQESSFVMMSPAPAEYK 255

Seqres LQQGTFLCANEYTGNYQCCHYTHITAKETLYRIDGAHLTKMSEYKGEVTDVFXKETSXTTIR 318
 3e9s.1.A LQQGTFLCANEYTGNYQCCHYTHITAKETLYRIDGAHLTKMSEYKGEVTDVFXKETSXTTIR 317

Fig.8 Alignment of the region of the Novel Coronavirus COVID-19 protein with template 3e9s of PDB database

The important templates that aligned with this 4409 amino acid residues of the whole genome of the Wuhan seafood market pneumonia virus isolate Wuhan-Hu-1 were 2a5i of the PDB database which is a crystal structure of SARS coronavirus main peptidase inhibited by an Aza-Peptide epoxide in the space group C2 (Lee et al 2005) and 3e9s of the PDB database which is new class of papain-like protease/deubiquitinase which when combined with ligand GRL0617 acts as inhibitors blocking SARS virus replication (Ratia et al 2008). The model with template 2a5i of the PDB database shows that Aza-Peptide Epoxide (APE; kinact/Ki=1900(±400) M⁻¹ s⁻¹) which is a known anti SARS agent can be used to develop a molecular target with irreversible inhibitor properties.

The protein ligand interaction analysis of the Novel Coronavirus C3 like peptidase and aza-peptide epoxide is shown in Fig.9



239

240 **Fig.9** Protein Ligand interaction between the C3 like peptidase with aza-peptide epoxide

241 The substrate binding properties and structural and chemical complementarity of this Aza-
 242 Peptide Epoxide can be explored as an anti - Coronavirus COVID-19 agent. The APE which
 243 is ethyl (2S)-4-[(3-amino-3-oxo-propyl)-[(2S)-2-[(2S)-4-methyl-2-
 244 phenylmethoxycarbonylamino-pentanoyl]amino]-3-phenyl-propanoyl]amino]amino]-2-
 245 hydroxy-4-oxo-butanoate structure is shown in Fig.10.




Fig. 10 Structure of Aza-Peptide Epoxide (APE) ethyl (2S)-4-[(3-amino-3-oxo-propyl)-[[[(2S)-2-[[[(2S)-4-methyl-2-phenylmethoxycarbonylamino-pentanoyl]amino]-3-phenylpropanoyl]amino]amino]-2-hydroxy-4-oxo-butanoate with possible anti Coronavirus activity – (Source <https://www.rcsb.org/ligand/AZP>)

The model with template 3e9s of the PDB database shows that the Coronavirus viral protein can have a ligand which is a papain-like protease (PLpro) that is known to be a potent inhibitor of viral replication in SARS (Ratia et al 2008).

The two parts of the Main protein from the whole genome of the Novel Coronavirus COVID-19 aligned with two SAR proteins and the ligand binding sites were similar, the alignment positions, number of amino acids and ligand and the interacting residues is given in Table 3

The main protein with a sequence length of 5509aa of the Wuhan Corona Virus showing structural alignment with two other proteins of SARS-CoV is given in Table 4

Table 4 Main Protein with a sequence length – 4409aa of Wuhan Corona Virus showing structural alignment with two other proteins of SARS-CoV

Template ID	Template Title	Alignment Positions	Number of aa	Ligands	Interacting Residues
3e9s.1	A new class of papain-like protease/deubiquitinase inhibitors blocks SARS virus replication	1568-1882	315	TTT	Chain A: L.1729, G.1730, D.1731, P.1814, P.1815, Y.1831, Y.1835, Q.1836, Y.1840, T.1868
2a5i.1	Crystal structures of SARS coronavirus main peptidase inhibited by an aza-peptide epoxide in the space group C2	3268-3573	306	AZP	Chain A: T.3292, T.3293, H.3308, M.3316, Y.3321, F.3407, L.3408, N.3409, G.3410, S.3411, C.3412, H.3430, H.3431, M.3432, E.3433, P.3435, H.3439, D.3454, R.3455, Q.3456, T.3457, A.3458, Q.3459

The complete genome of MN908947.3 Wuhan seafood market pneumonia virus isolate Wuhan-Hu-1 encodes a 4409aa long protein along with the other glycoproteins and polyproteins. The homology modelling of this protein showed sequence and structural alignment with two SARS proteases with structural accession numbers 3e9s.1 and 2a5i.1 at positions 1568-1882 and 3268-3573 respectively. Reports suggests inhibition of virus replication by TTT ligand and an aza-peptide epoxide inhibiting the main peptidase. The structural similarity of these templates are 83% and 96% respectively. The multiple sequence alignment shows complete conservation of the sequence suggesting a high degree of homology.

The protein ligand interaction analysis of the Novel Coronavirus non structural protein and papain-like protease is shown in Fig. 11




Fig.11 Protein Ligand interaction between the Novel Coronavirus non structural protein and papain-like protease

The Hydrophobic interaction, hydrogen bonding, salt bridges of the constructed model of the Novel Coronavirus protein from region 3268-3573 aa to ligand AZP is given in Suppl. Table 1, the Hydrophobic interaction, hydrogen bonding, salt bridges of the template 2a5i is given in Suppl. Table 2, when comparing both it is seen that the binding properties are the same except for the presence of water bridge in the template 2a5i.

The Hydrophobic interaction, hydrogen bonding, π -Stacking of the constructed model of the Novel Coronavirus protein from region 1568-1882 aa to ligand Small molecule Noncovalent Lead Inhibitor is given in Suppl. Table 3, the Hydrophobic interaction, hydrogen bonding, π -Stacking of the template 3e9s is given in Suppl. Table 4, when comparing both it is seen that the binding properties are the same except or an addition π -Stacking at Tyr in the template 2a5i. This shows that there is high possibility of binding of the these antiviral compounds with the regions of Novel Coronavirus protein that is in homology with the SARS protein.

289

290 from the in case of template 2a5i and π -Stacking in the case of template 3e9s

291 The targeting of this part of the genome of the Novel Coronavirus COVID-19 with the antiviral
292 compounds which have to shown to bind in the similar region of the SARS virus can have
293 implication in the development of an effective antiviral compound against the Novel
294 Coronavirus COVID-19 . The residues 254 to 13480 in Frame 2 with 4409 amino acids from
295 the Complete genome sequence of the Wuhan seafood market pneumonia virus isolate Wuhan-
296 Hu-1 shows homology with the SARS coronaviral proteases, papain-like protease (PLpro) and
297 3C-like protease (3CLpro), these proteins have the function of processing the viral polyprotein
298 and also they perform the function of stripping ubiquitin and the ubiquitin-like interferon
299 (IFN)-stimulated gene 15 (ISG15) from the hosts to facilitate coronavirus replication and help
300 in evading immune response of the host, these inhibitors can also have a role in disrupting
301 signalling cascades in infected cells and protecting the uninfected cells.

302 The chemical GRL0617 is 5-Amino-2-methyl-N-[(1R)-1-(1-naphthalenyl)ethyl]benzamide
303 and is known to inhibit the papainlike protease that is present in SARS CoV . This protease is
304 a potential target for antiviral compounds (Chaudhuri et al., 2011). We found the Novel
305 COVID-19 has homology with this and the binding sites for this in the structural protein of the
306 Novel COVID-19 is the same (Table 4). This compound inhibits the enzyme that is required
307 for the cleavage of the viral protein from the virus in SARS CoV, it also cleaves ubiquitin and
308 has a structural homology with the Deubiquitinases (DUBs) of the Ubiquitin-Specific Proteases
309 Compound GRL0617 binds in the S4 and S3 enzyme subsite that gets the C terminal tail of the
310 Ubiquitin (King and Finley 2014; Schauer et al., 2019). Our results show that Aza-Peptide
311 Epoxide an irreversible protease inhibitor and GRL0617 a viral replication inhibitor can be
312 used to develop inhibitors of the Novel Coronavirus COVID-19.

313

314 **References**

315 Bae, J.E., Kim, I.J., Kim, K.J. and Nam, K.H., 2018. Crystal structure of a substrate-binding
316 protein from *Rhodothermus marinus* reveals a single α/β -domain. Biochemical and
317 Biophysical Research Communications, 497(1), pp.368-373.

318 Baez-Santos, Y.M., John, S.E.S. and Mesecar, A.D., 2015. The SARS-coronavirus papain-like
319 protease: structure, function and inhibition by designed antiviral compounds. Antiviral
320 Research, 115, pp.21-38.

321 Benson, D. A., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., Rapp, B. A., & Wheeler, D. L.
322 (2000). GenBank. Nucleic Acids Rresearch, 28(1), 15-18.

323 Biasini, M., Schmidt, T., Bienert, S., Mariani, V., Studer, G., Haas, J., Johner, N., Schenk,
324 A.D., Philippsen, A. and Schwede, T., 2013. OpenStructure: an integrated software
325 framework for computational structural biology. Acta Crystallographica Section D:
326 Biological Crystallography, 69(5), pp.701-709.

327 Chaudhuri, R., Tang, S., Zhao, G., Lu, H., Case, D.A. and Johnson, M.E., 2011. Comparison
328 of SARS and NL63 papain-like protease binding sites and binding site dynamics:
329 inhibitor design implications. Journal of molecular biology, 414(2), pp.272-288.

330 Daczkowski, C.M., Dzimianski, J.V., Clasman, J.R., Goodwin, O., Mesecar, A.D. and Pegan,
331 S.D., 2017. Structural insights into the interaction of coronavirus papain-like proteases
332 and interferon-stimulated gene product 15 from different species. Journal of Molecular
333 Biology, 429(11), pp.1661-1683.

334 de Wit, E., Rasmussen, A.L., Falzarano, D., Bushmaker, T., Feldmann, F., Brining, D.L.,
335 Fischer, E.R., Martellaro, C., Okumura, A., Chang, J. and Scott, D., 2013. Middle East

336 respiratory syndrome coronavirus (MERS-CoV) causes transient lower respiratory tract
337 infection in rhesus macaques. *Proceedings of the National Academy of Sciences*,
338 110(41), pp.16598-16603.

339 de Wit, E., van Doremalen N., D. Falzarano, V. J. Munster, SARS and MERS: recent insights
340 into emerging coronaviruses. *Nat Rev Microbiol* 14, 523-534 (2016).

341 Du, L., He, Y., Zhou, Y., Liu, S., Zheng, B.J. and Jiang, S., 2009. The spike protein of SARS-
342 CoV—a target for vaccine and therapeutic development. *Nature Reviews*
343 Microbiology, 7(3), pp.226-236.

344 Fehr, A.R. and Perlman, S., 2015. Coronaviruses: an overview of their replication and
345 pathogenesis. In *Coronaviruses* (pp. 1-23). Humana Press, New York, NY.

346 Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R. D., & Bairoch, A. (2003).
347 ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucleic*
348 Acids Research, 31(13), 3784-3788.

349 Gasteiger, E., Hoogland, C., Gattiker, A., Wilkins, M. R., Appel, R. D., & Bairoch, A. (2005).
350 Protein identification and analysis tools on the ExPASy server. In *The proteomics*
351 *protocols handbook* (pp. 571-607). Humana press.

352 Hall, T., Biosciences, I., & Carlsbad, C. (2011). BioEdit: an important software for molecular
353 biology. *GERF Bull Biosci*, 2(1), 60-61.

354 King, R.W. and Finley, D., 2014. Sculpting the proteome with small molecules. *Nature*
355 chemical biology, 10(11), p.870.

356 Kirchdoerfer, R.N. and Ward, A.B., 2019. Structure of the SARS-CoV nsp12 polymerase
357 bound to nsp7 and nsp8 co-factors. *Nature Communications*, 10(1), pp.1-9.

358 Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: molecular
 359 evolutionary genetics analysis across computing platforms. *Molecular Biology and*
 360 *Evolution*, 35(6), 1547-1549.

361 Kumar, T. A. (2013). CFSSP: Chou and Fasman secondary structure prediction server. *Wide*
 362 *Spectrum*, 1(9), 15-19.

363 Lee, T.W., Cherney, M.M., Huitema, C., Liu, J., James, K.E., Powers, J.C., Eltis, L.D. and
 364 James, M.N., 2005. Crystal structures of the main peptidase from the SARS coronavirus
 365 inhibited by a substrate-like aza-peptide epoxide. *Journal of Molecular Biology*, 353(5),
 366 pp.1137-1151.

367 Luo, C.M., Wang, N., Yang, X.L., Liu, H.Z., Zhang, W., Li, B., Hu, B., Peng, C., Geng, Q.B.,
 368 Zhu, G.J. and Li, F., 2018. Discovery of novel bat coronaviruses in south China that
 369 use the same receptor as Middle East respiratory syndrome coronavirus. *Journal of*
 370 *Virology*, 92(13), pp.e00116-18.

371 Ratia, K., Pegan, S., Takayama, J., Sleeman K., Coughlin, M., Baliji, S., Chaudhuri, R., Fu,
 372 W., Prabhakar, B.S., Johnson, M.E. and Baker, S.C., 2008. A noncovalent class of
 373 papain-like protease/deubiquitinase inhibitors blocks SARS virus replication.
 374 *Proceedings of the National Academy of Sciences*, 105(42), pp.16119-16124.

375 Salentin, S., Schreiber, S., Haupt, V.J., Adasme, M.F. and Schroeder, M., 2015. PLIP: fully
 376 automated protein–ligand interaction profiler. *Nucleic acids research*, 43(W1),
 377 pp.W443-W447

378 Schauer, N.J., Magin, R.S., Liu, X., Doherty, L.M. and Buhrlage, S.J., 2019. Advances in
 379 Discovering Deubiquitinating Enzyme (DUB) Inhibitors. *Journal of medicinal*
 380 *chemistry*.

381 Schwede, T., Kopp, J., Guex, N., & Peitsch, M. C. (2003). SWISS-MODEL: an automated
 382 protein homology-modeling server. *Nucleic Acids Research*, 31(13), 3381-3385.

383 Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F.T.,
 384 de Beer, T.A.P., Rempfer, C., Bordoli, L. and Lepore, R., 2018. SWISS-MODEL:
 385 homology modelling of protein structures and complexes. *Nucleic Acids Research*,
 386 46(W1), pp.W296-W303.

387 World Health Organization (WHO) 2004. [Accessed 11 Feb 2020]
 388 https://www.who.int/csr/don/2004_05_18a/en/

389 World Health Organization (WHO). Coronavirus. Geneva: WHO; 2020 [Accessed 4 Feb
 390 2020]. Available from: <https://www.who.int/health-topics/coronavirus>

391 Yost, S.A. and Marcotrigiano, J., 2013. Viral precursor polyproteins: keys of regulation from
 392 replication to maturation. *Current Opinion in Virology*, 3(2), pp.137-142.