

Super-heavy elements ($Z = 100-109$) treated as calcium clusters: first ionization energies

Robson Fernandes de Farias

Universidade Federal do Rio Grande do Norte, Cx. Postal 1524, 59078-970, Natal-RN, Brasil. robdefarias@yahoo.com.br

Abstract

In the present work, it is shown that the first ionization energies for super-heavy elements ($Z= 100-109$) can be calculated by a semi-empirical method, if the super-heavy elements are treated (modelled) as calcium clusters. The values calculated in the present work are in good agreement compared with those from literature (modelling the super-heavy elements themselves, based on relativistic quantum mechanics).

Keywords: Super-heavy elements, calcium clusters, ionization energy, semi-empirical

Introduction

The search for new chemical elements have resulted in the discovery of the elements 111-118. Since their chemical and physical properties are difficult or even impossible to be accessed/measured directly, the so-called super-heavy elements have been the subject of a lot of theoretical investigations and some works have been dedicated to estimate/predict the properties of such elements [1-6].

Taking into account their higher Z values, trustable calculations involving the properties of super-heavy elements must, necessarily, include the relativistic contributions [7]. However, even today, with the development of faster computers, such calculations are not so easy from an exclusively computational point of view, and only a relatively few researchers/groups are really able/dedicated to them.

In the present work, is shown that the first ionization energies for super-heavy elements ($Z= 100$ -109) can be calculated by a semi-empirical method, if the super-heavy elements are treated as calcium clusters. The values calculated in the present work are compared with those from literature (mainly based on relativistic quantum mechanics calculations) as well experimental values, when available.

Methodology

All calculations were performed by using Spartan [8]. The calcium clusters were modelled as six membered rings. Since calcium has $Z= 20$, five calcium atoms were taken, and another element was “summed up” in order to provide a total of protons equals to the desired element. Hence, for element 103, for example, five calcium ($Z= 20$) atoms and one lithium ($Z= 3$) atom were employed, and the cluster Ca_5Li has a total of 103 protons, as the element 103.

The total number of neutrons is not the same, of course, but is considered here that only the elements charged particles affect the cluster energy levels.

The choice of calcium as the main clusters “building block” was not aleatory: calcium was chose taking into account that ^{40}Ca has a closed nuclear shell with protons and neutrons with the so called “magic configuration”, been close to the line of stability [9]. Furthermore, calcium has been employed as “building block” for experimental super-heavy elements synthesis since calcium is element 20, and calcium plus americium (element 95) yielded element 115, calcium plus curium (element 96) yielded element 116, and so on [10].

All clusters were modelled as neutrals or +1 cations, with the corresponding number of unpaired electrons.

All calculations were performed by semi-empirical (PM6) method. The SE-PM6 approach was chose taking into account its minor computation time consuming and its reliability for calculations involving inorganic systems, as verified for PtF_6 [11] and tin borates [12]. The only exception was the cluster Ca_5He^+ , since PM6 method do not support He. For such cluster, Hartree-Fock (G-3111 + G**) method was employed.

Results and discussion

The obtained results are summarized in Table 1 and compared with calculated values from literature (mainly based on relativistic quantum chemical calculations) and experimental values [13], when available. The obtained equilibrium (minimum energy) structure to Ca_5 (“emulating” element 100) cluster is shown in Figure 1.

Table 1. First ionization energies (eV) for super-heavy elements modelled as calcium clusters.

Element	Z	Modelled cluster	IE/eV _(cluster)	IE/eV _(ref. values)
Fm	100	Ca ₅	4.30	6.50 ^d
		Ca ₅ ⁺	8.70	
			6.50*	
Md	101	Ca ₅ H	5.78	6.58 ^d
		Ca ₅ H ⁺	7.40	
			6.59*	
No	102	Ca ₅ He ⁺	6.70	6.65 ^d
Lr	103	Ca ₅ Li	5.76	4.96 ^e
		Ca ₅ Li ⁺	7.65	
			6.71*	
Rf	104	Ca ₅ Be	4.52	6.01 ^a
		Ca ₅ Be ⁺	9.04	
			6.78*	
Db	105	Ca ₅ B ⁺	7.12	6.9 ^a
Sg	106	Ca ₅ C ⁺	8.44	7.8 ^a
Bh	107	Ca ₅ N ⁺	7.12	7.7 ^a
Hs	108	Ca ₅ O ⁺	7.85	7.6 ^a
Mt	109	Ca ₅ F ⁺	7.33	8.7 ^a

*Mean value of the IE calculated to the neutral and cationic (+) cluster; ^aRef. 9;

^bBased on estimated Clementi effective nuclear charges (Ref. 5); ^c(Ref. 13);

^dCRC Handbook (Ref. 14); ^eExperimental value (Ref. 13);

As can be verified, the SE results are (with exception of element 103) in good (sometimes very good) agreement with the relativistic quantum chemical ones, showing that the employed approach is suitable/reliable. Of course, from times to times, some calculations are improved and a specific value for a given element can change a little from one work to another, but the main result, ie., the agreement between the SE (for calcium clusters) and relativistic (for the elements themselves) values remains.

From elements 100-104 (with exception of No) the “correct” (taking into account a reference value from literature) ionization energy value is the mean of

the value calculated ones to the respective uncharged and cationic (+1) clusters.




Figure 1. Ca_5 cluster structure

This fact suggests that for such elements the “correct” energy level of the valence electron is a mean value of the uncharged and +1 cluster energy levels. In other words, the energy for the valence electron of the super-heavy element is a mean of the HOMO energies for the respective uncharged and cationic (+1) calcium cluster.

Curiously, this happens to the “lighter” (100, 101, 103 and 104 elements) but not the “heavier” 105-109 elements, suggesting that SE methods (calcium clusters) can provide trustable results (that is, comparable to those provided by the relativistic methods -the super-heavy elements themselves) as better as heavier is the considered element.

Of course, only the a profound and detailed analysis of the physical/mathematical “idiosyncrasies” of both theoretical approach can really provide a good explanation for such phenomena.

References

- [1] R.F. de Farias, Estimation of some physical properties for tennessine and tennessine hydride (TsH), *Chem. Phys. Lett.*, 667 (2017) 1-3.
- [2] R.F. de Farias, Estimation of the gas phase formation enthalpies for superheavy-elements (112, 113, 114, 117, 118, 119 and 120) and some of their +1 and -1 ions, *Chem. Res. J.*, 2(3) (2017) 108-111.
- [3] R.F. de Farias, The first and second ionization energies of the element 119: absolute hardness and Mulliken electronegativity for the cation 119+ based on an empirical equation involving absolute hardness, *Mendeleev Commun.*, 28 (2018) 306-307.
- [4] R.F. de Farias, Halides lattice energies and cationic hydration enthalpies for superheavy elements 119 and 120, *J. Atoms and Molecules.*, 8(2) (2018) 1160-1165.
- [5] R.F. de Farias, Estimation of Clementi effective nuclear charges and ionization energies for superheavy elements: explaining the variations for IE along period 7, *J. Atoms and Molecules.*, 8(1) (2018) 1155-1159.
- [6] R.F. de Farias, Estimating the Fermi Energies and Work Functions for the Super Heavy Elements 119 and 120 by using Cationic Absolute Hardness, *Chem. Res. J.*, 3(2) (2018) 56-59.
- [7] B.A. Hess, *Relativistic effects in heavy-element chemistry and physics*, Wiley, Chichester, 2003.
- [8] Wavefunction Inc., Irvine, California, USA.
- [9] M. Schädel, D. Shaughnessy (eds.), *The Chemistry of the Superheavy Elements* (2nd ed.), Springer, Heidelberg, 2014.
- [10] S. Kean, *Science*, A storied Russian lab is trying to push the periodic table past its limits- and uncover exotic new elements, doi:10.1126/science.aaw8425.

- [11] R.F. de Farias, Computational Gas-Phase Formation Enthalpy and Electron Affinity for Platinum Hexafluoride: Is Gaseous PtF_6 Diamagnetic because of a Relativistic Effect?, *Inorg. Chem.*, 55 (23) (2016) 12126-12127.
- [12] R.F. de Farias, On the trustability of semi-empirical methods to the calculation of gas phase formation enthalpies of inorganic compounds containing heavy metals: tin borates, *Mor. J. Chem.*, 6(2) (2018) 256-258.
- [13] T.K. Sato et al., Measurement of the first ionization potential of lawrencium, element 103, *Nature*, 520 (2015) 209-212.
- [14] CRC Handbook of Chemistry and Physics 96th ed, CRC Press - Taylor and Francis, Boca Raton, 2016.