

# Whole Genome Sequence Analysis and Homology Modelling of a 3C Like Peptidase and a Non-Structural Protein 3 of the SARS-CoV-2 Shows Protein Ligand Interaction with an Aza-Peptide and a Noncovalent Lead Inhibitor with Possible Antiviral Properties

Arun K. Shanker<sup>1</sup> \*, Divya Bhanu<sup>1,2</sup> and Anjani Alluri<sup>3</sup>

<sup>1</sup>ICAR - Central Research Institute for Dryland Agriculture

Santoshnagar, Hyderabad – 500059, India

Corresponding author email: [arunshank@gmail.com](mailto:arunshank@gmail.com)

<sup>2</sup> Centre for Plant Molecular Biology, Osmania University, Hyderabad, India

<sup>3</sup> Advanced Post Graduate Centre, Acharya N.G.Ranga Agricultural University, Guntur, India

## Abstract

The family of viruses belonging to Coronaviridae mainly consist of virulent pathogens that have a zoonotic property, Severe Acute Respiratory Syndrome (SARS-CoV) and Middle East Respiratory Syndrome (MERS-CoV) of this family have emerged before and now the SARS-CoV-2 has emerged in China. Characterization of spike glycoproteins, polyproteins and other viral proteins from viruses are important for vaccine development. Homology modelling of these proteins with known templates offers the opportunity to discover ligand binding sites and explore the possible antiviral properties of these protein ligand complexes. Any information emerging from these protein models can be used for vaccine development. In this study we did a complete bioinformatic analysis, sequence alignment, comparison of multiple sequences and homology modelling of the SARS-CoV-2 whole genome sequences, the spike protein and the polyproteins for homology with known proteins, we also analysed receptor binding sites in these models for possible binding with ligands that exhibit antiviral properties. Our results showed that the tertiary structure of the polyprotein isolate SARS-CoV-2\_HKU-SZ-001\_2020 had 98.94 percent identity with SARS-Coronavirus NSP12 bound to NSP7 and NSP8 co-factors. Our results indicate that a part of the viral genome (residues 3268 -3573 in Frame 2 with 306 amino acids) of the SARS-CoV-2 virus isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) when modelled with template 2a5i of the PDB database had 96 percent identity with a 3C like peptidase of SARS-CoV which has ability to bind with Aza-Peptide Epoxide (APE) which is known for irreversible inhibition of SARS-CoV main peptidase. This region was conserved in 250 genomes of SARS-CoV-2. The part of the genome (residues 1568-1882 in Frame 2 with 315 amino acids) when modelled with template 3e9s of the PDB database had 82 percent identity with a papain-like protease/deubiquitinase which when complexed with ligand GRL0617 acts as inhibitor which can block SARS-CoV replication. This region was conserved in 250 genomes of SARS-CoV-2. It is possible that these viral inhibitors can be used for vaccine development for the SARS-CoV-2.

## Introduction

More than a decade has passed since the emergence human Coronavirus that caused Severe Respiratory Syndrome (SARS-CoV) and it is about 7 years since the emergence of another

34 type of Coronavirus - Middle East Respiratory Syndrome (MERS-CoV) and now the SARS-  
35 CoV-2 has emerged in China. This repeated onslaught of these viruses goes to show that it can  
36 assume pandemic proportions at any time and at any place.

37 The family of viruses belonging to Coronaviridae mainly consist of virulent pathogens that  
38 have a zoonotic property and this large family of corona viruses, have been known to be  
39 circulating in animals including camels, cats and bats. It has been seen in the past that Severe  
40 Acute Respiratory Syndrome associated coronavirus (SARS-CoV) and Middle East  
41 Respiratory Syndrome-associated coronavirus (MERS-CoV) belonging to this family of  
42 viruses can be transmitted from animals to humans and can cause respiratory diseases. Human  
43 to human transmission on this virus has been a concern and due to this search for antiviral  
44 compounds and vaccine development for this family of virus becomes the need of the hour.

45 The SARS was first seen in 2002 in Guangdong province of China, and later spread globally  
46 and has caused close to about 8096 cases (WHO 2004, de Wit et al., 2016). In 2012, a novel  
47 betacoronavirus, designated Middle East Respiratory Syndrome coronavirus or MERS-CoV  
48 associated with severe respiratory disease in humans, emerged in the Arabian Peninsula (de  
49 Wit et al., 2013).

50 The World Health Organization (WHO), China Country Office was informed of cases of  
51 pneumonia of unknown aetiology in Wuhan City, Hubei Province, on 31 December 2019  
52 (WHO 2020). A novel coronavirus currently termed SARS-COV-2 was officially announced  
53 as the causative agent by Chinese authorities on 7 January 2020. As on 20 Feb 2020 China's  
54 National Health Commission reported that there are 74,280 confirmed cases in China (Fig.1).  
55 The World Health Organization reported 924 confirmed cases in 25 countries outside China  
56 (WHO Situation Report 29 2020). This novel corona virus has been designated as SARS-CoV-  
57 2.

58

59   Coronaviruses are RNA viruses and have large genomes structures and due to this they can  
60   have high error in replication as compared to host genomes. It is also known that various CoVs  
61   can do effective recombination of their genomes after infecting host cells (Luo et al 2018). This  
62   recombination can be a factor for their evolution to novel types which may have new animals  
63   as their intermediate hosts. These factors give the CoVs high adaptive ability and the capability  
64   to jump across species and have a relatively large host range.

65   Characterization of Spike glycoproteins from viruses are important for vaccine development.  
66   Any information coming from the protein model can be used for vaccine development. *In Silico*  
67   Epitope, polypeptide and spike protein-based peptide vaccine designing for infectious viruses  
68   is a way that can hasten the process of vaccine development. Spike (S) protein, polypeptide and  
69   other viral proteins of the SARS-CoV-2 as a target for the development of vaccines and  
70   therapeutics for the prevention and treatment of infection is an important approach. In the case  
71   of SARS-CoV, these proteins can mediate binding of the virus with its receptor and promotes  
72   the fusion between the viral and host cell membranes and virus entry into the host cell, hence  
73   peptides, antibodies, organic compounds and short interfering RNAs that interact with the spike  
74   protein can have a potential role in vaccine development (Du et al 2009).

75   There are multiple domain functions that are active in the replication of the coronavirus and  
76   these domains are present in a protein designated as Non-structural protein 3 (nsp3) which is  
77   the largest protein in the coronavirus genome (Chen et al 2015). 3C like protease (3CLpro) and  
78   Papain like Protease (PLpro) are two important class of proteases that are involved in the  
79   process of translation of the polypeptide from the genomic RNA to protein components that  
80   are required structurally or non-structurally for replication and packaging of new generation  
81   viruses (Liu et al 2020)

82 The main protease in the SARS virus is the key enzyme for processing of polyproteins of the  
83 virus. This has been the main target for antivirals in the past in SARS-CoV and we hypothesize  
84 that as this has high homology with the main protease of SARS-CoV-2, the same protein can  
85 be a target for antivirals in this virus as well. It has been known that viral replication can be  
86 blocked by inhibiting this protein (Anand et al 2003). The nonexistence of this proteins in  
87 humans makes it an even more attractive antiviral target as there can be no cytotoxicity to  
88 humans.

89 We hypothesised that there can be some proteins in the large chunk of proteins in the SARS-  
90 CoV-2 that could have homology with the Non-structural protein 3 (nsp3) SARS CoV and  
91 these proteins can possibly have binding sites with ligands that can bind with known ligand  
92 with antiviral properties.

93 Here in this study we did a complete bioinformatic analysis, sequence alignment, comparison  
94 of multiple sequences of the SARS-CoV-2 whole genome sequences, the Spike protein and  
95 the polyproteins for homology with known spike proteins and also analysed receptor binding  
96 sites for possible vaccine development.

## 97 **Materials and Methods**

98 Six complete viral genome sequences, seven polyproteins (RdRp region) and seven  
99 glycoproteins available on NCBI portal on 4 Feb 2020 were taken for analysis. The sequence  
100 details and GenBank accession numbers are listed in Supplementary Table 1. Amongst the  
101 seven polyproteins, five are of Wuhan pneumonia virus isolate SARS-COV-2 and two  
102 sequences are of Wuhan pneumonia virus isolate SI200040-SP. The seven Glycoproteins are  
103 of the same isolate, Wuhan pneumonia virus isolate SARS-COV-2.

104 The available polyproteins (RdRp region) and glycoprotein sequences were retrieved from  
105 Genbank, NCBI (Benson et al., 2000). These sequences were translated to amino acid

106 sequences using sorted six frame translation with Bioedit (Hall et al., 2011). Multiple sequence  
107 alignment of the translated protein sequences was performed and phylogenetic tree was  
108 constructed using Mega-X (Kumar et al., 2018). The alignment shows that amongst the seven  
109 polyproteins, five sequences were identical being from the same isolate and two other  
110 sequences of the other isolate are identical. Similar analysis of the seven glycoproteins was  
111 done, all the seven glycoprotein sequences were found to be identical. Therefore, further  
112 analysis was carried out for three sequences.

113 1. MN938385.1 SARS-CoV-2 virus isolate SARS-COV-2 \_HKU-SZ-001\_2020 ORF1ab  
114 polyprotein, RdRp region, (orf1ab) gene, partial cds: 0 to 284: Frame 3 95 aa

115 2. MN970003.1 SARS-CoV-2 virus isolate SI200040-SP orf1ab polyprotein, RdRP  
116 region, (orf1ab) gene, partial cds: 2 to 289: Frame 2 96 aa

117 3. MN938387.1 SARS-CoV-2 virus isolate SARS-COV-2 \_HKU-SZ-001\_2020 surface  
118 glycoprotein (S) gene, partial cds: 1 to 105: Frame 1 35 aa

119 Expasy proteomics server (Gasteiger et al., 2003) was used to study the protein sequence and  
120 structural details. These peptides were studied for their physio-chemical properties using the  
121 tool Protparam (Gasteiger et al., 2005). The secondary structure analysis was done using Chou  
122 and Fasman algorithm with CFSSP (Kumar, 2013). To generate the 3D structure from the fasta  
123 sequence, homology modelling was performed and the templates were identified. The model  
124 was built using the template with highest identity. Swiss-model (Schwede et al., 2003) was  
125 used to build and validate the 3D model, structural assessment was also performed to validate  
126 the model built.

127 Complete genome sequence of the SARS-CoV-2 virus isolate Wuhan-Hu-1 (Genbank  
128 Accession Number MN908947.3) which has 29903 bp ss-RNA linear was translated sorted  
129 6 frame with minimum ORF of 20 with any start codon and the resultant protein sequence was

used for homology modelling, homology models were done with large chunks of proteins 21503 to 25381 in Frame 2 with 1293 amino acids, 13450 to 21552 in Frame 1 with 2701 amino acids and 254 to 13480 in Frame 2 with 4409 amino acids.

SWISS-MODEL server was used for homology modelling (Waterhouse et al 2018) where computation was on ProMod3 engine which is based on Open Structure (Biasini et al 2013). Structural information is extracted from the template, sequence alignment is used to define insertions and deletions.

Protein ligand interaction profile with hydrogen bonding, hydrophobic interactions, salt bridges and  $\pi$ -Stacking was done with PLIP server (Salentin et al., 2015)

## **Results and Discussion**

The physico- chemical properties and primary structure parameters of the 7 polyproteins RdRp region of the SARS-CoV-2 virus isolate is given in Table 1. RdRP forms an important part of the viral genome where in the RNA viruses its function is to catalyze the synthesis of the RNA strand complementary to a given RNA template.

The isolates SI200040-SP orflab polyprotein and the isolate SI200121-SP orflab polyprotein had 2 reading frames as compared to the rest of the isolates which had 3 reading frames. The presence of multiple reading frames suggests the possibility of overlapping genes as seen in many virus and prokaryotes and mitochondrial genomes. This could affect how the proteins are made. The number of amino acid residues in all the polyproteins were the same except one isolate SI200040-SP which had one amino acid more than the other polyproteins. The extinction coefficients of the two isolates SI200040-SP orflab polyprotein and the isolate SI200121-SP orflab polyprotein was much higher compared to the rest of the polyproteins. The extinction coefficient is important when studying protein-protein and protein-ligand interactions. The instability index of these two isolates was also high when compared to the

others indicating the that these two isolates are instable. Regulation of gene expression by polyprotein processing is known in viruses and this is seen in many viruses that are human pathogens (Yost et al 2013).

The isolates here like many other viruses may be using replication strategy which could involve the translation of a large polyprotein with subsequent cleavage by viral proteases. The two isolates SI200040-SP orflab polyprotein and the isolate SI200121-SP orflab polyprotein also showed shorter half-lives as compared to the other isolates indicating that they are susceptible to enzymatic degradation.

The tertiary structure analysis of the isolate SARS-CoV-2 \_HKU-SZ-001\_2020 ORF1ab polyprotein is given in Table 2. It is seen that the polyprotein has a 98.94 percent identity with PDB structure 6nur.1.A which is a hetero-1-2-1-mer. The polyprotein is an RNA directed RNA polymerase. The protein is identical to the SARS-Coronavirus NSP12 bound to NSP7 and NSP8 co-factors (Kirchdoerfer and Ward 2019). In SARS it is basically a nonstructural protein with NSP12 being the RNA dependent RNA polymerase and the co factors NSP 7 and NSP 8 having the function of forming hexadecameric complexes and also act as processivity clamp for RNA polymerase and primase (Fehr et al., 2016). This structure as in SARS CoV here in SARS-CoV-2 may be involved in the machinery of core RNA synthesis and can be a template for exploring antiviral properties.

The phylogenetic tree of the seven polyproteins is shown in Fig.2. It is seen that two polyproteins were distinctly different from the rest. The Phylogenetic tree of the seven glycoproteins of the SARS-CoV-2 virus isolate is shown in Fig.3, it is seen that the glycoproteins are similar in all the isolates. Multiple alignment of the Polyproteins of the SARS-CoV-2 is shown in Supplementary Fig.1.

This structure as in SARS CoV here in SARS-CoV-2 may be involved in the machinery of core RNA synthesis and can be a template for exploring antiviral properties. Based on its functions

in the SARS CoV and its identity to the SARS-CoV-2, it is possible that it has the same functions in SARS-CoV-2 an RNA polymerase which does de novo initiation and primer extension with possible exonuclease activities, the activity itself being primer dependent useful for understanding the mechanism of SARS-CoV-2 replication and can be used as an antiviral target (Te Velhuis et al 2012; Te Velhuis et al 2010; Subissi et al 2014; Subissi et al 2014).

The polyprotein also has an identity of 19.74 percent with an ABC-type uncharacterized transport system periplasmic component-like protein, this protein is known to be a substrate binding protein and possible binding can be explored here (Bae et al 2019).

The homology model developed from the residues 254 to 13480 in Frame 2 with 4409 amino acids from the Complete genome sequence of the SARS-CoV-2 virus isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) which has 29903 bp with linear ss-RNA linear showed interesting template alignments, in all the model aligned with 50 templates from the PDB database with most of them being replicase polyprotein 1ab which is a SARS-CoV papain-like protease (Daczkowski 2017). The maximum similarity of 97.3 percent was with template structure of a Nsp9 protein from SARS-coronavirus indicating that this novel coronavirus has high degree of similarity with the SARS-coronavirus and this can be used for gaining insights into vaccine development. Nsp 9 is an RNA binding protein and has an oligosaccharide/oligonucleotide fold-like fold, this protein can have an important function in the replication machinery of the virus and can be important when designing antiviral for this virus (Egloff et al 2004).

Two models were developed, one from residues 3268 -3573 in Frame 2 with 306 amino acids and the other from the part of the genome residues 1568-1882 in Frame 2 with 315 amino acids of the SARS-CoV-2 virus isolate Wuhan-Hu-1 (Genbank Accession Number



203 MN908947.3). The models had similarity with the 3C like proteinase and a papain-like  
204 protease/deubiquitinase protein which are known antiviral drug targets. Ligand binding with  
205 these proteins and their action is on viral replication and inactivation can be useful in stopping  
206 the viral replication (Baez-Santos et al 2015).

207

208 The homology models of the 4409 amino acid residues of the whole genome of the SARS-  
209 CoV-2 virus isolate Wuhan-Hu-1 with the ligand association with templates 2a5i and 3e9s are  
210 shown in Fig. 4 and Fig. 5 respectively.

211 The statistics of structural comparison with PDB templates is given in Table 5, it is seen that  
212 the proteins from the SARS-CoV-2 are significantly close to the proteins of SARS CoV and  
213 the amino acid alignment in the binding region is the same in both the viruses.

214 The alignment of the 305 residues from 3268-3573 aa of the Novel Coronavirus COVI-19 with  
215 the template 2a5i is shown in Fig.6 and the alignment of the 315 residues from 1568-1882 aa  
216 of the Novel Coronavirus COVI-19 with the template 3e9s is shown in Fig.7.

217 A PSI-BLAST of a length of 306 amino acid residues 3268 -3573 in Frame 2 from the SARS-  
218 CoV-2 virus isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) was conducted  
219 to ascertain the conservation of these amino acids in 250 genome sequences of SARS-CoV-2  
220 and it was found that there was a complete match in these genomes of the virus. The fact that  
221 the region is conserved in all these SARS-CoV-2 sequences further emphasizes this ligand  
222 interaction of Aza-Peptide epoxide with the protein can be used as an antiviral in SARS-CoV-  
223 2. Similarly A PSI-BLAST of a length of 315 amino acid residues 3268 -3573 in Frame 2  
224 with 315 amino acid residues 1568-1882 in Frame 2 from SARS-CoV-2 virus isolate Wuhan-  
225 Hu-1 (Genbank Accession Number MN908947.3) was conducted to ascertain the conservation  
226 of these amino acids in 250 genome sequences of SARS-CoV-2 and it was found that there

was a complete match in these genomes of the virus. The fact that the region is conserved in all these SARS-CoV-2 sequences further emphasizes this ligand interaction of ligand GRL0617 with the protein can be used as an antiviral in SARS-CoV-2.

The important templates that aligned with this 4409 amino acid residues of the whole genome of the SARS-CoV-2 virus isolate Wuhan-Hu-1 were 2a5i of the PDB database which is a crystal structure of SARS coronavirus main peptidase inhibited by an Aza-Peptide epoxide in the space group C2 (Lee et al 2005) and 3e9s of the PDB database which is new class of papain-like protease/deubiquitinase which when combined with ligand GRL0617 acts as inhibitors blocking SARS virus replication (Ratia et al 2008). The model with template 2a5i of the PDB database shows that Aza-Peptide Epoxide (APE;  $k_{inact}/K_i=1900(\pm 400) \text{ M}^{-1} \text{ s}^{-1}$ ) which is a known anti SARS agent can be used to develop a molecular target with irreversible inhibitor properties. The protein ligand interaction analysis of the Novel Coronavirus C3 like peptidase and aza-peptide epoxide is shown in Fig.8. The substrate binding properties and structural and chemical complementarity of this Aza-Peptide Epoxide can be explored as an anti - Coronavirus SARS-COV-2 agent. The APE which is ethyl (2S)-4-[(3-amino-3-oxo-propyl)-[[[(2S)-2-[[[(2S)-4-methyl-2-phenylmethoxycarbonylamino-pentanoyl]amino]-3-phenyl-propanoyl]amino]amino]-2-hydroxy-4-oxo-butanoate structure is shown in Fig.9.

The model with template 3e9s of the PDB database shows that the Coronavirus viral protein can have a ligand which is a papain-like protease (PLpro) that is known to be a potent inhibitor of viral replication in SARS (Ratia et al 2008).

The two parts of the Main protein from the whole genome of the SARS-CoV-2 aligned with two SAR proteins and the ligand binding sites were similar, the alignment positions, number of amino acids and ligand and the interacting residues is given in Table 3

251 The complete genome of MN908947.3 SARS-CoV-2 virus isolate Wuhan-Hu-1 encodes a  
252 4409aa long protein along with the other glycoproteins and polyproteins. The homology  
253 modelling of this protein showed sequence and structural alignment with two SARS proteases  
254 with structural accession numbers 3e9s.1 and 2a5i.1 at positions 1568-1882 and 3268-3573  
255 respectively. Reports suggests inhibition of virus replication by TTT ligand and an aza-peptide  
256 epoxide inhibiting the main peptidase. The structural similarity of these templates are 83% and  
257 96% respectively. The multiple sequence alignment shows complete conservation of the  
258 sequence suggesting a high degree of homology. The protein ligand interaction analysis of the  
259 Novel Coronavirus non structural protein and papain-like protease is shown in Fig. 10.

260 The Comparison of Hydrophobic interaction, hydrogen bonding, salt bridges of the  
261 constructed model of the Novel Coronavirus protein from region 3268-3573 aa to ligand AZP  
262 with Hydrophobic interaction, hydrogen bonding, salt bridges of the template 2a5i is given in  
263 Suppl. Table 2, when comparing both it is seen that the binding properties are the same expect  
264 for the presence of water bridge in the template 2a5i.

265

266 The Comparison of Hydrophobic interaction, hydrogen bonding,  $\pi$ -Stacking of the constructed  
267 model of the Novel Coronavirus protein from region 1568-1882 aa to ligand Small molecule  
268 Noncovalent Lead Inhibitor with the Hydrophobic interaction, hydrogen bonding,  $\pi$ -Stacking  
269 of the template 3e9s is given in Suppl. Table 3, when comparing both it is seen that the binding  
270 properties are the same except or an additional  $\pi$ -Stacking at Tyr in the template 2a5i. This  
271 shows that there is high possibility of binding of these antiviral compounds with the regions of  
272 Novel Coronavirus protein that is in homology with the SARS protein.

273 Comparison of the hydrophobic interaction of the biding of the ligand AZP between the SARS-  
274 CoV-2 protein and the template 2a5i of SARS CoV is shown in Fig.11 and the comparison of

275 the hydrophobic interaction of the binding of the ligand AZP between the SARS-CoV-2 protein  
276 and the template 3e9s of SARS CoV is shown in Fig.12. It is seen that the interaction is the  
277 same in both proteins with the same amino acids participating in the interaction indicating that  
278 there is a possibility that these ligands with antiviral properties can bind to the new virus.

279 The similarity in the amino acids involved in the Hydrophobic interactions which are short range  
280 interactions and have an important role in the affinities of the ligands and receptors shows that  
281 the proteins of the SARS-CoV-2 may bind with the same affinity as seen in the SARS CoV  
282 and this also shows a similar action of the ligand as seen in SARS CoV, indicating that these  
283 ligands can be used as antivirals in the SARS-CoV-2.

284 The targeting of this part of the genome of the SARS-CoV-2 with the antiviral compounds  
285 which have shown to bind in the similar region of the SARS virus can have implication in the  
286 development of an effective antiviral compound against the SARS-CoV-2. The SARS-CoV-2  
287 shows homology with the SARS coronaviral proteases, papain-like protease (PLpro) and 3C-  
288 like protease (3CLpro), these proteins have the function of processing the viral polyprotein and  
289 also they perform the function of stripping ubiquitin and the ubiquitin-like interferon (IFN)-  
290 stimulated gene 15 (ISG15) from the hosts to facilitate coronavirus replication and help in  
291 evading immune response of the host, these inhibitors can also have a role in disrupting  
292 signalling cascades in infected cells and protecting the uninfected cells.

293 The chemical GRL0617 is 5-Amino-2-methyl-N-[(1R)-1-(1-naphthalenyl)ethyl]benzamide  
294 and is known to inhibit the papainlike protease that is present in SARS CoV. This protease is  
295 a potential target for antiviral compounds (Chaudhuri et al., 2011). We found the SARS-CoV-  
296 2 has homology with this and the binding sites for this in the structural protein of the SARS-  
297 CoV-2 is the same (Table 4). This compound inhibits the enzyme that is required for the  
298 cleavage of the viral protein from the virus in SARS CoV, it also cleaves ubiquitin and has a

structural homology with the Deubiquitinases (DUBs) of the Ubiquitin-Specific Proteases  
Compound GRL0617 binds in the S4 and S3 enzyme subsite that gets the C terminal tail of the  
Ubiquitin (King and Finley 2014; Schauer et al., 2019). Our results show that Aza-Peptide  
Epoxide an irreversible protease inhibitor and GRL0617 a viral replication inhibitor can be  
used to develop inhibitors of the Novel Coronavirus SARS-COV-2.

## References

- Anand, K., Ziebuhr, J., Wadhwani, P., Mesters, J.R. and Hilgenfeld, R., 2003. Coronavirus  
main proteinase (3CLpro) structure: basis for design of anti-SARS drugs. *Science*,  
300(5626), pp.1763-1767.
- Bae, J.E., Kim, I.J., Kim, K.J. and Nam, K.H., 2018. Crystal structure of a substrate-binding  
protein from *Rhodothermus marinus* reveals a single  $\alpha/\beta$ -domain. *Biochemical and  
Biophysical Research Communications*, 497(1), pp.368-373.
- Baez-Santos, Y.M., John, S.E.S. and Mesecar, A.D., 2015. The SARS-coronavirus papain-like  
protease: structure, function and inhibition by designed antiviral compounds. *Antiviral  
Research*, 115, pp.21-38.
- Benson, D. A., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., Rapp, B. A., & Wheeler, D. L.  
(2000). GenBank. *Nucleic Acids Research*, 28(1), 15-18.
- Biasini, M., Schmidt, T., Bienert, S., Mariani, V., Studer, G., Haas, J., Johnner, N., Schenk,  
A.D., Philippsen, A. and Schwede, T., 2013. OpenStructure: an integrated software  
framework for computational structural biology. *Acta Crystallographica Section D:  
Biological Crystallography*, 69(5), pp.701-709.

321 Chaudhuri, R., Tang, S., Zhao, G., Lu, H., Case, D.A. and Johnson, M.E., 2011. Comparison  
322 of SARS and NL63 papain-like protease binding sites and binding site dynamics:  
323 inhibitor design implications. *Journal of molecular biology*, 414(2), pp.272-288.

324 Chen, Y., Savinov, S.N., Mielech, A.M., Cao, T., Baker, S.C. and Mesecar, A.D., 2015. X-ray  
325 structural and functional studies of the three tandemly linked domains of non-structural  
326 protein 3 (nsp3) from murine hepatitis virus reveal conserved functions. *Journal of*  
327 *Biological Chemistry*, 290(42), pp.25293-25306.

328 Daczkowski, C.M., Dzimianski, J.V., Clasman, J.R., Goodwin, O., Mesecar, A.D. and Pegan,  
329 S.D., 2017. Structural insights into the interaction of coronavirus papain-like proteases  
330 and interferon-stimulated gene product 15 from different species. *Journal of Molecular*  
331 *Biology*, 429(11), pp.1661-1683.

332 de Wit, E., Rasmussen, A.L., Falzarano, D., Bushmaker, T., Feldmann, F., Brining, D.L.,  
333 Fischer, E.R., Martellaro, C., Okumura, A., Chang, J. and Scott, D., 2013. Middle East  
334 respiratory syndrome coronavirus (MERS-CoV) causes transient lower respiratory tract  
335 infection in rhesus macaques. *Proceedings of the National Academy of Sciences*,  
336 110(41), pp.16598-16603.

337 de Wit, E., van Doremalen N., D. Falzarano, V. J. Munster, SARS and MERS: recent insights  
338 into emerging coronaviruses. *Nat Rev Microbiol* 14, 523-534 (2016).

339 Du, L., He, Y., Zhou, Y., Liu, S., Zheng, B.J. and Jiang, S., 2009. The spike protein of SARS-  
340 CoV—a target for vaccine and therapeutic development. *Nature Reviews*  
341 *Microbiology*, 7(3), pp.226-236.

342 Egloff, M.P., Ferron, F., Campanacci, V., Longhi, S., Rancurel, C., Dutartre, H., Snijder, E.J.,  
343 Gorbalenya, A.E., Cambillau, C. and Canard, B., 2004. The severe acute respiratory  
344 syndrome-coronavirus replicative protein nsp9 is a single-stranded RNA-binding

345 subunit unique in the RNA virus world. Proceedings of the National Academy of  
 346 Sciences, 101(11), pp.3792-3796.

347 Fehr, A.R. and Perlman, S., 2015. Coronaviruses: an overview of their replication and  
 348 pathogenesis. In Coronaviruses (pp. 1-23). Humana Press, New York, NY.

349 Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R. D., & Bairoch, A. (2003).  
 350 ExPASy: the proteomics server for in-depth protein knowledge and analysis. Nucleic  
 351 Acids Research, 31(13), 3784-3788.

352 Gasteiger, E., Hoogland, C., Gattiker, A., Wilkins, M. R., Appel, R. D., & Bairoch, A. (2005).  
 353 Protein identification and analysis tools on the ExPASy server. In *The proteomics*  
 354 *protocols handbook* (pp. 571-607). Humana press.

355 Hall, T., Biosciences, I., & Carlsbad, C. (2011). BioEdit: an important software for molecular  
 356 biology. GEF Bull Biosci, 2(1), 60-61.

357 King, R.W. and Finley, D., 2014. Sculpting the proteome with small molecules. Nature  
 358 chemical biology, 10(11), p.870.

359 Kirchdoerfer, R.N. and Ward, A.B., 2019. Structure of the SARS-CoV nsp12 polymerase  
 360 bound to nsp7 and nsp8 co-factors. Nature Communications, 10(1), pp.1-9.

361 Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: molecular  
 362 evolutionary genetics analysis across computing platforms. Molecular Biology and  
 363 Evolution, 35(6), 1547-1549.

364 Kumar, T. A. (2013). CFSSP: Chou and Fasman secondary structure prediction server. Wide  
 365 Spectrum, 1(9), 15-19.

366 Lee, T.W., Cherney, M.M., Huitema, C., Liu, J., James, K.E., Powers, J.C., Eltis, L.D. and  
 367 James, M.N., 2005. Crystal structures of the main peptidase from the SARS coronavirus

368 inhibited by a substrate-like aza-peptide epoxide. *Journal of Molecular Biology*, 353(5),  
 369 pp.1137-1151.

370 Liu, W., Morse, J.S., Lalonde, T. and Xu, S., 2020. Learning from the Past: Possible Urgent  
 371 Prevention and Treatment Options for Severe Acute Respiratory Infections Caused by  
 372 2019-nCoV. *ChemBioChem*.

373 Luo, C.M., Wang, N., Yang, X.L., Liu, H.Z., Zhang, W., Li, B., Hu, B., Peng, C., Geng, Q.B.,  
 374 Zhu, G.J. and Li, F., 2018. Discovery of novel bat coronaviruses in south China that  
 375 use the same receptor as Middle East respiratory syndrome coronavirus. *Journal of*  
 376 *Virology*, 92(13), pp.e00116-18.

377 Ratia, K., Pegan, S., Takayama, J., Sleeman K., Coughlin, M., Baliji, S., Chaudhuri, R., Fu,  
 378 W., Prabhakar, B.S., Johnson, M.E. and Baker, S.C., 2008. A noncovalent class of  
 379 papain-like protease/deubiquitinase inhibitors blocks SARS virus replication.  
 380 *Proceedings of the National Academy of Sciences*, 105(42), pp.16119-16124.

381 Salentin, S., Schreiber, S., Haupt, V.J., Adasme, M.F. and Schroeder, M., 2015. PLIP: fully  
 382 automated protein–ligand interaction profiler. *Nucleic acids research*, 43(W1),  
 383 pp.W443-W447

384 Schauer, N.J., Magin, R.S., Liu, X., Doherty, L.M. and Buhrlage, S.J., 2019. Advances in  
 385 Discovering Deubiquitinating Enzyme (DUB) Inhibitors. *Journal of medicinal*  
 386 *chemistry*.

387 Schwede, T., Kopp, J., Guex, N., & Peitsch, M. C. (2003). SWISS-MODEL: an automated  
 388 protein homology-modeling server. *Nucleic Acids Research*, 31(13), 3381-3385.



389 Subissi, L., Imbert, I., Ferron, F., Collet, A., Coutard, B., Decroly, E. and Canard, B., 2014.  
390 SARS-CoV ORF1b-encoded nonstructural proteins 12–16: replicative enzymes as  
391 antiviral targets. *Antiviral research*, 101, pp.122-130.

392 Subissi, L., Posthuma, C.C., Collet, A., Zevenhoven-Dobbe, J.C., Gorbalenya, A.E., Decroly,  
393 E., Snijder, E.J., Canard, B. and Imbert, I., 2014. One severe acute respiratory syndrome  
394 coronavirus protein complex integrates processive RNA polymerase and exonuclease  
395 activities. *Proceedings of the National Academy of Sciences*, 111(37), pp.E3900-  
396 E3909.

397 Te Velhuis, A.J., Arnold, J.J., Cameron, C.E., van den Worm, S.H. and Snijder, E.J., 2010.  
398 The RNA polymerase activity of SARS-coronavirus nsp12 is primer dependent.  
399 *Nucleic acids research*, 38(1), pp.203-214.

400 Te Velhuis, A.J., van den Worm, S.H. and Snijder, E.J., 2012. The SARS-coronavirus nsp7+  
401 nsp8 complex is a unique multimeric RNA polymerase capable of both de novo  
402 initiation and primer extension. *Nucleic acids research*, 40(4), pp.1737-1747.


403 Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F.T.,  
404 de Beer, T.A.P., Rempfer, C., Bordoli, L. and Lepore, R., 2018. SWISS-MODEL:  
405 homology modelling of protein structures and complexes. *Nucleic Acids Research*,  
406 46(W1), pp.W296-W303.

407 World Health Organization (WHO) 2004. [Accessed 11 Feb 2020]  
408 [https://www.who.int/csr/don/2004\\_05\\_18a/en/](https://www.who.int/csr/don/2004_05_18a/en/)


409 World Health Organization (WHO). Coronavirus. Geneva: WHO; 2020 [Accessed 4 Feb  
410 2020]. Available from: <https://www.who.int/health-topics/coronavirus>

Yost, S.A. and Marcotrigiano, J., 2013. Viral precursor polyproteins: keys of regulation from replication to maturation. Current Opinion in Virology, 3(2), pp.137-142.


**Figure Captions**



**Fig.1** Countries, territories or areas with reported confirmed cases of SARS-COV-2 , 3 February 2020 Source WHO ([https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200219-sitrep-30-covid-19.pdf?sfvrsn=6e50645\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200219-sitrep-30-covid-19.pdf?sfvrsn=6e50645_2))




**Fig.2** Phylogenetic tree of the seven polyproteins of Severe acute respiratory syndrome coronavirus 2 isolate virus isolates



**Fig.3** Phylogenetic tree of the seven glycoproteins of Severe acute respiratory syndrome coronavirus 2 isolate virus isolates

427



428

429


430 **Fig. 4** Homology model with ligand binding of protein from amino acids 3268 -3573 in Frame  
431 2 with 306 amino acids of the Complete genome sequence of the SARS-CoV-2 virus  
432 isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) which has 29903 bp  
433 linear ss-RNA with 2a5i of the PDB database as template.

434

435

436

437



438

439 **Fig. 5** Homology model with ligand binding of protein from residues 1568-1882 in Frame 2  
440 with 315 amino acids of the Complete genome sequence of the SARS-CoV-2 virus  
441 isolate Wuhan-Hu-1 (Genbank Accession Number MN908947.3) which has 29903 bp  
442 linear ss-RNA with 3e9s of the PDB database as template.

443

444

445

```

Seqres  SGFRKMAFPSSGKVEGCMVQVTCGTTTLNGLWDDTVYCPRHVICTAEDMLNPNYEDLLIRKSNHSFLVQAGNVQLRVIGH 80
2a5i.1. (AB) SGFRKMAFPSSGKVEGCMVQVTCGTTTLNGLWDDTVYCPRHVICTAEDMLNPNYEDLLIRKSNHSFLVQAGNVQLRVIGH 80

Seqres  SMQNCCLRLKYDTSNPKTFKYKFVRIQPGQTFSVLACYNGSPSGVYQCAMRPNHTIKGSFLNGSCGSVGFNIIDYDCVSFC 160
2a5i.1. (AB) SMQNCCLRLKYDTSNPKTFKYKFVRIQPGQTFSVLACYNGSPSGVYQCAMRPNHTIKGSFLNGSCGSVGFNIIDYDCVSFC 160

Seqres  YMHMELPTGVHAGTDLEGKFYGFVDRQTAQAAGTDTITLNVLANLYAAVINGDRWFLNRFTTTLNDFNLVAMKYNFY 240
2a5i.1. (AB) YMHMELPTGVHAGTDLEGKFYGFVDRQTAQAAGTDTITLNVLANLYAAVINGDRWFLNRFTTTLNDFNLVAMKYNFY 240

Seqres  PLTQDHVDILGFLSAQTGIAVLDMCAALKELLQNGMNGRTILGSTILEDEFTPFDDVVRQCSSGVTFQ 306
2a5i.1. (AB) PLTQDHVDILGFLSAQTGIAVLDMCAALKELLQNGMNGRTILGSTILEDEFTPFDDVVRQCSSGVTFQ 306

```

446

447 **Fig. 6** Alignment of the 305 residues from 3268-3573 aa of the Novel Coronavirus COVI-19  
 448 with the template 2a5i

```

Seqres  ASMEVKTIKVFTTYDNTNLHTQLVDMSTYGGQFGFTYLDGADVTIKIKPHYNHEGKTFEVLPSDDTLRSEAFEYYHTLDESFLGR 85
3e9s.1.A ASMEVKTIKVFTTYDNTNLHTQLVDMSTYGGQFGFTYLDGADVTIKIKPHYNHEGKTFEVLPSDDTLRSEAFEYYHTLDESFLGR 85

Seqres  YMSALNHTKKWKFPQVGGTTSIKWADNNCYLSSVLLALQQLLEVAFNAPALQEAYYRARAGDAANFCALILAYSNNKTVGELGDVRE 170
3e9s.1.A YMSALNHTKKWKFPQVGGTTSIKWADNNCYLSSVLLALQQLLEVAFNAPALQEAYYRARAGDAANFCALILAYSNNKTVGELGDVRE 170

Seqres  TMTHLLQHANLESARKVNLNVVCKKCGQKTTTLTGVEAVMYMGTLSDNLKTVSIPCVCGRDATQYLVDQESSFVMM$APPAEYK 255
3e9s.1.A TMTHLLQHANLESARKVNLNVVCKKCGQKTTTLTGVEAVMYMGTLSDNLKTVSIPCVCGRDATQYLVDQESSFVMM$APPAEYK 255


Seqres  LQOGTFLCANEYTGNYQCCHYTHITAKETLYRIDGAHLTKMSEYKGPVTDVFKETSXTTIR 318
3e9s.1.A LQOGTFLCANEYTGNYQCCHYTHITAKETLYRIDGAHLTKMSEYKGPVTDVFKETSXTTIR 317

```

449


450 **Fig.7** the alignment of the 315 residues from 1568-1882 aa of the Novel Coronavirus COVI-  
 451 19 with the template 3e9s





452


453 **Fig.8** Protein Ligand interaction between the C3 like peptidase with aza-peptide epoxide of the  
 454 model with the template 2a5i




455

456 **Fig. 9** Structure of Aza-Peptide Epoxide (APE) ethyl (2S)-4-[(3-amino-3-oxo-propyl)-[[[(2S)-  
 457 2-[[[(2S)-4-methyl-2-phenylmethoxycarbonylamino-pentanoyl]amino]-3-phenyl-  
 458 propanoyl]amino]amino]-2-hydroxy-4-oxo-butanoate with possible anti Coronavirus  
 459 activity – (Source <https://www.rcsb.org/ligand/AZP>)





**Fig.10** Protein Ligand interaction between the Novel Coronavirus non structural protein and papain-like protease of the model with the template 3e9s




**Fig.11** Comparison of the hydrophobic interaction of the binding of the ligand AZP between the SARS-CoV-2 protein and the template 2a5i of SARS CoV

486

487

488



489

490 **Fig.12** comparison of the hydrophobic interaction of the binding of the ligand AZP between the  
491 SARS-CoV-2 protein and the template 3e9s of SARS CoV

492

493

