

The Chemoselective C–C σ -Bond Activation of the Arene Ring of Biphenylene

Richard Y. Kong, Mark R. Crimmin*

Department of Chemistry, Molecular Sciences Research Hub, Imperial College London, White City, London, W12 0BZ, UK.


ABSTRACT: The chemoselective cleavage of an arene ring in biphenylene is reported using an aluminium(I) complex. The reaction proceeds with complete integrity of the central 4-membered ring despite this ring containing the weakest C–C σ -bond in the hydrocarbon scaffold. A reaction intermediate derived from the (4+1) cycloaddition of the aluminium(I) complex to the π -system of biphenylene was isolated. Further experiments and DFT calculations suggest that this intermediate is involved in breaking of the C–C σ -bond. Activation strain analysis was used to understand the origins of the remarkable chemoselectivity of this system. Both the symmetry and diffuseness of the frontier molecular orbitals of the aluminium(I) fragment are implicated in its unusual reactivity with biphenylene.

Reactions that break the strong C–C σ -bonds of hydrocarbons are finding increasing use in synthesis.^{1,2} When more than one type of C–C σ -bond is present in the substrate, chemoselectivity becomes a central consideration. In nearly all cases, substrate bias dictates which C–C σ -bond breaks. Structural features such as strained ring systems prejudice the reagent toward specific sites and determine the chemoselectivity of the reaction. Examples where reagent control overrides substrate bias are essentially unknown.


Biphenylene has been the subject of detailed study in C–C σ -bond activation. The central C–C σ -bond of biphenylene has a low bond dissociation energy (BDE) of 65.4 kcal mol⁻¹.³ Both the antiaromatic character and strain of the four-membered ring contribute to the weakening of this C–C σ -bond. If biphenylene undergoes C–C σ -bond activation, this weakest link is always the site of reactivity. For example, the oxidative addition of a range of transition metal complexes to the central ring of biphenylene has been studied in great detail.^{4–7} More recently, the reaction of the central C–C σ -bond with an anionic aluminium(I) nucleophile has also been reported (Scheme 1a).⁸

The selectivity is unsurprising considering that alternative reactive sites of biphenylene require the dearomatisation of a benzene ring. The C–C bond of benzene is strong and is enforced by a combination of σ - and π -bonding. There are a number of transition metal^{9–16} and main group¹⁷ complexes that can effect the C–C σ -bond activation and dearomatisation of benzene units. Mechanistically, these systems show some common features: coordination of the metal complex to the arene π -manifold generates a strained metallocyclic intermediate from which C–C σ -bond activation evolves to form the observed products. None of these systems have been shown to react with benzene units in the presence of more strained ring systems containing weaker C–C σ -bonds.

We recently reported the activation of C–C σ -bonds of alkylidene cyclopropanes using the aluminium(I) complex, **1** (Scheme 1b).¹⁸ Here we demonstrate σ -C–C bond activation of biphenylene using **1**. Despite the substrate bias of biphenylene, the reaction is under reagent control and occurs with complete chemoselectivity for the strongest C–C



Scheme 1: a) Summary of previous studies on the C–C σ -bond activation of biphenylene. b) Ring-opening of alkylidene cyclopropanes by **1**. c) The work presented in this paper. Activated C–C σ -bonds are highlighted. Dip = 2,6-di-isopropylphenyl.



Scheme 2: C–C σ -Bond activation of biphenylene with **1**. Dip = 2,6-di-isopropylphenyl.

σ -bonds of the six-membered arene ring. The weakest C–C σ -bond of biphenylene remains intact through the reaction. We provide a clear mechanistic rationale for this selectivity that involves an initial reaction of **1** with the π -system of biphenylene.

The reaction of 2 equiv. of **1** with biphenylene at 100°C for 3 h yields a mixture of **2** and **3** in a 1:0.7 ratio (Scheme 2). **2** and **3** are derived from the cleavage of the C²–C³ and C⁴–C⁵ bonds in biphenylene respectively. Both products incorporate metallocycles in which aluminium(III) centers are in 5-membered rings. **2** crystallises readily from a concentrated toluene or THF solution of the reaction mixture at –35°C. **3** can be separated from **2** by extracting with THF and purified by recrystallising from toluene. Heating isolated samples of either **2** or **3** result in their decomposition and no interconversion is observed between these two compounds in either purified or crude mixtures.

Repeating the reaction of **1** with biphenylene at lower temperatures allowed the isolation of a reaction intermediate. After 7 days at 25°C addition of **1** to biphenylene yields **4** as a major product (Scheme 2). **4** is derived from a (4+1) cycloaddition of **1** to the C¹ and C⁴ positions of an arene ring of biphenylene. Based on cross-over experiments, the formation of **4** is, at least, partially reversible (See Supporting Information for further details). The formation of **4** can be explained by a pericyclic reaction, specifically a $[\pi 4_s + \pi 2_s]$ cycloaddition.¹⁹ Formation of **4** generates a highly strained and dearomatized hydrocarbon framework that is primed for further reactivity. Addition of a further equiv. of **1** to **4** and heating to 100°C generates the **2** and **3** with similar yields and selectivity to that observed directly from **1** and biphenylene.

All products have been characterised by multinuclear NMR spectroscopy and single-crystal X-ray diffraction. The connectivity of **2** is evident in the solid-state structure however the ring-opened biphenylene unit is disordered over four sites (See Supporting Information for further discussion). Regardless, the ¹H NMR spectrum is consistent with the structural assignment. Characteristic alkene signals for the protons

attached to C⁵ and C⁶ resonate at δ_H = 5.32 and 7.44 ppm respectively. Compound **2** contains two contiguous stereocentres at the C⁵ and C⁶ position. The relative stereochemistry is determined by the *cis*-fused ring junction and has been confirmed by 1D NOE experiments. The crystal structure of **3** clearly shows the rupture of the aromatic ring (Figure 1a). The aluminium centres of **3** are related by C₂-rotation around the perpendicular bisector of the C¹–C² bond. This compound again contains two contiguous stereocentres, now at the C¹ and C² positions, which impose a *trans*-fused ring junction and C₂-symmetry.




Figure 1: The structure of (a) **3** and (b) **4** determined by single crystal X-ray diffraction.




Figure 2: Calculated reaction pathway for the formation of **2** and **3** from **1** and biphenylene. Energies in kcal mol⁻¹. All calculations performed at the M06/6-31G** (C, H, N)/SDDAll (Al) level of theory with single point dispersion (D3) and solvent (benzene) corrections.

DFT calculations were used to explore the mechanism of C–C σ -bond activation (Figure 1). The (4 + 1) cycloaddition of **1** to biphenylene was found to proceed *via* a single transition-state **TS-1** ($\Delta G^\ddagger_{298K} = 28.5$ kcal mol⁻¹) and forms **4** directly. Disruption of the antiaromaticity of the central ring of biphenylene likely determines the selectivity of the (4+1) addition. NICS calculations show that this ring has less antiaromatic character in **4** (NICS(0) = 5.3) than in biphenylene (NICS(0) = 18.3) (Scheme 1). The (4+1) cycloaddition of **1** to the C³ and C⁶ positions of biphenylene was also considered computationally. Both the transition state and product were found to be higher in energy than the pathway toward the formation of **4** (Supporting Information, Figure S5).

Intermediate **4** has two chemically distinct alkene systems. **1** has previously been shown to perform (2+1) cycloadditions with alkenes.²⁰ Addition of **1** to the more sterically hindered alkene (C²–C³) of **4** was found to proceed *via* a low energy transition state **TS-2** to form **Int-1**.^{18,20} **Int-1** is the product of an overall 4e⁻ reduction of the arene ring of biphenylene with two equiv. of **1**. **Int-1** incorporates two aluminium centres at the bridgehead positions of a norbornane (Al^A) and norcarane (Al^B) skeleton on opposite faces of the reduced arene ring.

Subsequent isomerisation of **Int-1** to **Int-2** occurs *via* transition state **TS-3**. This reaction involves a concerted rearrangement of the ring framework of **Int-1** with two four-membered metallocycles forming from one three-membered and one five-membered metallocycle. The relief of strain from the

ring-expansion of the aluminocyclopropane is the likely thermodynamic driving force of this reaction (Figure 3a). From **Int-2** C–C σ -bond activation can occur by two different pathways. These pathways both involve a [1,3]-sigmatropic shift and a step that breaks the C–C σ -bond, however the order of these processes differs between the two.

C²–C³ σ -bond activation is possible directly from **Int-2** *via* **TS-4** to form **Int-3**. **TS-4** is an early transition state in which a key bond cleavage step triggers a series of bond migrations to form **Int-3** (Figure 3b). A subsequent [1,3]-sigmatropic shift of [Al^A] along the C³–C⁵ allyl system *via* **TS-5** results in formation of the observed product **2**.

Alternatively, [Al^A] can perform a [1,3]-sigmatropic shift along the C²–C⁶ allyl system in **Int-2** *via* **TS-6** to form **Int-4**. The aluminium centres in **Int-4** are related by C₂-rotation around the perpendicular bisector of the C⁴–C⁵ bond and this symmetry is preserved into the product. Activation of the C⁴–C⁵ σ -bond from **Int-4** occurs *via* **TS-7** and results in formation of **3**.

The formation of both **2** and **3** is calculated to be highly exergonic and is consistent with these reactions being non-reversible. The C–C σ -bond activation steps have the highest activation barriers in the formation of **2** ($\Delta G^\ddagger_{298K} = 29.4$ kcal mol⁻¹) and **3** ($\Delta G^\ddagger_{298K} = 29.6$ kcal mol⁻¹) from **1** and biphenylene. In both cases, C–C σ -bond activation originates from a highly strained metallocyclic intermediate in which aluminium

centers occupy positions in two separate metallocyclobutane rings. C–C cleavage *via* **TS-4** and **TS-7** relieves the strained 4-membered rings in **Int-2** and **Int-4** respectively. The computationally predicted selectivity for the formation of **2** in preference to **3** ($\Delta\Delta G^\ddagger_{298\text{K}} = 0.2 \text{ kcal mol}^{-1}$) is consistent with experiment. The small ($\Delta\Delta G^\ddagger \sim 1 \text{ kcal mol}^{-1}$) difference in activation energies between C–C σ -bond activation steps (**TS-4** and **TS-7**) and the (4+1) cycloaddition step (**TS-1**) is reflected experimentally: formation of **2** and **3** is observed during the synthesis of **4**.

The oxidative addition of **1** to the central C¹–C⁷ σ -bond of biphenylene was also explored computationally. While the overall reaction is highly exergonic ($\Delta G_{298\text{K}} = -64.9 \text{ kcal mol}^{-1}$), the transition state is high in energy (**TS-8**, $\Delta G^\ddagger_{298\text{K}} = 42.0 \text{ kcal mol}^{-1}$) and unlikely to be accessible under the reaction conditions (Supporting Information, Figure S4).

A deeper consideration of the key transition states can be used to rationalise the remarkable chemoselectivity of this system. Transition states for oxidative addition are typically destabilised by steric repulsion due to the requirement for a close – and side-on – approach of the C–X bond to the metal to achieve orbital overlap (e.g. X = C, H, Halogen). The oxidative addition of biphenylene to transition metal complexes has been calculated to involve a side-on approach and the synchronous breaking of the C–C σ -bond and formation of both M–C bonds along the reaction coordinate.^{21,22}

1 differs from common transition metal fragments in terms of both the symmetry and energy of the frontier molecular orbitals. The HOMO and the LUMO of **1** are orthogonal to one another and the 3p orbitals which contribute to them have a smaller radial extension than the 4d and 5d orbitals of 2nd and 3rd row transition metals. In line with these differences, **TS-8** is unusual. **TS-8** has an asymmetric geometry and involves close approach of the metal to only one carbon atom of the breaking C–C σ -bond (Figure 4b). Activation strain analysis (ASA) was used to interrogate this phenomenon further.²³ The ASA profile derived from **TS-8** was generated (Figure 4b). In this profile, E_{int} decreases modestly along the reaction coordinate and a high degree of distortion in the biphenylene fragment is required before favourable interaction between the two fragments is observed. This is further supported by the complex curvature of the reaction coordinate as a function of the Al–C bond distance. For comparison, the profile associated with **TS-1**, and the (4+1) cycloaddition, shows a steady increase in $E_{\text{dist}}(\text{Biph})$ that is offset by a steady decrease in E_{int} (Figure 4a). The data allow us to conclude that the oxidative addition of biphenylene to **1** is unfavourable due to the high degree of biphenylene reorganisation that is required for effective orbital overlap.




Figure 3: Activation strain analysis of the reaction profiles derived from **TS-8** (a) and **TS-1** (b).

The divergent reactivity of **1** from transition-metal systems is a direct consequence of the orthogonality and limited radial extension of the frontier molecular orbitals on aluminium and the associated strain required to reach an oxidative addition transition state.

ASSOCIATED CONTENT

The Supporting Information is available free of charge on the ACS Publications website. X-ray crystallographic data **2-4** are available from the Cambridge Crystallographic Data Centre (CCDC 2023760-2023762) and as a .cif file, full details of the experiments and calculations are available as a .pdf. NMR spectra and computational coordinates are available at dx.doi.org/10.14469/hpc/7325.

AUTHOR INFORMATION

Corresponding Author

Mark R. Crimmin – Department of Chemistry, Molecular Sciences Research Hub, Imperial College London, London W12 0BZ, U.K

Email: m.crimmin@imperial.ac.uk

Author

Richard Y. Kong – Department of Chemistry, Molecular Sciences Research Hub, Imperial College London, London W12 0BZ, U.K

Notes


The authors declare no competing financial interest.

ACKNOWLEDGMENT

We are grateful to the ERC (FluoroFix: 677367) for generous funding. We thank Imperial College London for the award of a President's Scholarship (RYK).

REFERENCES

- (1) Murakami, M.; Ishida, N. Potential of Metal-Catalyzed C–C Single Bond Cleavage for Organic Synthesis. *J. Am. Chem. Soc.* **2016**, *138*, 13759–13769.
- (2) Sarpong, R.; Wang, B.; Perea, M. A. Transition Metal-Mediated C–C Single Bond Cleavage: Making the Cut in Total Synthesis. *Angew. Chem., Int. Ed.* <http://dx.doi.org/10.1002/anie.201915657>.
- (3) Jones, W. D. Mechanistic Studies of Transition Metal-Mediated C–C Bond Activation. In *C–C Bond Activation*; Dong, G., Ed.; Topics in Current Chemistry; Springer Berlin Heidelberg: Berlin, Heidelberg, 2013; Vol. 346, pp 1–31.
- (4) Eisch, J. J.; Piotrowski, A. M.; Han, K. I.; Kruger, Carl.; Tsay, Y. H. Organic Chemistry of Subvalent Transition Complexes. 9. Oxidative Addition of Nickel(0) Complexes to Carbon-Carbon Bonds in Biphenylene: Formation of Nickelole and 1,2-Dinickelcyclopropanes. *Organometallics* **1985**, *4*, 224–231.
- (5) Perthuisot, C.; Jones, W. D. Catalytic Hydrogenolysis of an Aryl-Aryl Carbon-Carbon Bond with a Rhodium Complex. *J. Am. Chem. Soc.* **1994**, *116*, 3647–3648.
- (6) Perthuisot, C.; Edelbach, B. L.; Zubris, D. L.; Jones, W. D. C–C Activation in Biphenylene. Synthesis, Structure, and Reactivity of $(C_5Me_5)_2M_2(2,2'-Biphenyl) ($M = Rh, Co$). *Organometallics* **1997**, *16*, 2016–2023.$
- (7) Takano, H.; Ito, T.; Kanyiva, K. S.; Shibata, T. Recent Advances of Biphenylene: Synthesis, Reactions and Uses. *Eur. J. Org. Chem.* **2019**, *2019*, 2871–2883.
- (8) Koshino, K.; Kinjo, R. Construction of σ -Aromatic AlB2 Ring via Borane Coupling with a Dicoordinate Cyclic (Alkyl)(Amino)Aluminy Anion. *J. Am. Chem. Soc.* **2020**, *142*, 9057–9062.
- (9) Browning, J.; Green, M.; Laguna, A.; Smart, L. E.; Spencer, J. L.; Stone, F. G. A. A Platinum-Promoted Ring-Opening Reaction of Hexakis(trifluoromethyl)benzene Involving Carbon–Carbon Bond Cleavage; Crystal and Molecular Structure of Bis(*t*-Butyl Isocyanide)Platinacyclohepta-Cis-2,Cis-4,Cis-6-Triene. *J. Chem. Soc., Chem. Commun.* **1975**, No. 17, 723–724.
- (10) Ellis, D.; McKay, D.; Macgregor, S. A.; Rosair, G. M.; Welch, A. J. Room-Temperature C–C Bond Cleavage of an Arene by a Metallocarborane. *Angew. Chem., Int. Ed.* **2010**, *49*, 4943–4945.
- (11) Sattler, A.; Parkin, G. Cleaving Carbon–Carbon Bonds by Inserting Tungsten into Unstrained Aromatic Rings. *Nature* **2010**, *463*, 523–526.
- (12) Zeits, P. D.; Fiedler, T.; Gladysz, J. A. Ring Opening Metathesis Polymerization of an H_4 -Benzene Complex: A Direct Synthesis of a Polyacetylene with a Regular Pattern of π Bound Metal Fragments. *Chem. Commun.* **2012**, *48*, 7925–7927.
- (13) Hu, S.; Shima, T.; Hou, Z. Carbon–Carbon Bond Cleavage and Rearrangement of Benzene by a Trinuclear Titanium Hydride. *Nature* **2014**, *512*, 413–415.
- (14) Meola, G.; Braband, H.; Hernández-Valdés, D.; Gotzmann, C.; Fox, T.; Spingler, B.; Alberto, R. A Mixed-Ring Sandwich Complex from Unexpected Ring Contraction in $[Re(H_6-C_6H_5Br)(H_6-C_6R_6)](PF_6)$. *Inorg. Chem.* **2017**, *56*, 6297–6301.
- (15) Jakoobi, M.; Halcovitch, N.; Whitehead, G. F. S.; Sergeev, A. G. Selective Arene Cleavage by Direct Insertion of Iridium into the Aromatic Ring. *Angew. Chem., Int. Ed.* **2017**, *56*, 3266–3269.
- (16) Jakoobi, M.; Tian, Y.; Boulatov, R.; Sergeev, A. G. Reversible Insertion of Ir into Arene Ring C–C Bonds with Improved Regioselectivity at a Higher Reaction Temperature. *J. Am. Chem. Soc.* **2019**, *141*, 6048–6053.
- (17) Hicks, J.; Vasko, P.; Goicoechea, J. M.; Aldridge, S. Reversible, Room-Temperature C–C Bond Activation of Benzene by an Isolable Metal Complex. *J. Am. Chem. Soc.* **2019**, *141*, 11000–11003.
- (18) Kong, R. Y.; Crimmin, M. R. Activation and Functionalization of C–C σ Bonds of Alkylidene Cyclopropanes at Main Group Centers. *J. Am. Chem. Soc.* **2020**, *142*, 11967–11971.
- (19) Bakewell, C.; Garçon, M.; Kong, R. Y.; O'Hare, L.; White, A. J. P.; Crimmin, M. R. Reactions of an Aluminum(I) Reagent with 1,2-, 1,3-, and 1,5-Dienes: Dearomatization, Reversibility, and a Pericyclic Mechanism. *Inorg. Chem.* **2020**, *59*, 4608–4616.
- (20) Bakewell, C.; White, A. J. P.; Crimmin, M. R. Reversible Alkene Binding and Allylic C–H Activation with an Aluminium(I) Complex. *Chem. Sci.* **2019**, *10*, 2452–2458.
- (21) Gu, Z.; Boursalian, G. B.; Gandon, V.; Padilla, R.; Shen, H.; Timofeeva, T. V.; Tongwa, P.; Vollhardt, K. P. C.; Yakovenko, A. A. Activated Phenacenes from Phenyls by Nickel-Catalyzed Alkyne Cycloadditions. *Angew. Chem., Int. Ed.* **2011**, *50*, 9413–9417.
- (22) Takano, H.; Sugimura, N.; Kanyiva, K. S.; Shibata, T. DFT Studies on the Mechanism of the Iridium-Catalyzed Formal [4 + 1] Cycloaddition of Biphenylene with Alkenes. *ACS Omega* **2017**, *2*, 5228–5234.
- (23) This approach balances the energy penalty associated with distortion of the reagent fragments, (E_{dist}), with the energy gain from their interaction (E_{int}) along the reaction coordinate. As two bonds are forming over the course of both reactions, these values have been plotted against the lengths of the weaker Al–C bond at the transition state (See Supporting Information for further discussion).



Insert Table of Contents artwork here