

Pulse Dipolar EPR Reveals Double-Histidine Motif

Spin-labelling is Robust Against Competitor Ions

Joshua L. Wort,^{a,b} Swati Arya,^{b,c} Katrin Ackermann,^{a,b} Alan J. Stewart,^{b,c} Bela E. Bode^{a,b}*

^aEaStCHEM School of Chemistry, ^bBiomedical Sciences Research Complex, Centre of Magnetic Resonance, University of St Andrews, North Haugh, St Andrews, KY16 9ST, UK. ^cSchool of Medicine, University of St Andrews, North Haugh, St Andrews, KY16 9TF, UK

Corresponding Author

*beb2@st-andrews.ac.uk

ABSTRACT. Pulse-dipolar EPR is an appealing strategy for structural characterization of complex systems in solution that complements other biophysical techniques. Significantly, the emergence of genetically encoded self-assembling spin labels exploiting exogenously introduced double-histidine motifs in conjunction with Cu^{II}-chelates offers high precision distance determination in systems non-permissive to thiol-directed spin labelling. However, the non-covalency of this interaction exposes potential vulnerabilities to competition from adventitious divalent metal ions, and pH sensitivity. Herein, a combination of room-temperature isothermal titration calorimetry (ITC) and cryogenic relaxation-induced dipolar modulation enhancement (RIDME) measurements are applied to the model protein *Streptococcus sp.* group G. protein G, B1 domain (GB1). Results demonstrate double-histidine motif spin labelling using Cu^{II}-nitrilotriacetic acid (Cu^{II}-NTA) is robust against the competitor ligand Zn^{II}-NTA at >1000-fold molar excess, and high nM binding affinity is surprisingly retained under acidic and basic conditions even though room temperature affinity shows a stronger pH dependence. This indicates the strategy is well-suited for diverse biological applications, particularly metalloproteins with divalent metal ion cofactors.

KEYWORDS RIDME; Competitive Binding; GB1; EPR; Dissociation Constant; ITC

As the complexity of biomolecular assemblies implicated in health and disease has increased, so too has interest in pulse-dipolar EPR (PD-EPR) as a robust strategy for solution-state structural characterization of proteins^{1,2} and nucleic acids^{3,4} in the nanometer distance regime.^{5,6} PD-EPR is a powerful tool that complements X-ray crystallography, NMR, cryo-EM and Förster Resonance Energy Transfer (FRET) data by providing structural insight outwith crystallization, size-limitation, or structurally perturbative labels. Hence, PD-EPR has been applied to study conformational equilibria,⁷ oligomerization degree,^{8,9} complexation events,¹⁰⁻¹² and competing structural models.¹³

Pairs of paramagnetic moieties are commonly introduced into diamagnetic systems of interest using thiol-based site-directed spin labelling.¹⁴ Cysteine residues are typically covalently modified, as for the nitroxide R1 sidechain (figure 1a top). This strategy is suboptimal in systems containing essential cysteine residues, non-permissive to post-translational reduction. However, Cu^{II}-based genetically-encodable self-assembling spin labels using double-histidine motifs have emerged as an alternative labelling strategy.^{15,16} Additionally, the bipedal mode of Cu^{II}-chelate attachment at the double-histidine motif (figure 1a bottom) results in significantly improved precision and accuracy in the distance domain. Cu^{II}-nitrilotriacetic acid (Cu^{II}-NTA) spin labelling of double histidine motifs for PD-EPR has been applied successfully to enzymes,¹⁷ metalloproteins,¹⁸ and nucleoprotein complexes.¹⁹




Figure 1. Spin label structures and the GB1 constructs used in this work. a) MTSL nitroxide conjugated to a cysteine residue, resulting in the R1 sidechain (top) and Cu^{II} -NTA coordinated to a double-histidine motif (bottom). b) Cartoon representation of the I6R1/K28H/Q32H GB1 construct, with the R1 nitroxide and Cu^{II} -NTA spin labels shown in stick representation. c) Cartoon representation of the K28H/Q32H GB1 construct, with the Cu^{II} -NTA shown in stick representation.

Despite this success, optimization of the spin labelling approach is non-trivial, because the non-covalency of the interaction predisposes sensitivity to variations in binding affinity. For instance, different buffer conditions influence the double-histidine motif labelling efficiency with Cu^{II} -NTA.²⁰ Furthermore, while the influence of pH upon formation of Cu^{II} -chelates has been characterized by CW-EPR previously,²¹ current literature has not shown how pH variations influence binding at the double-histidine motif, particularly under cryogenic temperatures. Similarly, current literature has not addressed competition for double-histidine motif sites by adventitious divalent metal ions, and so warrants investigation. In the current study, *Streptococcus sp.* Group G. protein G, B1 domain (GB1) constructs (I6R1/K28H/Q32H (figure 1b) and K28H/Q32H (figure 1c) were used as biological model systems, in Cu^{II} -nitroxide relaxation-induced dipolar modulation enhancement (RIDME)²² pseudo-titrations,^{10,23} and isothermal titration calorimetry (ITC) measurements, respectively.

Measurements were first performed in presence of the model competitor ligand, Zn^{II}-NTA, which was chosen because i) it is a weak ligand for double-histidine motifs compared to Cu^{II}-NTA, and ii) it is diamagnetic, so does not contribute to the detected EPR signal. An EPR silent competitor ligand is desirable because analysis of pseudo-titration data is simplified (see SI section 1.6). Room temperature ITC data (figure 2a) fitted to a one-site model where binding stoichiometry could vary, indicated a binding affinity of 513 μ M. Using the determined enthalpy change (ΔH), the binding affinity was extrapolated to 235 K (i.e., the temperature at which the binding equilibrium is found to freeze out in our samples, such that diffusional processes cease, meaning our EPR data reflects equilibria at 235 K),²² to determine the influence of the competitor ligand upon double-histidine loading efficiency with Cu^{II}-NTA under PD-EPR conditions.




Figure 2. Zn^{II} -NTA competitor RIDME pseudo-titration. a) ITC data performed at 298 K, 800 μM 28H/32H GB1 titrated against 12 mM Zn^{II} -NTA. b) RIDME dipolar evolution functions, with the corresponding fits shown in dotted black. Modulation depths (Δ) are indicated. c) Validated RIDME distance distributions, corresponding to the dipolar evolution functions shown in b). The colour scheme is the same in b) and c). The concentrations of Zn^{II} -NTA are indicated. Colour bars represent reliability ranges (green: shape reliable; yellow: mean and width reliable; orange: mean reliable; red: no quantification possible). d) A univariate fit of the competitor dissociation constant (32 μM) is shown in solid black. Experimental points are shown as the blue scatter, and 95% confidence intervals are shown as the red error bars.

The corresponding RIDME pseudo-titration was performed at 1 μM protein concentration in presence of 10 μM Cu^{II} -NTA (to ensure quantitative loading and negligible ligand depletion²⁴ (see SI section 1.6)) and varying Zn^{II} -NTA concentrations. Importantly, the dipolar evolution functions (figure 2b) and distance distributions (figure 2c) show that in all cases, the expected

peak at ~ 2.5 nm is retrieved as the only significant feature following data validation. The fitted competitor K_D value ($32 \mu\text{M}$) is within 2-fold of that determined from ITC when extrapolated to 235 K ($48 \mu\text{M}$) (figure 2d). This suggests that Cu^{II} -NTA is robust against adventitious divalent metals in vast excesses, >1000 -fold, even at low μM protein concentrations. Additionally, this benchmarks quantitation of Cu^{II} -nitroxide RIDME modulation depths for remotely determining binding affinities of EPR silent ligands, in a competition assay format.

Next, the influence of pH upon double-histidine motif loading efficiency with Cu^{II} -NTA was investigated by measuring ITC and RIDME at pH 6.4. Since only deprotonated histidine residues can coordinate Cu^{II} -NTA, it follows that binding affinity should decrease under acidic conditions. Indeed, room-temperature ITC performed at pH 5, below the approximate pK_A of solvent-exposed histidine,²⁵ shows negligible binding (see SI section 2.3), and measurements at pH 6.4, fitted to a one-site model, indicated a 20-fold reduction in affinity compared to previous work²³ (figure 3a). Extrapolating ΔH to 235 K suggested a binding affinity of $\sim 4 \mu\text{M}$.




Figure 3. pH 6.4 RIDME pseudo-titration. a) ITC data performed at 298 K, 75 μM K28H/Q32H GB1 titrated against 2 mM Cu^{II} -NTA. b) RIDME dipolar evolution functions, with the corresponding fits shown in dotted black. Modulation depths (Δ) are indicated. c) Validated RIDME distance distributions, corresponding to the dipolar evolution functions shown in b). The colour scheme is the same in b) and c). The concentrations of Cu^{II} -NTA are indicated. d) A bivariate fit of the dissociation constant (0.31 μM) is shown in solid black. Experimental points are shown as the blue scatter, and 95% confidence intervals are shown as the red error bars.

A RIDME pseudo-titration was performed at 5 μM protein concentration to validate the room-temperature ITC prediction of reduced affinity under PD-EPR conditions. Significantly, the dipolar evolution functions (figure 3b) show Cu^{II} -NTA binding is only marginally reduced at lower pH, with one equivalent Cu^{II} -NTA saturating $\sim 70\%$ of available double-histidine motifs. This is further borne out by the fitted dissociation constant (figure 3d), 0.31 μM compared to 0.14 μM in previous work at pH 7.4.²³ The affinity reduced by only 2-fold, indicating that the

influence of pH upon double-histidine motif loading may be attenuated at lower temperatures. A possible explanation is that histidine protonation is endothermic,²⁶ driving the equilibrium towards the deprotonated state at lower temperatures, compensating for reduced pH and facilitating double-histidine loading. Importantly, this would also imply significantly tighter binding at higher pH, where histidine deprotonation is already favored.




Figure 4. pH 8.4 RIDME pseudo-titration. a) ITC data performed at 298 K, 75 μM K28H/Q32H GB1 titrated against 2.5 mM Cu^{II} -NTA. b) RIDME dipolar evolution functions, with the corresponding fits shown in dotted black. Modulation depths (Δ) are indicated. c) Validated RIDME distance distributions, corresponding to the dipolar evolution functions shown in b). The colour scheme is the same in b) and c). The concentrations of Cu^{II} -NTA are indicated. d) A bivariate fit of the dissociation constant (0.091 μM) shown in solid black. Experimental points are shown as the blue scatter, and 95% confidence intervals are shown as the red error bars.

To clarify the disparity between ITC and PD-EPR data at pH 6.4, room-temperature ITC was also performed at pH 8.4 (figure 4a), fitted to a one-site model, where a 20-fold increase in affinity was predicted (*via* improved thermodynamic favorability of binding) compared to previous work. Another RIDME pseudo-titration was performed at 2 μM protein concentration, with dipolar evolution functions (figure 4b) suggesting modest improvement in binding affinity. The fitted dissociation constant (figure 4d) of 0.091 μM indicates binding affinity is approximately 2-fold higher than at pH 7.4, consistent with observation at pH 6.4 that the influence of pH upon binding affinity is attenuated with decreasing temperature. While an endothermic protonation process would suggest much tighter binding is to be anticipated at pH 8.4, consider that at this pH <1% of histidine δ -nitrogen atoms should remain protonated. This may explain why the relative increase in binding affinity is smaller than expected, since the deprotonation is already driven toward completion by the high pH.

While the data suggests that spin-labelling and measurement at pH 8.4 will afford enhanced loading and sensitivity, it should be noted that the stoichiometry of binding is ~ 2 , compared to ~ 1 at pH 6.4. This may arise from deprotonation of the protein surface that promotes non-specific binding. This would explain the increased exothermic nature of the binding, if non-specific or additional binding events contributed to the isotherm and would further inflate the binding affinity when extrapolated to cryogenic temperatures. However, the corresponding distance distributions (figure 4c) do not contain additional peaks to support this hypothesis.

Perhaps most significantly, these results clearly show that Cu^{II} -NTA binding affinity for double-histidine motifs is not strongly perturbed from the high nM concentration regime by fluctuations of pH between 6.4-8.4. Coupled with measurements in presence of competitor ligand Zn^{II} -NTA, findings support that Cu^{II} -NTA is a highly robust spin label when combined with α -helical

double-histidine motifs. This is encouraging for the widespread application of double-histidine motifs in metalloproteins, or in systems where divalent metal cofactors are necessary.

Additionally, the benchmarking of a competition assay using PD-EPR is particularly exciting because it allows remote detection of binding interactions with diamagnetic ligands. This will be promising in cases where paramagnetic ligand analogues are not available or cause structural perturbation. PD-EPR also has greater sensitivity than ITC and the coupling of thermodynamic and structural information allows for the facile monitoring of non-specific and competitor ligand interactions.²⁷ Traditionally, monitoring competitive ligand binding has required expensive radio-labelling and judicious selection of appropriate isotopes.^{24,28} PD-EPR may complement these strategies, while obviating potential cost and safety considerations.

ASSOCIATED CONTENT

Supporting Information.

The following files are available free of charge.

I Experimental Procedures: Construct Design, Expression and Purification; Pulse EPR Sample Preparation; Metal Chelate Spin Label Preparation; Mass Spectrometry; Pulse EPR Measurements; Competitive Binding Model; Isothermal Titration Calorimetry; UV-visible Spectroscopy. II Results and Discussion: Inversion Recovery Measurements; 5-pulse RIDME Measurements; Influence of Differential pH upon Double-Histidine Motif Affinity; Influence of Differential pH upon Cu^{II}-NTA Complex Formation; Optimization of Cu^{II}-IDA Complex Formation. III References (PDF)

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

The authors thank Dr Hassane El Mkami for help with EPR. JLW is supported by the BBSRC DTP Eastbio. We thank the Leverhulme Trust for support (RPG-2018-397). This work was supported by equipment funding through the Wellcome Trust (099149/Z/12/Z) and BBSRC (BB/R013780/1). We gratefully acknowledge ISSF support to the University of St Andrews from the Wellcome Trust.

REFERENCES

- (1) Edwards, D. T.; Huber, T.; Hussain, S.; Stone, K. M.; Kinnebrew, M.; Kaminker, I.; Matalon, E.; Sherwin, M. S.; Goldfarb, D.; Han, S. Determining the Oligomeric Structure of Proteorhodopsin by Gd³⁺-Based Pulsed Dipolar Spectroscopy of Multiple Distances. *Structure* **2014**, 22 (11), 1677–1686. <https://doi.org/10.1016/j.str.2014.09.008>.
- (2) Verhalen, B.; Dastvan, R.; Thangapandian, S.; Peskova, Y.; Koteiche, H. A.; Nakamoto, R. K.; Tajkhorshid, E.; McHaourab, H. S. Energy Transduction and Alternating Access of the Mammalian ABC Transporter P-Glycoprotein. *Nature* **2017**. <https://doi.org/10.1038/nature21414>.
- (3) Schiemann, O.; Piton, N.; Mu, Y.; Stock, G.; Engels, J. W.; Prisner, T. F. A PELDOR-Based Nanometer Distance Ruler for Oligonucleotides. *J. Am. Chem. Soc.* **2004**. <https://doi.org/10.1021/ja0393877>.
- (4) Babaylova, E. S.; Malygin, A. A.; Lomzov, A. A.; Pyshnyi, D. V.; Yulikov, M.; Jeschke, G.; Krumkacheva, O. A.; Fedin, M. V.; Karpova, G. G.; Bagryanskaya, E. G. Complementary-Addressed Site-Directed Spin Labeling of Long Natural RNAs. *Nucleic*

- Acids Res.* **2016**. <https://doi.org/10.1093/nar/gkw516>.
- (5) Jeschke, G. The Contribution of Modern EPR to Structural Biology. *Emerg. Top. Life Sci.* **2018**. <https://doi.org/10.1042/etls20170143>.
- (6) Sanabria, H.; Rodnin, D.; Hemmen, K.; Peulen, T. O.; Felekyan, S.; Fleissner, M. R.; Dimura, M.; Koberling, F.; Kühnemuth, R.; Hubbell, W.; Gohlke, H.; Seidel, C. A. M. Resolving Dynamics and Function of Transient States in Single Enzyme Molecules. *Nat. Commun.* **2020**. <https://doi.org/10.1038/s41467-020-14886-w>.
- (7) Joseph, B.; Sikora, A.; Cafiso, D. S. Ligand Induced Conformational Changes of a Membrane Transporter in E. Coli Cells Observed with DEER/PELDOR. *J. Am. Chem. Soc.* **2016**. <https://doi.org/10.1021/jacs.5b13382>.
- (8) Hagelueken, G.; Ingledeu, W. J.; Huang, H.; Petrovic-Stojanovska, B.; Whitfield, C.; ELMkami, H.; Schiemann, O.; Naismith, J. H. PELDOR Spectroscopy Distance Fingerprinting of the Octameric Outer-Membrane Protein Wza from Escherichia Coli. *Angew. Chemie - Int. Ed.* **2009**. <https://doi.org/10.1002/anie.200805758>.
- (9) Bode, B. E.; Margraf, D.; Plackmeyer, J.; Dürner, G.; Prisner, T. F.; Schiemann, O. Counting the Monomers in Nanometer-Sized Oligomers by Pulsed Electron-Electron Double Resonance. *J. Am. Chem. Soc.* **2007**. <https://doi.org/10.1021/ja065787t>.
- (10) Giannoulis, A.; Oranges, M.; Bode, B. E. Monitoring Complex Formation by Relaxation-Induced Pulse Electron Paramagnetic Resonance Distance Measurements. *ChemPhysChem* **2017**, *18* (17), 2318–2321. <https://doi.org/10.1002/cphc.201700666>.

- (11) Ackermann, K.; Giannoulis, A.; Cordes, D. B.; Slawin, A. M. Z.; Bode, B. E. Assessing Dimerisation Degree and Cooperativity in a Biomimetic Small-Molecule Model by Pulsed EPR. *Chem. Commun.* **2015**. <https://doi.org/10.1039/c4cc08656b>.
- (12) Giannoulis, A.; Ackermann, K.; Spindler, P. E.; Higgins, C.; Cordes, D. B.; Slawin, A. M. Z.; Prisner, T. F.; Bode, B. E. Nitroxide-Nitroxide and Nitroxide-Metal Distance Measurements in Transition Metal Complexes with Two or Three Paramagnetic Centres Give Access to Thermodynamic and Kinetic Stabilities. *Phys. Chem. Chem. Phys.* **2018**. <https://doi.org/10.1039/c8cp01611a>.
- (13) Theillet, F. X.; Binolfi, A.; Bekei, B.; Martorana, A.; Rose, H. M.; Stuiiver, M.; Verzini, S.; Lorenz, D.; Van Rossum, M.; Goldfarb, D.; Selenko, P. Structural Disorder of Monomeric α -Synuclein Persists in Mammalian Cells. *Nature* **2016**. <https://doi.org/10.1038/nature16531>.
- (14) Hubbell, W. L.; Altenbach, C. Investigation of Structure and Dynamics in Membrane Proteins Using Site-Directed Spin Labeling. *Curr. Opin. Struct. Biol.* **1994**. [https://doi.org/10.1016/S0959-440X\(94\)90219-4](https://doi.org/10.1016/S0959-440X(94)90219-4).
- (15) Cunningham, T. F.; Putterman, M. R.; Desai, A.; Horne, W. S.; Saxena, S. The Double-Histidine Cu²⁺-Binding Motif: A Highly Rigid, Site-Specific Spin Probe for Electron Spin Resonance Distance Measurements. *Angew. Chemie - Int. Ed.* **2015**. <https://doi.org/10.1002/anie.201501968>.
- (16) Ghosh, S.; Lawless, M. J.; Rule, G. S.; Saxena, S. The Cu²⁺-Nitrilotriacetic Acid Complex Improves Loading of α -Helical Double Histidine Site for Precise Distance

- Measurements by Pulsed ESR. *J. Magn. Reson.* **2018**.
<https://doi.org/10.1016/j.jmr.2017.12.005>.
- (17) Lawless, M. J.; Pettersson, J. R.; Rule, G. S.; Lanni, F.; Saxena, S. ESR Resolves the C Terminus Structure of the Ligand-Free Human Glutathione S-Transferase A1-1. *Biophys. J.* **2018**, *114* (3), 592–601. <https://doi.org/10.1016/j.bpj.2017.12.016>.
- (18) Gamble Jarvi, A.; Cunningham, T. F.; Saxena, S. Efficient Localization of a Native Metal Ion within a Protein by Cu²⁺-Based EPR Distance Measurements. *Phys. Chem. Chem. Phys.* **2019**. <https://doi.org/10.1039/c8cp07143h>.
- (19) Sameach, H.; Ghosh, S.; Gevorkyan-Airapetov, L.; Saxena, S.; Ruthstein, S. EPR Spectroscopy Detects Various Active State Conformations of the Transcriptional Regulator CueR. *Angew. Chemie - Int. Ed.* **2019**. <https://doi.org/10.1002/anie.201810656>.
- (20) Gamble Jarvi, A.; Casto, J.; Saxena, S. Buffer Effects on Site Directed Cu²⁺-Labeling Using the Double Histidine Motif. *J. Magn. Reson.* **2020**.
<https://doi.org/10.1016/j.jmr.2020.106848>.
- (21) Lawless, M. J.; Ghosh, S.; Cunningham, T. F.; Shimshi, A.; Saxena, S. On the Use of the Cu²⁺-Iminodiacetic Acid Complex for Double Histidine Based Distance Measurements by Pulsed ESR. *Phys. Chem. Chem. Phys.* **2017**. <https://doi.org/10.1039/c7cp02564e>.
- (22) Milikisyants, S.; Scarpelli, F.; Finiguerra, M. G.; Ubbink, M.; Huber, M. A Pulsed EPR Method to Determine Distances between Paramagnetic Centers with Strong Spectral Anisotropy and Radicals: The Dead-Time Free RIDME Sequence. *J. Magn. Reson.* **2009**, *201* (1), 48–56. <https://doi.org/10.1016/j.jmr.2009.08.008>.

- (23) Wort, J. L.; Ackermann, K.; Giannoulis, A.; Stewart, A. J.; Norman, D. G.; Bode, B. E. Sub-Micromolar Pulse Dipolar EPR Spectroscopy Reveals Increasing CuII-Labeling of Double-Histidine Motifs with Lower Temperature. *Angew. Chemie - Int. Ed.* **2019**. <https://doi.org/10.1002/anie.201904848>.
- (24) Hulme, E. C.; Trevethick, M. A. Ligand Binding Assays at Equilibrium: Validation and Interpretation. *British Journal of Pharmacology*. 2010. <https://doi.org/10.1111/j.1476-5381.2009.00604.x>.
- (25) Cohen, J. S.; Griffin, J. H.; Schechter, A. N. Nuclear Magnetic Resonance Titration Curves of Histidine Ring Protons. IV. The Effects of Phosphate and Sulfate on Ribonuclease. *J. Biol. Chem.* **1973**.
- (26) Tanokura, M.; Tasumi, M.; Miyazawa, T. NMR STUDY ON THE PROTONATION OF IMIDAZOLE RING OF N -ACETYL-L-HISTIDINE METHYLAMIDE, A MODEL FOR HISTIDINE RESIDUES EXPOSED TO AQUEOUS SOLVENT . *Chem. Lett.* **1978**. <https://doi.org/10.1246/cl.1978.739>.
- (27) Bankston, J. R.; DeBerg, H. A.; Stoll, S.; Zagotta, W. N. Mechanism for the Inhibition of the CAMP Dependence of HCN Ion Channels by the Auxiliary Subunit TRIP8b. *J. Biol. Chem.* **2017**. <https://doi.org/10.1074/jbc.M117.800722>.
- (28) Maguire, J. J.; Kuc, R. E.; Davenport, A. P. Radioligand Binding Assays and Their Analysis. *Methods Mol. Biol.* **2012**. https://doi.org/10.1007/978-1-61779-909-9_3.