

Synthesis of 1-(1-Arylvinyl)pyridin-2(1H)-ones from Ketones and 2-Fluoropyridine

Takuji Kawamoto,* Shunya Ikeda, and Akio Kamimura

Department of Applied Chemistry, Yamaguchi University, Ube, Yamaguchi 755-8611, Japan

ABSTRACT: Pyridone skeletons are found in numerous biologically active molecules and pharmaceuticals. 1-(1-Arylvinyl)pyridin-2(1H)-ones are synthetic intermediates derived from the enamide moiety, and only few examples of the synthesis of 1-(1-arylvinyl)-2-pyridones have been reported. In this work, a simple and efficient procedure for the synthesis of N-vinyl-substituted pyridones from ketones and 2-fluoropyridine in the presence of trifluoromethane sulfonic anhydride, followed by base treatment is described. Various ketones with electron-donating or -withdrawing groups at the benzene rings can be used in this reaction. A preliminary mechanistic study indicates that it is not very likely that both vinyl triflates and vinyl cations play major roles as intermediates in this transformation. The thus obtained pyridones can be subsequently transformed via C–H arylation and radical alkylation reactions.


Pyridone, i.e., pyridin-2(1H)-one, skeletons exist in numerous biologically active molecules and pharmaceuticals.^{1–7} The development of efficient synthetic methods for the straightforward construction of this structural motif has thus received considerable interest from the synthetic community, and several synthetic routes to N-substituted pyridones have been reported.^{8,9} Although 1-(1-arylvinyl)pyridin-2(1H)-ones are useful synthetic intermediates derived from the enamide moiety, only few examples of the synthesis of 1-(1-arylvinyl)-2-pyridones have been reported.¹⁰ Undheim and co-workers have reported the reaction of 1-bromophenylacetylene with the sodium salt of 2-pyridinone (eq. 1).¹¹ Rao *et al.* have reported the nucleophilic addition to cyano-substituted aryl alkynes leading to 1-(1-arylvinyl)-2-pyridones.¹²

On the other hand, 2-halopyridinium salts are easily converted into 2-pyridones.^{13–18} Due to the toxicity of 2-chloropyridine,¹⁹ the development of efficient synthetic routes to vinyl-2-fluoropyridinium salts are highly desirable in order to construct 1-(1-arylvinyl)-2-pyridones. However, the synthesis of 1-(1-arylvinyl)-pyridinium salts is still limited. Relles has reported the synthesis of vinyl pyridinium salts from electron-rich acetophenones with SOCl₂ using pyridine as the solvent (eq. 2).²⁰ Novella and Alvarez-Builla have checked the generality of Relles's work.²¹ Substrates with electron-donating groups (e.g., 4-OMe) afford polymers, while substrates with electron-withdrawing groups (e.g., 4-F) do not afford vinyl pyridinium salts (eq. 3). Recently, Zhao and co-workers have reported the synthesis of vinyl pyridinium salts from styrenes with fluoropyridinium salts in the presence of an organoselenium catalyst (eq. 4).²² Very recently, Majirek has reported the pyridination of alkynes in the presence of TfOH (eq. 5). However, this system is limited to alkynes that contain electron-donating groups such as OMe,^{22,23} and


pyridinium salts were not observed using phenylacetylene (eq. 6).

Trifluoromethanesulfonic anhydride (Tf₂O) is a highly electrophilic reagent,²⁴ and its reactions with ketones and base furnish vinyl triflates.^{25,26} Yields can often be improved when using sterically hindered non-nucleophilic bases such as 2,6-di-(*t*-butyl)-4-methylpyridine (DTBMP),²⁷ albeit that the latter is relatively expensive. During our study on vinyl triflates,^{28,29} we obtained N-vinyl-substituted pyridinium salts from acetophenones with Tf₂O in the presence of 2-fluoropyridine^{30–34} as a mild nucleophilic base. Subsequent treatment of the resulting pyridinium salts under basic conditions furnished 1-(1-arylvinyl)pyridin-2(1H)-ones (eq. 7). Herein, we describe the synthesis of 1-(1-arylvinyl)pyridin-2(1H)-ones from acetophenones that contain electron-donating or -withdrawing groups at the benzene ring.


Undheim 1979, Rao 1989




R = H, EWG = Br
R = 4-Cl, 4-OMe, EWG = CN
 Relles 1973 and Novella 1985



R = H, 4-Me
 4-Cl, 4-Br, etc.




Zhao 2017




R = H, 4-Me, 4-OMe
 4-F, 4-Cl, 4-Br, etc.


Majireck 2019



R = 4-OMe




This Work



R = 4-OMe, 4-CO₂Me, 4-CN
 4-F, 4-Cl, 4-Br, etc.

For our initial study, we chose 4'-chloroacetophenone (**1a**) as a model substrate (Table 1). When a CH₂ClCH₂Cl solution of **1a**, 2-fluoropyridine (**2a**) (1.0 equiv), and Tf₂O (1.5 equiv) was stirred for 1 h at 80 °C, and subsequently treated with NaOH aq., vinyl triflate **3a** was obtained in 92% yield together with vinyl pyridone **4aa** in 4% yield (Table 1, entry 1). Increasing the amount of **2a** to e.g. 3.0 equiv resulted in the formation of **4aa** in 79% yield after column chromatography on silica gel (entry 3).

Table 1. Initial Study




entry	2a (equiv)	conv. of 1a ^a	3a ^a	4aa ^{a,b}
1	1.0	97%	92%	4%
2	2.0	86%	42%	53%
3	3.0	99%	5%	93% (79%)

^a Determined by ¹H NMR spectroscopy using 1,1,2,2-tetrachloroethane as an internal standard. ^b The isolated product yield, after flash column chromatography on SiO₂, is given in parentheses.


With the optimized conditions (Table 1, entry 3) in hand, we next examined the substrate scope for the synthesis of vinyl pyridones and the results are summarized in Table 2. Halo-substituted acetophenones **1a–1f** afforded **4aa–4fa** in good yield (entries 1–6). The reaction of acetophenone (**1g**) furnished vinyl pyridone **4ga** in 85% yield (entry 7). *p*-Methoxy-, *o*-methyl-, and *p*-phenyl-substituted acetophenones also worked well (entries 8–10). Substrates with electron-withdrawing groups at the benzene ring (**1k–1m**) afforded the corresponding products (**4ka–4ma**) in moderate to good yield (entries 11–13). For example, the reaction of 1-(4-(methylsulfonyl)phenyl)ethan-1-one (**1l**) furnished **4la** in 52% yield (entry 12). Increasing the proportion of 4-fluoropyridine (**2a**) to 5 equiv generated vinyl pyridone **4la** in 64% yield (entry 13). 1-(Naphthalen-1-yl)ethan-1-one (**1n**) and 2-(naphthalen-1-yl)ethan-1-one (**1o**) also worked well (entries 16 and 17). The reaction of **1f** and 2-fluoro-4-methylpyridine (**2b**) with Tf₂O gave the corresponding pyridine (**4fb**) in good yield (entry 18). The reaction of 1-(4-chlorophenyl)propan-1-one (**1p**) and 1,2-diphenylethan-1-one (**1q**) afforded the corresponding products with high stereoselectivity in 83% and 60% yield, respectively (entries 19 and 20). The present synthesis of vinyl pyridones can also be carried out on the gram scale, furnishing **4fa** in 88% yield and **4ha** in 97% yield (eq. 3 and 4).

Table 2. Substrate Scope^a




entry	1	2	4	yield [b,c]
1	4-Cl (1a)	2a	4aa	93% (79%)
2	3-Cl (1b)	2a	4ba	77% (66%)
3	2-Cl (1c)	2a	4ca	69% (56%)
4	R ¹ = 4-F (1d)	2a	4da	99% (89%)
5	R ¹ = 4-Br (1e)	2a	4ea	91% (84%)
6	R ¹ = 4-I (1f)	2a	4fa	99% (88%)
7	R ¹ = H (1g)	2a	4ga	(85%)
8	R ¹ = 4-OMe (1h)	2a	4ha	99% (96%)
9	R ¹ = 2-Me (1i)	2a	4ia	(90%)
10	R ¹ = 4-Ph (1j)	2a	4ja	99% (98%)
11	R ¹ = 4-CO ₂ Me (1k)	2a	4ka	74% (58%)
12	R ¹ = 4-SO ₂ Me (1l)	2a	4la	(52%)
13	R ¹ = 4-CN (1m)	2a	4ma	54% (46%)
14 [d]	R ¹ = 4-SO ₂ Me (1l)	2a	4la	(63%)
15 [d]	R ¹ = 4-CN (1m)	2a	4ma	72% (64%)
16	1n	2a	4na	(99%)
17	1o	2a	4oa	90% (92%)
18	1f	2b	4fb	64% (54%)
19	1p	2a	4pa	86% (83%) Z/E = 85/15 [e]
20	1q	2a	4qa	72% (60%) Z/E = 88/12 [e]

^a **1** (0.5 mmol), **2** (1.5 mmol, 3 equiv), Ti_2O_3 (1.5 equiv), $\text{CH}_2\text{ClCH}_2\text{Cl}$ (2 mL), 80°C , 1 h. ^b Determined by ^1H NMR spectroscopy using 1,1,2,2-tetrachloroethane as the internal standard. ^c The isolated product yield, after flash column chromatography on SiO_2 , is given in parentheses. ^d **2a** (5 equiv) was used. ^e The stereoselectivity was determined by ^1H NMR analysis of the crude reaction mixture.



According to previous work on vinyl triflates by other groups, vinyl triflates that contain electron-donating groups at the *para* position of the benzene ring easily afford vinyl cations, which can be trapped by nucleophiles such as nitriles³⁵ or sulfoxides.³⁶ Therefore, we examined the reaction using vinyl triflates **3a**, **3g**, and **3k** or aryl alkynes **1h'**, **1g'**, and **1k'** to gain mechanistic insights. When a $\text{CH}_2\text{ClCH}_2\text{Cl}$ solution of **3a**, **3g**, and **3k** and 2-fluoropyridine was stirred at 80°C for 1 h, followed by treatment with NaOH aq., the desired product was obtained in 1%, 12%, and 0% yield, respectively. Similar to the result of Majireck *et al.*,²³ the reaction of aryl alkynes **1g'** and **1k'** with TiOH and 2-fluoropyridine did not afford vinyl pyridones effectively. These results suggest that it is not very likely that both vinyl triflates (**3**) and vinyl cations play major roles as intermediates in this transformation (Scheme 1).

Scheme 1. Control experiments




When a pyridinium salt, which was obtained from treating 2-fluoropyridine (**2a**) with Ti_2O_3 ,³⁴ in CDCl_3 was exposed to acetophenone **1a** (4-Cl) or **1h** (4-OMe) at rt for 1 h, the corresponding vinyl pyridinium salts were generated.

Although the details of the underlying mechanism still remain unclear at this stage, a plausible mechanism is shown in Scheme 2. The generated trifluoromethanesulfonyl pyridinium salt reacts with a ketone via a concerted pathway to give pyridinium intermediate **A**. Another 2-fluoropyridine abstracts a proton to give vinyl pyridinium **B**, which undergoes hydrolysis to furnish the final product. An alternative mechanism is based on Neuhoof's work.³⁷ The reaction of


acetophenone with TiF_2O gives carbocation intermediate, which is trapped by 2-fluoropyridine to afford pyridinium intermediate **A**.

Scheme 2. Plausible reaction mechanism for the synthesis of 1-(1-arylvinyl)pyridin-2(1*H*)-ones from ketones and 2-fluoropyridine



As the products contain enamide moieties, they are amenable to further transformations (Scheme 3). For example, a palladium-catalyzed C–H arylation reaction³⁸ of **4ha** with aryl iodide **5** forms trisubstituted alkene **6ha** in 81% yield. A black-light-induced Heck-type reaction with bromomalonate furnished the corresponding alkylated product (**8ha**) in 56% yield.

Scheme 3. Subsequent transformation of the obtained vinyl pyridones



In summary, we have developed a synthetic route to vinyl pyridones from acetophenones and 2-fluoropyridine with TiF_2O as an activator. This strategy is characterized by an excellent functional-group tolerance and procedural simplicity. The generated vinyl pyridones are easily transformed via C–H arylation and radical alkylation reactions. Further

mechanistic studies are currently in progress and the corresponding results will be reported in due course.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedure, spectroscopic data, and copies of spectra (PDF)

AUTHOR INFORMATION

Corresponding Author

Takuji Kawamoto – Department of Applied Chemistry, Yamaguchi University, Ube, Yamaguchi 755-8611, Japan; or-cid.org/0000-0002-1845-4700; E-mail: tak102@yamaguchi-u.ac.jp

Authors

Shunya Ikeda – Department of Applied Chemistry, Yamaguchi University, Ube, Yamaguchi 755-8611, Japan

Akio Kamimura – Department of Applied Chemistry, Yamaguchi University, Ube, Yamaguchi 755-8611, Japan; or-cid.org/0000-0002-3060-4265

ACKNOWLEDGMENT

This work was partially supported by a JSPS Grant-in-Aid for Young Scientists (B) (18K14222), the Yamagin Regional Enterprise Support Foundation, and the Tobe Maki Scholarship Foundation. The authors would like to thank Central Glass Co., Ltd. (Japan) for a generous gift of TiF_2O .

REFERENCES

- Hibi, S.; Ueno, K.; Nagato, S.; Kawano, K.; Ito, K.; Norimine, Y.; Takenaka, O.; Hanada, T.; Yonaga, M. Discovery of 2-(2-Oxo-1-Phenyl-5-Pyridin-2-Yl)-1,2-Dihydropyridin-3-Yl)Benzonitrile (Perampanel): A Novel, Noncompetitive α -Amino-3-Hydroxy-5-Methyl-4-Isoxazolepropanoic Acid (AMPA) Receptor Antagonist. *Journal of Medicinal Chemistry* **2012**, 55 (23), 10584–10600. <https://doi.org/10.1021/jm301268u>.
- Li, L.; Jiang, X.; Huang, S.; Ying, Z.; Zhang, Z.; Pan, C.; Li, S.; Wang, X.; Zhang, Z. Discovery of Highly Potent 2-Sulfonyl-Pyrimidinyl Derivatives for Apoptosis Inhibition and Ischemia Treatment. *ACS Medicinal Chemistry Letters* **2017**, 8 (4), 407–412. <https://doi.org/10.1021/acsmedchemlett.6b00489>.
- Cox, R. M.; Toots, M.; Yoon, J.-J.; Sourimant, J.; Ludeke, B.; Fearn, R.; Patti, J.; Lee, E.; Vernachio, J.; Plemper, R. K.; Bourque, E. Development of an Allosteric Inhibitor Class Blocking RNA Elongation by the Respiratory Syncytial Virus Polymerase Complex. *J. Biol. Chem.* **2018**, 293 (43), 16761–16777. <https://doi.org/10.1074/jbc.RA118.004862>.
- Wang, Z.; Yu, Z.; Kang, D.; Zhang, J.; Tian, Y.; Daelemans, D.; De Clercq, E.; Pannecouque, C.; Zhan, P.; Liu, X. Design, Synthesis and Biological Evaluation of Novel Acetamide-Substituted Doravirine and Its

- Prodrugs as Potent HIV-1 NNRTIs. *Bioorg. Med. Chem.* **2019**, *27* (3), 447–456. <https://doi.org/10.1016/j.bmc.2018.12.039>.
- (5) Ettari, R.; Cerchia, C.; Maiorana, S.; Guccione, M.; Novellino, E.; Bitto, A.; Grasso, S.; Lavecchia, A.; Zappalà, M. Development of Novel Amides as Noncovalent Inhibitors of Immunoproteasomes. *ChemMedChem* **2019**, *14* (8), 842–852. <https://doi.org/10.1002/cmdc.201900028>.
 - (6) Namasivayam, V.; Vanangamudi, M.; Kramer, V. G.; Kurup, S.; Zhan, P.; Liu, X.; Kongsted, J.; Byraredy, S. N. The Journey of HIV-1 Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs) from Lab to Clinic. *Journal of Medicinal Chemistry* **2019**, *62* (10), 4851–4883. <https://doi.org/10.1021/acs.jmedchem.8b00843>.
 - (7) Dorich, S.; Cox, J. H.; Burch, J. D.; Chagnon, F.; Chen, B.; Léger, S.; St-Onge, M.; Fader, L. D. Novel Pyridone EP4 Agonists Featuring Allylic Alcohol \$omega\$-Chains. *Bioorg. Med. Chem. Lett.* **2020**, *30* (10), 127104. <https://doi.org/10.1016/j.bmcl.2020.127104>.
 - (8) Hillenbrand, J.; Ham, W. S.; Ritter, T. C-H Pyridonation of (Hetero-)Arenes by Pyridinium Radical Cations. *Organic Letters* **2019**, *21* (13), 5363–5367. <https://doi.org/10.1021/acs.orglett.9b02054>.
 - (9) Kuriyama, M.; Hanazawa, N.; Abe, Y.; Katagiri, K.; Ono, S.; Yamamoto, K.; Onomura, O. N - And O -Arylation of Pyridin-2-Ones with Diaryliodonium Salts: Base-Dependent Orthogonal Selectivity under Metal-Free Conditions. *Chem. Sci.* **2020**, *11* (31), 8295–8300. <https://doi.org/10.1039/d0sc02516j>.
 - (10) Sun, C.; Qi, X.; Min, X.-L.; Bai, X.-D.; Liu, P.; He, Y. Asymmetric Allylic Substitution-Isomerization to Axially Chiral Enamides: Via Hydrogen-Bonding Assisted Central-to-Axial Chirality Transfer. *Chem. Sci.* **2020**, *11* (37), 10119–10126. <https://doi.org/10.1039/d0sc02828b>.
 - (11) Breivik, H.; Ottersen, T.; Undheim, K. Regioselective Addition of 2-Pyridinone to C-2 of a Haloalkyne. *Acta Chemica Scandinavica* **1979**, *33b*, 365–369. <https://doi.org/10.3891/acta.chem.scand.33b-0365>.
 - (12) Yadla, R.; Rehman, H.; Rao, J. M.; Mahesh, V. K. Aromatic Vs Diene Reactivity of 2(1H)-Pyridinone and Its Derivatives. *Tetrahedron* **1989**, *45* (22), 7093–7098. [https://doi.org/10.1016/S0040-4020\(01\)89177-8](https://doi.org/10.1016/S0040-4020(01)89177-8).
 - (13) Mukaiyama, T.; Usui, M.; Shimada, E.; Saigo, K. A CONVENIENT METHOD FOR THE SYNTHESIS OF CARBOXYLIC ESTERS. *Chemistry Letters* **1975**, *4* (10), 1045–1048. <https://doi.org/10.1246/cl.1975.1045>.
 - (14) Bald, E.; Saigo, K.; Mukaiyama, T. A FACILE SYNTHESIS OF CARBOXAMIDES BY USING 1-METHYL-2-HALOPYRIDINIUM IODIDES AS COUPLING REAGENTS. *Chemistry Letters* **1975**, *4* (11), 1163–1166. <https://doi.org/10.1246/cl.1975.1163>.
 - (15) Mukaiyama, T.; Ikeda, S.; Kobayashi, S. A NOVEL METHOD FOR THE PREPARATION OF VARIOUS 2-PYRIDYL SULFIDES FROM ALCOHOLS. *Chemistry Letters* **1975**, *4* (11), 1159–1162. <https://doi.org/10.1246/cl.1975.1159>.
 - (16) Kobayashi, S.; Tsutsui, M.; Mukaiyama, T. A CONVENIENT METHOD FOR THE TRANSFORMATION OF ALCOHOLS TO ALKYL IODIDES USING 2-FLUOROPYRIDINIUM SALT. *Chemistry Letters* **1976**, *5* (4), 373–374. <https://doi.org/10.1246/cl.1976.373>.
 - (17) Hu, G.; Xu, J.; Li, P. Synthesis of N-Alkylated 2-Pyridones through Pummerer Type Reactions of Activated Sulfoxides and 2-Fluoropyridine Derivatives. *Org. Biomol. Chem.* **2018**, *16* (22), 4151–4158. <https://doi.org/10.1039/C8OB00860D>.
 - (18) Liao, C.; Li, J.; Chen, X.; Lu, J.; Liu, Q.; Chen, L.; Huang, Y.; Li, Y. Selective Synthesis of Pyridyl Pyridones and Oxydipyridines by Transition-Metal-Free Hydroxylation and Arylation of 2-Fluoropyridine Derivatives. *Org. Biomol. Chem.* **2020**, *18* (6), 1185–1193. <https://doi.org/10.1039/C9OB02661D>.
 - (19) Gehring, P. J.; Torkelson, T. R.; Oyen, F. A Comparison of the Lethality of Chlorinated Pyridines and a Study of the Acute Toxicity of 2-Chloropyridine. *Toxicology and Applied Pharmacology* **1967**, *11* (2), 361–371. [https://doi.org/10.1016/0041-008X\(67\)90079-8](https://doi.org/10.1016/0041-008X(67)90079-8).
 - (20) Relles, H. M. Thionyl Chloride-Pyridine Chemistry. II.¹⁸ Synthesis and Reactions of N- α - Styrylpyridinium Salts. *The Journal of Organic Chemistry* **1973**, *38* (8), 1570–1574. <https://doi.org/10.1021/jo00948a026>.
 - (21) Novella, J. L.; Alvarez-Builla, J. Synthesis of 1-(1-Arylethenyl) Pyridinium Salts as Potential Anthelmintic Agents. *Anales de Quimica, Serie C Quimica Organica y Bioquimica* **1985**, *81*, 186–187.
 - (22) Shapiro, J.; Sonberg, J.; Schafer, B.; Williams, C.; Ferris, H.; Reinheimer, E.; Van Wynsberghe, A.; Kriley, C.; Majireck, M. Synthesis, Characterization, and Computational Modeling of N-(1-Ethoxyvinyl)Pyridinium Triflates, an Unusual Class of Pyridinium Salts. *Molecules* **2018**, *23* (2), 413–414. <https://doi.org/10.3390/molecules23020413>.
 - (23) Blades, A. M.; McConnell, D. L.; Gomes Rodrigues, D.; Keyes, P. V.; Sonberg, J. C.; Anthony, C. E.; Rachad, S.; Simone, O. M.; Sullivan, C. F.; Shapiro, J. D.; Williams, C. C.; Schafer, B. C.; Goodman, J. A.; Majireck, M. Synthesis of Bench-Stable N-Quaternized Ketene N,O-Acetals and Preliminary Evaluation as Reagents in Organic Synthesis; preprint; 2019. <https://doi.org/10.26434/chemrxiv.7381952.v1>.
 - (24) Martínez, A. G.; Subramanian, L. R.; Hanack, M.; Williams, S. J.; Régnier, S. Trifluoromethanesulfonic Anhydride. 1–17. <https://doi.org/10.1002/047084289X.rt247.pub3>.
 - (25) Ritter, K. Synthetic Transformations of Vinyl and Aryl Triflates. *Synthesis* **1993**, *1993* (08), 735–762. <https://doi.org/10.1055/s-1993-25931>.
 - (26) Baraznenok, I. L.; Nenajdenko, V. G.; Balenkova, E. S. Chemical Transformations Induced by Triflic Anhydride. *Tetrahedron* **2000**, *56* (20), 3077–3119. [https://doi.org/10.1016/S0040-4020\(00\)00093-4](https://doi.org/10.1016/S0040-4020(00)00093-4).
 - (27) Stang, P. J.; Treptow, W. Single-Step Improved Synthesis of Primary and Other Vinyl Trifluoromethanesulfonates. *Synthesis* **1980**, *1980* (4), 283–284. <https://doi.org/10.1055/s-1980-28991>.
 - (28) Kawamoto, T.; Sasaki, R.; Kamimura, A. Synthesis of A-Trifluoromethylated Ketones from Vinyl Triflates in the Absence of External Trifluoromethyl Sources. *Angew. Chem. Int. Ed.* **2017**, *56* (5), 1342–1345. <https://doi.org/10.1002/anie.201608591>.

- (29) Kawamoto, T.; Sasaki, R.; Kamimura, A.; Matsubara, H. The Regioselective Trifluoromethylation of 1,3-Bis(Vinyl Triflates) in the Absence of External Trifluoromethyl Sources. *J. Fluorine Chem.* **2019**, *221*, 66–69. <https://doi.org/10.1016/j.jfluchem.2019.04.002>.
- (30) Medley, J. W.; Movassaghi, M. Direct Dehydrative N-Pyridinylation of Amides. *The Journal of Organic Chemistry* **2009**, *74* (3), 1341–1344. <https://doi.org/10.1021/jo802355d>.
- (31) Bechara, W. S.; Pelletier, G.; Charette, A. B. Chemoselective Synthesis of Ketones and Ketimines by Addition of Organometallic Reagents to Secondary Amides. *Nature Chem.* **2012**, *4*, 228–234. <https://doi.org/10.1038/nchem.1268>.
- (32) Xiao, K.-J.; Wang, A.-E.; Huang, P.-Q. Direct Transformation of Secondary Amides into Secondary Amines: Triflic Anhydride Activated Reductive Alkylation. *Angewandte Chemie International Edition* **2012**, *51* (33), 8314–8317. <https://doi.org/10.1002/anie.201204098>.
- (33) Peng, B.; Geerdink, D.; Farès, C.; Maulide, N. Chemoselective Intermolecular α -Arylation of Amides. *Angewandte Chemie International Edition* **2014**, *53* (21), 5462–5466. <https://doi.org/10.1002/anie.201402229>.
- (34) White, K. L.; Mewald, M.; Movassaghi, M. Direct Observation of Intermediates Involved in the Interruption of the Bischler–Napieralski Reaction. *The Journal of Organic Chemistry* **2015**, *80* (15), 7403–7411. <https://doi.org/10.1021/acs.joc.5b01023>.
- (35) Martinez, A. G.; Herrera, A.; Martinez, R.; Teso, E.; Garcia, A.; Osio, J.; Pargada, L.; Unanue, R.; Subramanian, L. R.; Hanack, M. A New and Convenient Synthesis of Alkyl and Aryl Pyrimidines. *J. Heterocycl. Chem.* **1988**, *25*, 1237–1241.
- (36) Zawodny, W.; Teskey, C. J.; Mishevskaya, M.; Völkl, M.; Maryasin, B.; González, L.; Maulide, N. α -Functionalisation of Ketones Through Metal-Free Electrophilic Activation. *Angewandte Chemie International Edition* **2020**, *59* (47), 20935–20939. <https://doi.org/10.1002/anie.202006398>.
- (37) Müller, E.; Langerbeck, A.; Neuhoff, H. Überführung α , β -Ungesättigter Ketone in Triene. *Berichte der deutschen chemischen Gesellschaft (A and B Series)* **1944**, No. 3–4, 141–152. <https://doi.org/10.1002/cber.19440770301>.
- (38) Zhang, W.; Yu, W.; Yan, Q.; Liu, Z.; Zhang, Y. Synthesis of Substituted Oxazoles via Pd-Catalyzed Tandem Oxidative Cyclization. *Org. Chem. Front.* **2017**, *4* (12), 2428–2432. <https://doi.org/10.1039/C7QO00517B>.

Authors are required to submit a graphic entry for the Table of Contents (TOC) that, in conjunction with the manuscript title, should give the reader a representative idea of one of the following: A key structure, reaction, equation, concept, or theorem, etc., that is discussed in the manuscript. Consult the journal's Instructions for Authors for TOC graphic specifications.

