

Quantitative profiling and dynamics of mRNA modifications in *Escherichia coli*

Dimitar Plamenov Petrov¹, Steffen Kaiser^{2,3}, Stefanie Kaiser^{2,3*} and Kirsten Jung^{1*}

¹ Department of Biology I, Microbiology, Ludwig-Maximilians-University Munich, Martinsried, Germany.

² Department of Chemistry, Ludwig-Maximilians-University Munich, Munich, Germany. ³ Department of Pharmacy, Goethe-University Frankfurt, Frankfurt, Germany.

* To whom correspondence should be addressed. Email: jung@lmu.de; kellner@pharmchem.uni-frankfurt.de

ABSTRACT

mRNA methylation is an important regulator of many physiological processes in eukaryotes but has not been studied in depth in prokaryotes. In contrast to the large number of eukaryotic mRNA modifications that have been described, *N*⁶-methyladenosine (*m*⁶A) is the only modification of bacterial mRNA identified to date. Here, we used a gel electrophoresis-based RNA separation method and quantitatively analyzed the mRNA-specific modification profile of *Escherichia coli* using mass spectrometry. In addition to *m*⁶A, we provide evidence for the presence of 7-methylguanosine (*m*⁷G), and we found first hints for 5-methylcytidine (*m*⁵C), *N*⁶,*N*⁶-dimethyladenosine (*m*^{6,6}A), 1-methylguanosine (*m*¹G), 5-methyluridine (*m*⁵U), and pseudouridine (Ψ) in the mRNA of *E. coli*, which implies that *E. coli* has a complex mRNA modification pattern. Furthermore, we observed changes in the abundance of some mRNA modifications during the transition of *E. coli* from the exponential growth to the stationary phase as well as upon exposure to stress. This study reveals a previously underestimated level of regulation between transcription and translation in bacteria.

INTRODUCTION

Chemical modification of DNA is a well-known epigenetic regulatory mechanism. The recent development of highly sensitive analytical techniques has discovered increasing numbers of RNA modifications leading to new aspects of RNA regulation in the emerging field of epitranscriptomics. To date, more than 160 RNA modifications have been described in all domains of life (1,2). Eukaryotes carry both reversible and irreversible RNA modifications in their messenger RNA (mRNA). These mRNA modifications play an important role in RNA editing, splicing, and 5'-capping, and some also have regulatory effects on RNA stability and translational fidelity [reviewed by (3) (4)].

In eukaryotes, tRNA and rRNA modifications can affect the whole protein synthesis in the cell, whereas several mRNA modifications have been shown to be posttranscriptional regulators of specific gene expression programs involved in cell development and homeostasis (5), (6). Complex machineries, including modification-specific methyltransferases, demethylases, and binding proteins, recognize or change the methylation state of specific mRNAs and consequently affect their fates and functions (7). *N*⁶-methyladenosine (*m*⁶A) is the most abundant mRNA modification in eukaryotes and is the best studied example of how mRNA methylation regulates both mRNA stability and gene expression [reviewed by (8)].

Pseudouridine (Ψ), one of the most abundant modifications found in all types of RNA, is also present in the coding sequence of mRNAs and has been reported to affect translational speed and tRNA selection by ribosomes (9).

Although bacteria have been used as model organisms to study modifications and modifying enzymes of tRNA and rRNA (10,11), knowledge regarding the mRNA modification profile of prokaryotes is extremely limited. Working with bacterial mRNA is challenging because it lacks a poly(A)-tail, which can be used for its enrichment in experimental methods, and its average half-life time is very short (in the range of minutes) (12). Despite these challenges, in 2015, high-resolution transcriptome-wide *m*⁶A profiling for the two bacterial model organisms *Escherichia coli* and *Pseudomonas aeruginosa* revealed both conserved and distinct distribution patterns of this modification (13). The *m*⁶A/A ratios observed (>0.2%) were comparable to those of mammals (0.1%–0.4%) and meiotic yeast (0.25%). Most *m*⁶A-modified transcripts were associated with respiration, amino acid metabolism, stress response, and small RNA species, suggesting a potential regulatory role for mRNA methylation in bacteria. Therefore, it was suggested that RNA methylation might be involved in the adaptation of bacteria to different conditions and developmental stages, as has been demonstrated for eukaryotic cells (13). Unlike *E. coli* and *Pseudomonas* spp., two other Gram-negative cyanobacteria (*Anabaena*

sp. PCC 7120 and *Synechocystis* sp. PCC 6803) and representative species of Gram-positive bacteria (*Staphylococcus aureus* and *Bacillus subtilis*) showed low m⁶A/A ratios (<0.04% and <0.08%, respectively) (13). Finally, 5-methylcytidine (m⁵C) has been mapped in the mRNA of the archaeon *Sulfolobus solfataricus* (14).

Because m⁶A is the only described mRNA modification in bacteria, we were interested whether other modifications found in eukaryotes are also present in bacterial mRNA. To address this question, we used a gel electrophoresis-based method to separate RNA from *E. coli* and analyzed the individual fractions by isotope dilution nucleoside mass spectrometry (LC-MS/MS). Our results provide evidence for the presence of not only m⁶A but also 7-methylguanosine (m⁷G), and first hints for 5-methylcytidine (m⁵C), N⁶,N⁶-dimethyladenosine (m^{6,6}A), 1-methylguanosine (m¹G), 5-methyluridine (m⁵U), and pseudouridine (Ψ) in *E. coli* mRNA. The abundance of some of these modifications is altered during the transition of cells from the exponential growth to the stationary phase. Furthermore, stress conditions like oxygen limitation or acid stress led to changes in the abundance of some mRNA modifications. This study sheds light on a thus far neglected layer of regulation between transcription and translation in bacteria.

RESULTS

Profiling of RNA modifications in *E. coli* MG1655

To investigate the complexity of mRNA modification in bacteria, we used the model organism *E. coli* MG1655. Since RNA sequencing is only possible for a limited number of modifications (15), and negative controls are lacking because the modification machinery in bacteria is unknown, we used an LC-MS/MS-based approach. As a starting point, we performed qualitative profiling and comparison of the modifications in the three main RNA types: mRNA (including long noncoding RNA), rRNA (23S and 16S rRNA), and tRNA (including small RNA species less than 110 nucleotides (nt) in length). For this purpose, *E. coli* cells were grown in LB medium and harvested in the early exponential growth phase. Total RNA was isolated using phenol–chloroform–isoamyl alcohol (PCI) extraction and separated into tRNA and large RNA species containing both rRNA and mRNA (rRNA+mRNA fraction) by an established size exclusion chromatography (SEC) method (16). The successful removal of tRNA from the

rRNA+mRNA fraction was verified by chip gel electrophoresis. We observed a clean distinction between the tRNA and rRNA+mRNA fractions (Supplemental Fig. S1A). As an additional control for the efficiency of tRNA separation, we measured m⁶A levels in the tRNA and rRNA+mRNA fractions in the *E. coli* wild-type and the mutant $\Delta trmM$, which lacks the only known methyltransferase for m⁶A methylation of tRNA (17). Whereas the m⁶A abundance in the rRNA+mRNA fraction remained similar in both strains, only substoichiometric amounts of m⁶A were detectable after LC-MS/MS analysis in the tRNA fraction of the mutant in contrast to a high level in the wild type, which shows efficient separation between the tRNA and rRNA+mRNA fractions (Supplemental Figs. S1B-C). Next, the purified rRNA+mRNA fraction was separated into rRNA and mRNA using a commercially available kit for the oligonucleotide-based removal of rRNA. The efficiency of separation was verified by RT-qPCR, and for each mRNA sample, we observed at least a 500-fold decrease in the rRNA amount compared to samples taken before oligonucleotide-based rRNA depletion (Supplementary Fig. S2). Finally, each RNA fraction was enzymatically hydrolyzed, and the resulting nucleoside mixtures were subjected to LC-MS/MS for qualitative analysis of 54 RNA modifications (Fig. 1).

For each of the three studied RNA types—tRNA, rRNA, and mRNA—we observed a complex profile of modifications, as summarized in Fig. 1. As expected, we found the highest chemical diversity in the tRNA fraction, with 26 different modified nucleosides detected. We confirmed numerous known tRNA modifications, and in addition, we detected N¹-methyladenosine (m¹A), m^{6,6}A, 2'-O-methyladenosine (Am), and m⁵C in *E. coli* tRNA. A lower chemical diversity was observed in the mRNA fraction. Here, we detected 15 nucleosides overlapping with either tRNA or rRNA. Alongside the only previously known modification m⁶A, in the mRNA of *E. coli* MG1655, we identified the presence of m¹A, m²A, m^{6,6}A, Am, m⁵C, 2'-O-methylcytidine (Cm), m¹G, m²G, m⁷G, 2'-O-methylguanosine (Gm), m⁵U, inosine (I), and Ψ. In addition, we detected a modified cytidine derivative that is modified at both the ribose and the nucleobase (m^xCm). According to the literature, this modification is most likely N⁴,2'-O-dimethylcytidine (m⁴Cm), but comparison with synthetic standards of m⁴Cm and m⁵Cm revealed co-elution in our system, and so the existence of m⁵Cm cannot be excluded. In the SEC-purified rRNA fraction (16), we detected 16 modified nucleosides.

Three of these modifications, namely, N^6 , 2'-O-dimethyladenosine (m^6Am), 5-formylcytidine (f^5C), and inosine, have not been observed previously in the rRNA of *E. coli* (10). Taken together, the modifications m^2A , m^6A , $m^{6,6}A$, m^5C , m^xCm , Cm , m^1G , m^7G , Gm , m^5U , I , and Ψ were detected in all three types of RNA in *E. coli* MG1655 (Fig. 1). Furthermore, m^2G was detectable in the rRNA and mRNA, but not in the tRNA fraction, whereas m^1A and Am were found in the tRNA and mRNA fractions but not in the rRNA. The modifications m^6Am and m^3U

were detected exclusively within the rRNA fraction. However, owing to their low abundance, they cannot be used as internal controls for rRNA contamination. N^6 -threonylcarbamoyladenosine (t^6A), 2-thiocytidine (s^2C), 4-thiouridine (s^4U), and dihydrouridine (D) were found only in the tRNA fraction and might be suitable indicators for contamination with tRNA. Under our test conditions, 25 modifications known from both prokaryotes and eukaryotes were below the limit of detection in our RNA samples (Fig. 1).




Figure 1. RNA modification profile of early log-phase *E. coli* MG1655. Cells were aerobically cultivated in LB medium and harvested, and total RNA was isolated. The main RNA types were separated by size exclusion chromatography and subsequent rRNA depletion with commercial kits. After enzymatic hydrolysis, mass spectrometry was used to screen for 54 known modified nucleosides. 29 modified nucleosides were found in the mRNA, rRNA*, and tRNA fractions as indicated (left), and 25 were not found (right). *The results for rRNA represent the modifications found in the rRNA+mRNA fraction after tRNA removal by SEC, where the minor contribution of mRNA to the total modification levels is neglectable due to the excessive amounts of rRNA.

Preparation of samples for quantitative analysis of mRNA and rRNA modifications

Our qualitative profiling of modifications in the three major RNA types obtained by commonly applied RNA purifications technologies revealed an unexpected high chemical diversity of modified nucleosides of mRNA in *E. coli*. However, the high sensitivity of the MS method depends on RNA samples with well-defined purity, because e.g., tRNA or rRNA

contaminations would go unnoticed. Furthermore, it is important to mention that RNA handling itself can induce the formation of RNA modification artefacts or loss of certain modified nucleosides (18). For a rigorous qualitative and quantitative assessment of mRNA modifications, both aspects need to be considered already during the preparation of bacterial mRNA for subsequent MS-based analysis. We decided that agarose gel electrophoresis-based size selection of RNAs is

the most stringent method for successful separation of mRNA and rRNA, which bears the lowest risk for RNA modification artefact formation.

Total bacterial RNA was loaded on denaturing agarose gels, and after electrophoresis, four gel pieces per lane were excised. The RNA was extracted using a commercially available kit. According to size we obtained the following fractions: fraction 1 (23S rRNA), fraction 2 (mRNAs and noncoding RNAs of 1,700 to 2,700 bases), fraction 3 (16S rRNA) and fraction 4 (mRNAs and noncoding RNAs of 400 to 1,100 bases) (Fig. 2A). The quality of each fraction was tested by chip gel electrophoresis and by running the extracted RNA on a denaturing agarose gel (Fig. 2A). These controls indicated that we obtained efficient separation between the rRNA and mRNA fractions.

Due to their abundance, 16S and 23S rRNAs are a constant source for the contamination of mRNA modification signals (19). Therefore, we used RT-qPCR and RNA deep sequencing (RNA-Seq) to analyze the composition of the mRNA-containing fractions 2 and 4. We first determined the ratio of mRNA:rRNA using the two different internal calibrators *alaS* and *recA* for the long and short mRNA fractions, respectively. We observed at least a 1,000-fold increase in the mRNA:rRNA ratio in both fractions when compared to the total RNA samples (Fig. 2B, Supplemental Fig. S3A). We got a similar result for mRNA samples obtained using the commercial oligonucleotides-based kits, however, these kits appear to selectively deplete 16S rRNA while substantial amounts of 23S rRNA remain (Supplemental Fig. S2).

A second RT-qPCR experiment focused on the possibility of a co-purification of rRNA fragments in the mRNA-containing fractions 2 and 4. We performed RT-qPCR using primer pairs binding to three different regions of the rRNA sequences (5'-, middle and 3'-region). These results revealed that 16S rRNA and, to a lesser extent, 23S rRNA had been partially degraded at the 3' end (Supplemental Fig. S3B). We found a contamination of fractions 2 and 4

with approximately 10% rRNA fragments (Supplemental Fig. S3C).

To further define the purity of our mRNA fractions 2 and 4, we subjected all four fractions as well as total RNA to deep sequencing. As expected, fraction 1 (23S rRNA) and fraction 3 (16S rRNA) contained the expected rRNA, but in addition 1.4% and 1.3% mRNA, respectively (Fig. 2C). Worryingly, the RNA-Seq results of the total read samples revealed an mRNA:rRNA ratio of only 14:86 for fraction 2 and 16:83 for fraction 4 (Fig. 2C), which is clearly different from our RT-qPCR data (Fig. 2B). However, because the mRNA in a given fraction is a mixture of many different sequences with low abundance, it is likely that some mRNA targets were amplified less efficiently than the rRNAs with many of the same targets during library preparation for RNA-Seq. This factor can be compensated by removing duplicates that accumulate during the PCR amplification steps of library preparation. Taking this factor into account, we obtained an mRNA:rRNA ratio of 53:47 for fraction 2 and 58:42 for fraction 4 (Fig. 2D). A comparison of the 23S and 16S rRNA reads coverage between the total RNA and fractions 2 and 4 indicated also distribution of different rRNA fragments in the gel extracted samples (Fig. 2E). For example, the middle fragment of the 23S rRNA sequence (700-1900 nt) was overrepresented in fraction 2, whereas read coverage of the 3' end (1000-1542 nt) of 16S rRNA was strongly underrepresented in fraction 4 (Fig. 2E). Considering the localization of known modifications in 23S and 16S rRNA (Fig. 2E), these results were important to judge which modifications might be the result of rRNA contaminations in fractions 2 and 4.

In summary, depending on the used method, our quality controls revealed different degrees of contamination with rRNA of the mRNA-enriched fractions 2 and 4. Nevertheless, according to our RNA-Seq controls, the mRNA levels in fractions 2 and 4 were increased about 15-fold compared with fractions 1 and 3 in the worst-case scenario and up to 50-fold in the best-case scenario (Fig. 2).




Figure 2. Analysis of RNA species from *E. coli* after separation by denaturing agarose gel electrophoresis. (A) After denaturing gel electrophoresis, total RNA was separated in four fractions: fraction 1 (23S rRNA), fraction 2 (mRNAs and noncoding RNAs of 1,700 to 2,700 bases), fraction 3 (16S rRNA) and fraction 4 (mRNAs and noncoding RNAs of 400 to 1,100 bases). The fractions were cut from the gel and after extraction analyzed by automated chip gel electrophoresis (Bioanalyzer, Agilent). The fractions were reloaded onto an agarose gel to confirm the efficiency of separation. (B) Percentage of RNA species in the mRNA-enriched fractions 2 and 4 based

on RT-qPCR analysis. Values were calculated based on an assumed ratio of 1% mRNA and 90% rRNA in the total RNA (see *Material and Methods* for detailed description). (C) Molar ratios of RNA species in the four fractions and the total RNA based on RNA-Seq analysis. (D) Molar ratios of RNA species found in the four fractions and the total RNA based on RNA-Seq analysis after duplicate removal. (E) Schematic map of the localization of modifications in 23S rRNA and 16S rRNA in *E. coli*, and distribution of the RNA-seq reads coverage of 23S rRNA and 16S rRNA of total RNA and fractions 2 and 4. B-D represent average results of three biological replicates with error bars of standard deviation from the means in B. E shows one representative example.

Quantitative analysis of mRNA and rRNA modifications


We moved on to quantitative LC-MS/MS analysis of the gel-extracted RNA fractions. The quantification of modifications detected in rRNA-enriched fractions 1 (23S rRNA) and 3 (16S rRNA) served as an internal control for our approach. We found m⁶A only in 16S rRNA, whereas m⁶A, m¹G, m⁵U, and m²A were found in 23S rRNA (Fig. 3, Supplemental Fig. S4B, Table 1), which is in agreement with the methylation patterns reported in previous studies [reviewed by (20)]. Moreover, the stoichiometry of the modifications in the 23S and 16S rRNA fractions matched with the theoretical expectations based on the number of modification sites (Supplemental Fig. S4B). When calculating the expected abundance, we considered the fact that 23S rRNA is twice as long as 16S rRNA as the modification abundance would be inversely proportional to the length of the molecule. This effect was well demonstrated by the observed ~40% lower abundance of m⁷G in 23S rRNA (1 site: 1.38‰ ± 0.11‰) compared to 16S rRNA (1 site: 2.63‰ ± 0.07‰), reflecting a ~40% difference in length between the two rRNA types (Fig. 3A, Table 1).

Quantitative LC-MS/MS analysis of the gel-extracted mRNA-containing fractions 2 and 4 was consistent with our qualitative results. We detected m⁶A, m⁶.⁶A, m¹G, m⁷G, m⁵C, m⁵U, and Ψ in both the long and short mRNA-enriched fractions 2 and 4 in stoichiometrically relevant quantities using a stable isotope-labeled internal standard from *Saccharomyces cerevisiae* tRNA (SILIS) (21) (Fig. 3A and Table 1). m²A was also detected in the mRNA fractions at a similar abundance as in the 23S rRNA fraction (Supplemental Fig. S5). However, due to the absence of m²A in the yeast-derived SILIS, further analysis and quantitative assessment of m²A was not possible.

The quantification results for fractions 2 and 4 showed that m⁶A is not the most abundant mRNA modification in *E. coli*. We measured comparable amounts of m⁶A (on average 1.1‰)

to those published previously by Deng *et al.*, (2015), but modifications such as m¹G, m⁷G, m⁵C, m⁵U, and Ψ had higher abundances (Table 1). The most abundant modification in the mRNA-enriched fractions 2 and 4 is Ψ (approximately 12‰). The m⁵C abundance detected in the fractions containing mRNAs was about 3.5‰. The abundance of m⁶.⁶A was comparable to that of m⁶A. For several modifications, we observed different methylation levels even between the two mRNA-enriched fractions. In fraction 2, containing long mRNAs, we measured a 2.5-times higher m⁶A abundance compared to fraction 4, containing short mRNAs. The abundance of m¹G in fraction 2 was also higher than that in fraction 4. Conversely, for m⁷G, we observed a lower abundance in fraction 2 compared to that in fraction 4 (Fig. 3A and Table 1). For other modifications such as m⁵C and Ψ, we did not observe significant differences between the long and short mRNA-enriched fractions.

Since our quality controls have indicated the presence of rRNA contamination in fractions 2 and 4, we assumed a worst-case scenario (using the highest degree of contamination, Fig. 2C) and checked whether such contamination could be sufficient to explain the modification levels observed in the mRNA-enriched fractions 2 and 4. First, we calculated the abundance of each modification coming from the rRNA contamination and subtracted the value from the determined values in fractions 2 or 4. We included in this calculation the location of the known modification sites and correlated this number with the distribution of reads coverage (Fig. 2E). Taking this worst-case scenario, the abundancies of the modifications measured in fractions 2 and 4 could not be explained with co-purified rRNAs. For example, the rRNA-contamination accounts for only about 50% of the measured abundance of m⁷G and m⁶.⁶A in fractions 2 and 4. For modifications such as m⁶A as well as m⁵U or Ψ, the measured levels in fraction 4 were higher than the assumed worst-case rRNA-contamination.



Overall, our mRNA-enriched fractions 2 and 4 showed a reproducible modification profile that is different from the 23S and 16S rRNA modification profiles and quantitatively decoupled (Fig. 3A). For example, m⁶A is lower in fraction 2 compared to the preceding 23S rRNA fraction, whereas modifications such as m⁷G, m⁵C, m⁵U and Ψ are higher. In contrast, m⁶A, m¹G, m⁷G, m⁵U and Ψ are higher in fraction 4 compared with the preceding 16S rRNA fraction, whereas m⁵C is lower and of similar abundance as in fraction 2.

In parallel, we subjected the same total RNAs used for gel electrophoresis of mRNA enrichment to the established oligonucleotide-based rRNA depletion protocol. Despite relying on different strategies, both approaches resulted in the qualitative identification of nearly identical mRNA modifications after LC-MS/MS analysis (Fig. 3B). In addition to m⁶A, both approaches identified the presence of m^{6.6}A, m¹G, m²G, m⁷G, m⁵C, m⁵U, Am, Gm, Cm, m^xCm and Ψ in the mRNA from *E. coli*. Inosine (I), a potential artefact derived from chemical adenosine deamination, was only detected in the rRNA-depletion approach and in the gel-extracted long mRNA-enriched fraction 2, but not in fraction 4.

When comparing the quantitative results of the two approaches, we generally observed

higher modification abundances in the gel-extracted samples compared to those using the rRNA depletion approach (Supplemental Fig. S4A). Nevertheless, the results of both approaches showed similar profiles: Ψ was the most abundant modification, followed by m⁵C, and m⁵U, whereas m⁶A was among the least abundant modifications found in the mRNA-enriched fractions of *E. coli*. Importantly, the mRNA profile was distinct from the tRNA as well as the 23S and 16S rRNA profiles of *E. coli* (Supplemental Fig. S4). The clearest differences between the profiles of the three RNA types were observed in the ratios between different modifications: m^{6.6}A:m⁵C:m⁵U:m⁷G:Ψ ratio was 0:0.4:1:6 in tRNA, 0:2:3:1:15 in 23S rRNA, 5:5:0:2:3 in 16S rRNA, and 2:4:3:2:12 in mRNA (Supplemental Fig. S4). These ratios of RNA modifications are an additional indication that we were able to prepare substantially mRNA-enriched fractions.

In summary, although we do not know exactly to what extent the mRNA fractions are contaminated with full-length rRNAs or their various fragments, the differences in modification quantities and profiles found in rRNA and in the mRNA-enriched fractions give us confidence that m⁶A and m⁷G, but also m¹G, m⁵C, m^{6.6}A m⁵U, and Ψ modifications are present in the mRNA of *E. coli*.

Table 1. Quantitative LC-MS/MS analysis of the fractions 1-4 (Fig. 3A), tRNA after SEC, and mRNA after rRNA depletion isolated from total RNA of *E. coli* MG1655. Cells were cultivated in LB medium for 2 h (exponential growth phase) or 8 h (stationary phase). The values represent the quantities of modified nucleosides after normalization to their respective canonical precursors (per 1,000 nucleosides). Results represent the mean and standard deviation (SD) of at least three biological replicates measured in technical triplicates.

Time	RNA type	m ⁶ A	m ² A	m ^{6.6} A	m ¹ G	m ⁷ G	m ⁵ C	m ⁵ U	Ψ
2h	mRNA (rRNA depletion)	0.50 ± 0.15	1.44 ± 0.08	1.18 ± 0.77	0.85 ± 0.29	0.92 ± 0.22	1.17 ± 0.66	1.35 ± 0.83	5.88 ± 1.83
2h	fraction 2 (long mRNA)	1.42 ± 0.28	1.08 ± 0.18	1.75 ± 0.70	2.09 ± 0.47	2.34 ± 0.45	3.74 ± 0.65	2.73 ± 0.51	12.15 ± 1.78
8h	fraction 2 (long mRNA)	2.02 ± 0.28	1.30 ± 0.41	0.84 ± 0.24	3.38 ± 0.32	2.25 ± 0.55	2.87 ± 0.55	3.55 ± 0.66	15.40 ± 2.28
2h	fraction 4 (short mRNA)	0.56 ± 0.09	1.23 ± 0.36	2.01 ± 0.55	1.48 ± 0.40	3.08 ± 0.47	3.37 ± 0.61	2.12 ± 0.77	11.94 ± 2.94
8h	fraction 4 (short mRNA)	0.69 ± 0.19	1.16 ± 0.21	2.55 ± 0.66	0.96 ± 0.10	3.61 ± 0.29	4.75 ± 0.55	2.12 ± 0.25	12.21 ± 1.78
2h	fraction 1 (23S rRNA)	1.85 ± 0.42	1.33 ± 0.13	0.16 ± 0.19	2.81 ± 0.27	1.38 ± 0.11	2.16 ± 0.35	2.52 ± 0.57	13.71 ± 1.21
8h	fraction 1 (23S rRNA)	2.09 ± 0.07	1.30 ± 0.10	0.19 ± 0.24	3.05 ± 0.14	1.46 ± 0.09	2.45 ± 0.69	2.52 ± 0.20	14.50 ± 1.01
2h	fraction 3 (16S rRNA)	0.19 ± 0.09	0.10 ± 0.03	5.32 ± 0.97	0.14 ± 0.11	2.63 ± 0.07	5.82 ± 0.35	0.33 ± 0.10	4.07 ± 0.91
8h	fraction 3 (16S rRNA)	0.32 ± 0.07	0.23 ± 0.03	5.47 ± 0.84	0.53 ± 0.15	2.82 ± 0.22	6.22 ± 0.50	0.75 ± 0.15	5.85 ± 0.94
	23S rRNA (theoretical)	2.62	1.31	0	1.10	1.10	1.57	3.30	16.60

	16S rRNA (theoretical)	0	0	5.14	0	2.06	5.68	0	3.17
2h	tRNA	1.69 ± 0.20	NA	0.44 ± 0.10	7.46 ± 0.30	25.55 ± 0.56	0.34 ± 0.08	97.14 ± 1.71	136.29 ± 4.04

Dynamics of mRNA modification

In eukaryotes, mRNA methylation is a dynamic, reversible process that has regulatory functions and involves a complex system of writers, readers, and erasers (8) (22). Therefore, we were interested in whether the newly identified modifications of bacterial mRNA change in relative abundance depending on the growth phase. We compared the modification patterns between the early exponential growth phase (2 h) and the stationary phase (8 h) (Fig. 3A). We observed statistically significant increases in m⁶A, m¹G, m⁵C, and Ψ, and a significant decrease in m^{6,6}A in some of the mRNA-enriched fractions of stationary phase cells compared with mRNA derived from the exponential growth phase cells. Whereas for m⁶A the increase of modification abundance towards the stationary phase was observed in fractions 2 and 4, for m¹G and Ψ, we observed an increase only in fraction 2, and for m⁵C only in fraction 4.

In contrast, the abundances of rRNA modifications in *E. coli* between exponential and stationary phase cells were mostly unchanged, as shown by the semi-quantitative analysis of fraction 1 and fraction 3 (Fig. 3A). We only measured a small increase in the abundance of m⁶A, m¹G, and Ψ in the 16S rRNA-containing fraction 3 of stationary phase cells. However, for example, m⁶A is not known to exist in 16S rRNA, and the changes in the measured amounts are similar to the changes observed for the mRNA-enriched fractions 2 and 4. Therefore, it is likely that these results hail from the co-purification of mRNAs of the same size.

As the abundances of modifications in rRNA were comparable between 2 h and 8 h of cultivation, we decided to analyze the mRNA+rRNA fractions obtained by size exclusion chromatography to study growth phase-dependent changes in mRNA modifications (Supplemental Fig. S6). For m⁶A and m¹G, we measured a steady increase, with the highest values observed after 24 hours of growth (representing the late stationary phase). In contrast, for m⁵C, we observed a slight decrease in abundance over the course of growth of *E. coli*. For m⁷G, the abundance remained nearly unchanged over the measured time course, but there was a temporal peak

during the transition of *E. coli* to stationary phase. Overall, the quantitative changes observed in the mRNA-enriched gel preparations were also found in the m+rRNA fractions. Thus, quantitative analysis of the m+rRNA fraction could be an interesting approach for future studies to determine mRNA modification profiles in other bacteria under different environmental stress conditions.

Next, we tested whether the modification profile of mRNA in *E. coli* changes under the influence of external factors such as pH or oxygen availability. We cultivated cells in tryptone-based (LB) medium at pH 7.6 or buffered at pH 5.4 under aerobic or microaerobic (oxygen limited) conditions, purified mRNA using denaturing gel electrophoresis, and compared the absolute abundance of mRNA modifications during the exponential growth phase and shortly before the transition into the stationary phase. We found that most of the epitranscriptomic marks remained at similar abundances independent of oxygen limitation: Changes of the mRNA modifications observed during the transition from exponential growth to stationary phase under aerobic conditions (Fig. 3A) were also found under oxygen limiting conditions (Supplemental Fig. S7). Specifically, we observed a statistically significant increase in m⁶A, m¹G, and m⁵U and a decrease in m⁵C in the long mRNA-enriched fraction 4 of stationary phase cells both under aerobic and microaerobic conditions.

However, we noted a sharp decrease for the m^{6,6}A levels in 16S rRNA upon oxygen limitation. During aerobic growth the m^{6,6}A abundance in fraction 3 (16S rRNA) was 5.8‰ ± 0.35‰ (Fig. 3A), and under microaerobic conditions there was a more than threefold reduction to 1.51‰ ± 0.35‰ (Fig. 4). These results indicate a correlation between environmental stress adaptation and rRNA modification in *E. coli*.

Comparison of cells cultivated under physiological condition (pH 7.6) and acid stress (pH 5.4) revealed stress-specific changes for two modifications: m^{6,6}A and Ψ (Fig. 4). The cultivation of *E. coli* at pH 5.4 led to a threefold reduction in the m^{6,6}A abundance in fraction 3 (16S rRNA) and to a smaller extent in fraction 2 (long mRNAs). An even more dramatic effect was observed for Ψ. Cultivation of *E. coli* under aerobic or microaerobic conditions at pH 5.4

resulted in a loss of Ψ in fraction 4 (short mRNAs) (Fig. 4). This effect was target-specific and not caused by a general issue with RNA as the levels of other modifications in this fraction were not affected (data not shown). In addition, the Ψ level of fraction 2 (long-mRNAs) was not

significantly altered by acid stress. These results suggest repression of specific subsets of mRNAs between 400 and 1,200 nt under acid stress, which are otherwise targets of Ψ modification.




Figure 4. Impact of oxygen limitation and acid stress on the RNA modifications in *E. coli*. Changes in the absolute levels of modifications in mRNA and rRNA-enriched fractions extracted after denaturing gel electrophoresis of total RNA from *E. coli* MG11655 cells cultivated in complex LB media buffered at pH 7 or pH 5 under aerobic or microaerobic conditions. 1, 23S rRNA; 2, RNAs 1,700 to 2,700 nts; 3, 16S rRNA; 4, RNAs 400 to 1,100 nts. The results of three biological replicates are presented. p-values were calculated by paired Student's t-test with * ≤ 0.05 , ** ≤ 0.01 , and *** ≤ 0.001 .

DISCUSSION

In this study, we analyzed the mRNA modification profile of the model bacterium *E. coli* MG1655. In addition to the known modification m⁶A, we provide evidence for the presence of previously undescribed modifications in the mRNA of *E. coli*. Using two different mRNA enrichment methods in combination with targeted LC-MS/MS, we obtained evidence that modifications previously

found only in the rRNA and/or tRNA of *E. coli*, such as m⁶A, m^{6,6}A, m¹G, m⁵C, m⁷G, m⁵U, and Ψ also occur as mRNA modifications. m⁵C, m⁷G, and Ψ have been previously described as mRNA modifications in eukaryotes, and our quantitative MS analysis revealed comparable levels of these modifications in bacterial mRNA. For example, m⁵C has been found in the mRNA of the archaeon *S. solfataricus* (14) and in human mRNA with an abundance of 4.3‰ (23), which is similar to the abundances of 3.25‰ \pm

0.84‰ (fraction 2, long mRNA) and 3.65‰ ± 0.33‰ (fraction 4, short mRNA) measured in this study (Fig. 3A). Likewise, the determined abundance of Ψ in the mRNA of *E. coli* (1%) was comparable with the levels of 0.2%–0.6% that have been reported for human mRNA (24), and the abundance of m⁷G in the mRNA of *E. coli* (2‰–3‰) was comparable to the abundance of 4.5‰ of m⁷G found in human mRNA (25). In mitochondria, which have 70S ribosomes as prokaryotes, Ψ is also found in the mRNA (26). A previous screen found no evidence for internal m⁷G modifications in *E. coli*'s mRNA (27), suggesting that our result might reflect a capping modification, similar to the m⁷G cap at the 5' end of mRNA in eukaryotes (28). On the other hand, we obtained evidence that m^{6,6}A and m⁵U are present in the mRNA of a prokaryote. Hence, our work implies a high complexity of mRNA modifications in *E. coli*, which resembles the profiles previously observed in eukaryotic species (7) (4). Future studies should focus on identifying the enzymes involved in bacterial mRNA modification and on clarifying whether certain mRNA structures are also substrate of the thus far known RNA methyltransferases.

We also show that the abundance of modifications such as m⁶A, m¹G, m⁵C, and m⁵U in mRNA changes during the growth of *E. coli*. Moreover, we observed clear effects resulting from acid stress on the quantity of Ψ in short mRNAs from *E. coli*. Since the quantity of Ψ in the long mRNAs was not changed, it is suggested that a set of genes, which is modified under normal conditions, is repressed under acid stress. Variation in mRNA modification levels during different stages of development has been previously observed in mammalian cells, and disruption of the plasticity of RNA modification leads to severe defects in tissue development and homeostasis (6). Additionally, the presence of Ψ in mRNA has been shown to increase the mRNA half-life in the parasite *Toxoplasma gondii* (24).

A potential regulatory function of the *E. coli* mRNA methylation status is supported by the observed different abundances of modifications between long and short mRNA-enriched fractions, which suggest a target-specific RNA methylation. If a modification is universal and is present in all mRNAs in equal amounts per molecule, such as some capping modifications, its abundance would be “diluted” and would thus show a lower amount in longer mRNAs. However, we observed the opposite scenario, with a higher abundance of m⁶A and m¹G in long mRNAs compared to short mRNAs, which indicates that these may be gene-specific

modifications that could affect target-specific processes within the cell.

Previously, analysis of the modification profile of bacterial mRNA has been limited by the lack of reliable methods for its purification and quantification. In this work, we validated a simple and low-cost method with the potential to overcome some of the existing difficulties. One of the main advantages of our gel extraction approach is the direct, simultaneous separation of different RNA subtypes, which allows a comparison of RNA fractions that underwent identical chemical treatment. Hence, it avoids possible biases due to the usage of different separation methods for each RNA type, primer selection, or low antibody specificity. Yet, some of the challenges for bacterial mRNA purification remain, and the contamination of mRNA fractions with rRNA cannot be excluded and is thus the major limitation of our purification platform.

The separation of RNA in different fractions gave us strong evidence that Ψ is present in mRNA, when we consider the drastic reduction of Ψ levels only in the short mRNA-enriched fraction 4 under acid stress. Since Ψ is well distributed along the sequences of both 23S and 16S rRNA, the unchanged levels of other modifications in fraction 4 together with the unchanged Ψ levels in the other three fractions confirmed that under this condition the absence of Ψ is not a result of changes in the rRNA modification levels or different levels of rRNA-contamination in this fraction. Creating modification profiles by measuring several modifications at once is useful also for assessing potential fraction-to-fraction contamination with tRNA. For examples, the m^{6,6}A:m⁵C:m⁵U:m⁷G: Ψ ratio of tRNA is substantially different to the ratio in mRNA-enriched fractions which might serve as an indication of low to no tRNA contamination in the mRNA-enriched fractions.

Size-dependent RNA separation allows for the comparison of modification abundances between the fractions enriched in long or short mRNA, which can indicate whether modifications are specific or widespread. In the future, owing to the size selectivity of mRNA purification, the gel extraction approach could also be combined not only with MS but also with third-generation RNA sequencing methods to enrich certain target fractions of interest and to determine the location of observed modifications in transcripts.

Taken together, our results show that mRNA modification is not a specific regulatory feature of eukaryotes but also appears to have a regulatory function in prokaryotes. In bacteria, which exhibit rapid mRNA turnover and no

compartmentalization, alteration of subsets of functionally related mRNAs with different modifications that specifically affect mRNA stability or translation may be an efficient way to prioritize certain cellular programs.

MATERIAL AND METHODS

Strains and growth conditions – In this study, we used *E. coli* K-12 MG1655 and *E. coli* BW25113 $\Delta trmN$ carrying chromosomal deletion of *trmN* (29). All strains were grown overnight in LB medium (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl, pH 7.6). Cells from the overnight culture were used to inoculate fresh LB medium or LM-MES medium (LB plus 0.1 M MES, pH 5.4). Bacteria were grown under agitation (200 rpm) at 37°C, and growth was monitored over time by measuring the optical density at 600 nm (OD₆₀₀). For microaerobic conditions, the cells were cultivated without shaking in closed, full vessels without excess air (Schott Duran bottles GL45, Jena, Germany).

Total RNA purification – RNA was isolated using the PCI protocol (30) with modifications. Each bacterial pellet was washed in 1 mL of ice-cold AE buffer (20 mM sodium acetate buffer, pH 5.2, 1 mM EDTA) and was resuspended in 500 μ L of the same buffer. Then, 500 μ L of pre-warmed PCI for RNA extraction (Carl Roth, Karlsruhe, Germany) and 25 μ L of 10% (w/v) SDS were added, and the mixture was incubated for 5 min at 60°C under vigorous agitation. Samples were cooled for 2 h on ice and centrifuged for 1 h at 16,000 $\times g$. The supernatant was transferred to phase-lock tubes (Phase Lock Gel, QuantaBio, USA), and 1.0 volume of PCI and 0.1 volume of 3 M sodium acetate (pH 5.2) were added before centrifugation for 15 min. The supernatant was collected, mixed with 2.3 volumes of ethanol, and placed in a freezer at –80°C overnight. After centrifugation at 16,000 $\times g$ for 1 h, the supernatant was discarded, and the pellet was washed twice with 70% (v/v) ethanol, dried, and resuspended in 100 μ L of RNase-free water. After treatment of the samples with RNase-free DNase (Roche) according to the manufacturer's protocol, the samples were purified with another round of precipitation. The integrity of RNA was assessed via chip gel electrophoresis (Bioanalyzer 2100, RNA Nano chip kit, Agilent, Waldbronn). To evaluate the extent of remaining buffer and DNA contaminations, the RNA preparation samples were tested by performing standard spectroscopic measurements (Nanodrop One)

as well as with RT-qPCR analysis, where the cDNA synthesis step was not applied.

RNA separation with size exclusion chromatography – The separation of tRNA by size exclusion chromatography (SEC) was performed as described previously (31) with minor modifications. Briefly, total RNA was loaded on a Thermo Scientific UltiMate 3000 LC system equipped with a diode array detector set to 260 nm, an autosampler, a column thermostat (60°C), and a fraction collector. A size exclusion column (Agilent Bio SEC-3, 3 μ m, 300 Å, 7.8 \times 300 mm, Agilent, Waldbronn) allowed the collection of the RNA fractions after isocratic elution with 100 mM ammonium acetate at pH 7. The peaks representing the mRNA plus rRNA fraction as well as the tRNA fraction were collected and concentrated in a vacuum concentrator (Eppendorf Concentrator 5301). 5 M NH₄OAc was added to a final concentration of 0.5 M, and after addition of 2 \times Vol. ice-cold ethanol (100%), the RNA was precipitated overnight at –20°C. After centrifugation at 12,000 $\times g$ for 30 min at 4°C, the RNA pellet was subjected to an additional ethanol (80%, v/v) wash step to verify the complete removal of the ammonium acetate and was then resuspended in pure water. The quality of the isolated tRNA was verified with chip gel electrophoresis (BioAnalyzer 2100, RNA Pico chip, Agilent, Waldbronn), RT-qPCR analysis, agarose gel electrophoresis, and usage of deletion mutants lacking specific RNA modifications on either the tRNA or the mRNA + rRNA fractions. RNA concentration was determined by NanoDrop ND1000 spectrophotometer (peqlab, Germany).

Gel electrophoresis and RNA extraction – For gel electrophoretic extraction of RNA, denaturing 1.2% (w/v) agarose gels (agarose – Serva, Germany), 20 mM MOPS, 1.1% formaldehyde dissolved in DEPC-treated water) were used. Before loading 10 μ g of RNA, each sample was mixed 1:1 with sample buffer (64% [v/v] desalted formamide, 8.35% [v/v] formaldehyde, 26 mM MOPS, and 0.05% [v/v] ethidium bromide). After the addition of 1/10 volume RNA-marker (50% [v/v] glycerol, 1 mM EDTA, bromophenol blue, and xylene cyanol), the samples were denatured for 5 min at 65°C, followed by immediate cooling on ice. The gels were run in running chambers (Bio-Rad) using 20 mM MOPS as a running buffer at 5 V/cm for 3 h. As a size reference, the RiboRuler high range RNA ladder (Thermo Scientific) was used. RNA fragments with sizes from 3,200 to 2,800 nt (23S rRNA), 2,700 to 1,700 nt (long mRNAs), 1,600 to 1,400 nt (16S rRNA), and 1,100 to 400

nt (short mRNAs) were cut from the gel, leaving physical space between each of them. Cutting was documented before and after the excision of the fragments. RNA was isolated from the gel fragments with a commercially available kit (Hi Yield Gel/PCR DNA Fragment Extraction Kits, Suedlabor Gauting, Germany). Elution from the column was performed in two consecutive steps using 25 μ L of water. Finally, the quality of each fraction was analyzed using chip gel electrophoresis (Bioanalyzer 2100, RNA Pico chip kit, Agilent, Waldbronn) or by reloading of the isolated RNA on a denaturing agarose gel. Finally, the efficiency of depletion was tested using Bioanalyzer chip electrophoresis and RT-qPCR.

Oligonucleotide-based depletion of rRNA –

After removal of tRNA and small noncoding RNAs by size exclusion chromatography, the remaining mRNA plus rRNA mixture was subjected to two consecutive rounds of rRNA depletion according to the manufacturer protocols for oligonucleotide-based depletion of rRNA (RiboPOOL oligo pool for *E. coli*, siTOOLS, Martinsried, Germany). Finally, the efficiency of depletion was tested by Bioanalyzer chip electrophoresis and RT-qPCR. For the comparison of modification abundancies before and after rRNA-depletion with oligonucleotide-based kit presented in Fig. S5 we used the RiboZero RNA Removal Kit, Bacteria (Illumina, USA), which was unfortunately discontinued in the course of this work and was therefore replaced for later studies with the RiboPOOL oligo pool for *E. coli* (siTOOLS, Germany).

MS analysis of RNA methylation – Qualitative (no internal standard, no calibration), semi-quantitative (internal standard, no calibration) and absolute quantitative analyses (internal standard and calibration) of the prepared RNA fractions were performed by LC-MS/MS as previously described (21). After the enzymatic hydrolysis of the RNA to nucleosides using a mixture of benzonase, snake venom phosphodiesterase and calf intestine phosphatase (21), the improved gen² ¹³C/¹⁵N stable isotope labeled internal standard (SILIS) from *S. cerevisiae* tRNA was added (32) for semi-quantitative and absolute quantitative analysis. The resulting ribonucleoside mixture was separated using a Synergy Fusion RP column, with 2.5- μ m particle size, 100-Å pore size, 100-mm length, and 2-mm inner diameter from Phenomenex (Torrance, CA, USA), on an Agilent 1290 Infinity II series UHPLC. Mobile phase A was 5 mM ammonium acetate adjusted to pH 5.3 with glacial acetic acid, and

mobile phase B was pure acetonitrile. Gradient elution started with 100% A for 1 min, increased to 10% B after 4 min, 40% after 7 min, maintained for 1 min and re-establishment of the starting conditions with 100% A for an additional 2.5 min. The flow rate was 0.35 mL/min, and the column temperature was 35°C. For MS measurements, an Agilent 6470 Triple Quadrupole mass spectrometer set to dynamic multiple reaction monitoring mode was used. The MS was operated in positive ion mode with the following parameters: skimmer voltage of 15 V, cell accelerator voltage of 5 V, N₂ gas temperature of 230 °C and N₂ gas flow of 6 L/min, sheath gas (N₂) temperature of 400 °C with a flow of 12 L/min, capillary voltage of 2500 V, nozzle voltage of 0 V, and nebulizer at 40 psi.

RT-qPCR analysis – For RT-qPCR analysis, equal amounts of the isolated total RNA or purified RNA were converted to cDNA with the iScript Advanced Script (Bio-Rad) according to the manufacturer's protocol. The samples were mixed with SsoAdvanced Univ SYBR Green Supermix (Bio-Rad), dispensed in triplicate onto a 96-well PCR plate (Bio-Rad), and subjected to qPCR in a Bio-Rad CFX real-time cycler (see table S1 for primers). The same amount of starting RNA was used for cDNA synthesis. For additional internal calibration, we evaluated the data according to the $\Delta\Delta C_t$ method using 16S rRNA and/or mRNA targets such *recA* or *alaS* as internal references (33).

Analysis of mRNA enrichment by RT-qPCR

– To evaluate the level of RNA separation and mRNA enrichment in the mRNA fractions extracted after denaturing gel electrophoresis (fractions 2 and 4) and after oligonucleotide-based rRNA depletion, we evaluated the mRNA:16S rRNA as well as the mRNA:23S rRNA ratios in the samples by RT-qPCR. To calculate the level of rRNA removal, we compared changes in the mRNA:rRNA ratios before (total RNA or the mRNA + rRNA fraction) and after gel electrophoresis or oligonucleotide-based rRNA depletion (mRNA fractions). To avoid potential errors due to imprecise determination of the RNA concentrations of the total RNA and mRNA samples, which could affect the results if directly comparing 16S rRNA with 23S rRNA, we used the $\Delta\Delta C_t$ method, taking 16S and 23S rRNA as “genes” of interest and mRNA targets as reference genes for internal calibration (the reverse scenario of a standard calculation of gene expression). We used the *alaS* gene (2,631 bp) as an internal reference for the long mRNA fractions and the *recA* gene (1,062 bp) for the short mRNA fractions or the mRNA after oligo-

based rRNA depletion with RiboPOOL. Additionally, as a control for the separation of RNA by gel electrophoresis, we tested the amounts of *recA* and *alaS* in the fractions in which these mRNAs should not be expected based on their size-long mRNA and short mRNA, respectively. Indeed, the higher C_T values for *alaS* in the short mRNA fraction compared to that in the long mRNA fractions and the higher C_T values for *recA* in the long mRNA fractions than those in the short mRNA fractions indicate that after gel separation, *recA* was predominantly in the short mRNA fraction and *alaS* was in the long mRNA fraction.

To calculate the mRNA:rRNA ratios as percentages, we used the fold-change in the ratio of *alaS* (long mRNA) or *recA* (short mRNA) versus 23S rRNA or 16S rRNA between the total RNA and mRNA-containing fractions 2 and 4, on the basis that 90% rRNA (45% 23S rRNA, 45% 16S rRNA) and 1% mRNA are present in the total RNA sample (34). Therefore, we have 1:45 ratios of mRNA:23S rRNA and mRNA:16S rRNA. Then, the calculated fold-changes for the *recA*:16S rRNA and *alaS*:23S rRNA ratios between the total RNA samples and the gel-extracted mRNA-enriched fractions were used to calculate the final ratio of mRNA:rRNA in the mRNA-enriched fractions based on the initial estimated ratio of 1:45. For example, at a fold-change of 90 between *alaS*:23S rRNA, we would have a mRNA:23S rRNA ratio of 2:1, and similarly, for a fold-change of 90 between *recA*:16S rRNA, we would have a mRNA:16S rRNA ratio of 2:1. Taken together, this would mean that we have a ratio of mRNA:23S rRNA:16S rRNA of 2:1:1 or a total of 50% mRNA and 50% rRNA.

Analysis of mRNA enrichment by RNA sequencing – To evaluate the level of contaminating rRNA degradation products in the mRNA-enriched fractions 2 and 4 from our denaturing agarose gel electrophoresis experiments, we converted the RNA from all gel extracted fractions into cDNA libraries using the NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs, USA), involving fragmentation of the RNA into 200 nt fragments. As a control, we also prepared libraries from the corresponding total RNA. The libraries were analyzed in an HiSeq Illumina sequencer with an average sequencing depth of ~8 million single-end 50 nt-long reads.

Data analysis was performed with the CLC Genomic workbench software (Qiagen, Germany) with or without duplicate removal and both data sets were analyzed independently. To evaluate the approximate molar ratios of each RNA type in the samples, the quotient of (reads for a specific gene) / (total reads) was divided by the length of the specific gene. For the calculation of potential 23S rRNA and 16S rRNA contamination in the gel-extracted mRNA-enriched fractions 2 and 4, we used the percentage of reads binding to 23S rRNA or 16S rRNA, and multiplied the respective percentage with the abundance of each modification found in the 23S rRNA-enriched or 16S-enriched fraction. Because the rRNA sequences had rather uniform coverage in the total RNA samples, but there were differences in the coverage of rRNA sequences in the gel-extracted fractions 2 and 4, the calculations included the enrichment of distinct fragments of rRNA. To determine which fragments of the 23S rRNA and 16S rRNA sequences were over- or underrepresented, the results were compared with the coverage in the total RNA samples. From the number of reads covering these fragments, the percentage of rRNA contamination for each fraction (or the individual modifications within that fraction) was calculated.

ACKNOWLEDGEMENT

We are thankful to Florian Schelter, Matthias Hei and Thomas Carell, Department of Chemistry at LMU Munich, for the MS-data presented in Fig. 3 and Supplemental Fig. S1.

FUNDING

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation): SFB1309, project number 325871075 (S.K.) and JU270/21-1, project number 464582101 (K.J.).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

- Kadumuri, R.V. and Janga, S.C. (2018) Epitranscriptomic code and its alterations in human disease. *Trends Mol Med*, **24**, 886-903.
- Boccalletto, P., Machnicka, M.A., Purta, E., Piatkowski, P., Baginski, B., Wirecki, T.K., de Crécy-Lagard, V., Ross, R., Limbach, P.A., Kotter, A. *et al.* (2018) MODOMICS: a database of RNA modification pathways. 2017 update. *Nucleic Acids Res*, **46**, D303-d307.
- Li, S. and Mason, C.E. (2014) The pivotal regulatory landscape of RNA modifications. *Annu Rev Genomics Hum Genet*, **15**, 127-150.
- Roundtree, I.A., Evans, M.E., Pan, T. and He, C. (2017) Dynamic RNA Modifications in Gene Expression Regulation. *Cell*, **169**, 1187-1200.
- Wang, X. and He, C. (2014) Dynamic RNA modifications in posttranscriptional regulation. *Mol Cell*, **56**, 5-12.
- Frye, M., Harada, B.T., Behm, M. and He, C. (2018) RNA modifications modulate gene expression during development. *Science*, **361**, 1346-1349.
- Gilbert, W.V., Bell, T.A. and Schaening, C. (2016) Messenger RNA modifications: Form, distribution, and function. *Science*, **352**, 1408-1412.
- Zaccara, S., Ries, R.J. and Jaffrey, S.R. (2019) Reading, writing and erasing mRNA methylation. *Nat Rev Mol Cell Biol*, **20**, 608-624.
- Eyler, D.E., Franco, M.K., Batool, Z., Wu, M.Z., Dubuke, M.L., Dobosz-Bartoszek, M., Jones, J.D., Polikanov, Y.S., Roy, B. and Koutmou, K.S. (2019) Pseudouridylation of mRNA coding sequences alters translation. *Proc Natl Acad Sci U S A*, **116**, 23068-23074.
- Sergeeva, O.V., Bogdanov, A.A. and Sergiev, P.V. (2015) What do we know about ribosomal RNA methylation in *Escherichia coli*? *Biochimie*, **117**, 110-118.
- Nachtergaele, S. and He, C. (2017) The emerging biology of RNA post-transcriptional modifications. *RNA Biol*, **14**, 156-163.
- Rauhut, R. and Klug, G. (1999) mRNA degradation in bacteria. *FEMS Microbiol Rev*, **23**, 353-370.
- Deng, X., Chen, K., Luo, G.Z., Weng, X., Ji, Q., Zhou, T. and He, C. (2015) Widespread occurrence of N6-methyladenosine in bacterial mRNA. *Nucleic Acid Res*, **43**, 6557-6567.
- Edelheit, S., Schwartz, S., Mumbach, M.R., Wurtzel, O. and Sorek, R. (2013) Transcriptome-wide mapping of 5-methylcytidine RNA modifications in bacteria, archaea, and yeast reveals m5C within archaeal mRNAs. *PLoS Genet*, **9**, e1003602.
- Motorin, Y. and Helm, M. (2019) Methods for RNA Modification Mapping Using Deep Sequencing: Established and New Emerging Technologies. *Genes*, **10**.
- Hagelskamp, F., Borland, K., Ramos, J., Hendrick, A.G., Fu, D. and Kellner, S. (2020) Broadly applicable oligonucleotide mass spectrometry for the analysis of RNA writers and erasers in vitro. *Nucleic Acids Res*, **48**, e41.
- Golovina, A.Y., Sergiev, P.V., Golovin, A.V., Serebryakova, M.V., Demina, I., Govorun, V.M. and Dontsova, O.A. (2009) The *yfiC* gene of *E. coli* encodes an adenine-N6 methyltransferase that specifically modifies A37 of tRNA^{Val}(cmo⁵UAC). *RNA*, **15**, 1134-1141.
- Kaiser, S., Byrne, S.R., Ammann, G., Asadi Atoi, P., Borland, K., Brecheisen, R., DeMott, M.S., Gehrke, T., Hagelskamp, F., Heiss, M. *et al.* (2021) Strategies to Avoid Artifacts in Mass Spectrometry-Based Epitranscriptome Analyses. *Angew Chem Int Ed*, **60**, 23885-23893.
- Legrand, C., Tuorto, F., Hartmann, M., Liebers, R., Jacob, D., Helm, M. and Lyko, F. (2017) Statistically robust methylation calling for whole-transcriptome bisulfite sequencing reveals distinct methylation patterns for mouse RNAs. *Genome Res*, **27**, 1589-1596.
- Sergiev, P.V., Aleksashin, N.A., Chugunova, A.A., Polikanov, Y.S. and Dontsova, O.A. (2018) Structural and evolutionary insights into ribosomal RNA methylation. *Nat Chem Biol*, **14**, 226-235.
- Borland, K., Diesend, J., Ito-Kureha, T., Heissmeyer, V., Hammann, C., Buck, A.H., Michalak, S. and Kellner, S. (2019) Production and Application of Stable Isotope-Labeled Internal Standards for RNA Modification Analysis. *Genes*, **10**, 26.
- Esteve-Puig, R., Bueno-Costa, A. and Esteller, M. (2020) Writers, readers and erasers of RNA modifications in cancer. *Cancer Lett*, **474**, 127-137.
- Squires, J.E., Patel, H.R., Nusch, M., Sibbritt, T., Humphreys, D.T., Parker, B.J., Suter, C.M. and Preiss, T. (2012) Widespread occurrence of 5-methylcytosine in human coding and non-coding RNA. *Nucleic Acids Res*, **40**, 5023-5033.
- Li, X., Zhu, P., Ma, S., Song, J., Bai, J., Sun, F. and Yi, C. (2015) Chemical pulldown reveals dynamic pseudouridylation of the

- mammalian transcriptome. *Nat Chem Biol*, **11**, 592-597.
25. Zhang, L.S., Liu, C., Ma, H., Dai, Q., Sun, H.L., Luo, G., Zhang, Z., Zhang, L., Hu, L., Dong, X. *et al.* (2019) Transcriptome-wide mapping of internal N⁷-methylguanosine methylome in mammalian mRNA. *Mol Cell*, **74**, 1304-1316.e1308.
 26. Antonicka, H., Choquet, K., Lin, Z.Y., Gingras, A.C., Kleinman, C.L. and Shoubbridge, E.A. (2017) A pseudouridine synthase module is essential for mitochondrial protein synthesis and cell viability. *EMBO Rep* **18**, 28-38.
 27. Enroth, C., Poulsen, L.D., Iversen, S., Kirpekar, F., Albrechtsen, A. and Vinther, J. (2019) Detection of internal N⁷-methylguanosine (m⁷G) RNA modifications by mutational profiling sequencing. *Nucleic Acids Res*, **47**, e126.
 28. Furuichi, Y. (2015) Discovery of m⁷G-cap in eukaryotic mRNAs. *Proc Jpn Acad Ser B Phys Biol Sci*, **91**, 394-409.
 29. Grenier F, Matteau D, Baby V, Rodrigue S. (2014) Complete genome sequence of *Escherichia coli* BW25113. *Genome Announc*. **2**, 01038-14.
 30. Sambrook, J. and Russell, D.W. (2006) Purification of nucleic acids by extraction with phenol:chloroform. *Cold Spring Harb Protoc*, **2006**.
 31. Heiss, M., Reichle, V.F. and Kellner, S. (2017) Observing the fate of tRNA and its modifications by nucleic acid isotope labeling mass spectrometry: NAIL-MS. *RNA Biol*, **14**, 1260-1268.
 32. Heiss, M., Borland, K., Yoluç, Y. and Kellner, S. (2021) Quantification of Modified Nucleosides in the Context of NAIL-MS. *Methods Mol Biol*. **2298**, 279-306.
 33. Schmittgen, T.D. and Livak, K.J. (2008) Analyzing real-time PCR data by the comparative C_T method. *Nat Protoc*, **3**, 1101-1108.
 34. Westermann, A.J., Gorski, S.A. and Vogel, J. (2012) Dual RNA-seq of pathogen and host. *Nat Rev Microbiol*, **10**, 618-630.