
A NEW METHOD FOR DETERMINING STOICHIOMETRIC COEFFICIENTS USING MINORS OF A MATRIX

Rainier Lombaard

Email: rlombaard.ac@gmail.com


Abstract

Spinel materials often have complex structures and as a result, balancing of reactions with these compounds by traditional methods become very time consuming. A method to calculate the stoichiometric coefficients for chemical reactions using first a modified matrix-inverse method and then an optimised method is proposed. Both methods are explored using linear algebra and the result demonstrated using a typical chromite reduction reaction.

Introduction

One of the first concepts that students of chemistry learn is that of conservation of mass. Physical changes that demonstrate that mass is conserved during phase change are typically demonstrated, after which chemical reactions are introduced. In a chemical reaction the mass and elements are still conserved but the initial substances, reactants, will be transformed into new substances, products. In pioneering work done during the late 1700's, Antoine Lavoisier showed conclusively that the law of conservation of mass also applies to chemical reactions with the publication of his *Traité élémentaire de chimie* "In all the operations of art and nature, nothing is created; an equal quantity of matter exists both before and after the experiment; the quality and quantity of the elements remain precisely the same." (Lavoisier, 1789).

It was then understood that since the quantities of each element cannot change, even though they would form part of different substances, the amounts of reactants and products must be scaled by integer coefficients for the law of conservation of mass to hold (IUPAC, 1997). This insight led to the development of stoichiometry which studies the relationship between the amounts of reactants consumed and products produced in chemical reactions (Brown *et al*, 2015). As demonstration, consider the hypothetical single displacement reaction below:




The stoichiometric coefficients x_1, x_2, x_3 and x_4 must be some integers such that the number of atoms of elements A , B and C are the same on both sides of the arrow. The dominant method taught at high school and even university level to find the values of the coefficients is solving by inspection (Toth, 1997). This casual approach works well with simple chemical equations such as the one shown above. But for even slightly more complex reactions involving larger quantities of reactants and products, this inspection method becomes slow and unreliable.

Most students of chemistry would also be familiar with linear algebra and its techniques. To demonstrate some of the applications of linear algebra, systems of linear equations are introduced and then applied to chemical equations. This algebraic method of using a matrix of coefficients results in a much more systematic approach to balancing chemical equations and with the advent of modern scientific calculators that can store and manipulate matrices, has become the standard. However, this system of equations method becomes cumbersome to work with for large reaction equations with many substances and elements. Therefore, it is the goal is to develop a general method that can be applied to all chemical equations with ease.

Discussion

This project came about during the study of metallothermic reduction of metal oxide ores. These ores typically consist of complicated mineral structures which, when undergoing reduction, become very burdensome to balance by hand. Consider the following chromite molecule and its subsequent reduction with aluminium:



It is readily apparent that solving the following by inspection would be a very time-consuming task. However, below we outline a new augmented matrix method to calculate the coefficients using a simple spreadsheet application with ease.

Methodology


The steps that can be followed to solve the stoichiometric coefficients are outlined below:

- 1) Construct a composition array for the reaction
- 2) From this array extract the coefficient matrix K_{ij}
- 3) Construct the augmented matrix A_{nm} according to the following rules:
 - If $n \leq i$ and $m \leq j$ then $a_{nm} = c_{ij}$
 - If $n > i$ and $m \leq j$ insert the reversal matrix J_{aj}
 - If $n > i$ and $m > j$ insert the null matrix 0_{ad}
 - If $n = i$ and $m > j$ insert the matrix $H_{1d} \begin{bmatrix} 1 & 0 & \cdots & 0 \end{bmatrix}$
 - If $n \leq i - 1$ and $m > j$ insert reversal matrix $J_{(i-1)d}$ with first column of zeros
 - Where $k = \text{Max}(i, j) - \text{Min}(i, j)$ then $a = i + k + 2$ and $d = j + k + 2$
- 4) Extract the final coefficient matrix B_{xy} such that $x = y$ according to the rules:
 - If $i < j$ then $x = i + k$ and $y = j$ or,
 - If $i > j$ then $x = i + 1$ and $y = j + k + 1$ or,
 - If $i = j$ then $x = i + 1$ and $y = j + 1$
- 5) Determine the inverse matrix B_{xy}^{-1} as well as the determinant $\det(B_{xy})$
- 6) Transpose the last column of B_{xy}^{-1} into a new matrix C_{1n}
- 7) Multiply C_{1n} by the absolute value of the determinant e.g. $|\det(B_{xy})| \cdot C_{1n} = D_{1n}$
- 8) Divide each element of D_{1n} the greatest common divisor e.g. $D_{1n} / \text{gcd}(d_{11}, \dots, d_{1n}) = E_{1n}$
- 9) The resulting matrix E_{1n} now contains the stoichiometric coefficients from left to right.

The above method may seem needlessly complex, but the repeatability will become readily apparent. In the final section of the paper an optimised method for determining the stoichiometric coefficients is revealed. First, we will demonstrate the above methodology using a simple double displacement reaction:

The reaction of iron(III) oxide with sulfuric acid

This reaction can be represented with coefficients as follows:



Step 1:

Construct a composition array for the reaction:

	Fe_2O_3	H_2SO_4	$Fe_2(SO_4)_3$	H_2O
Fe	2	0	-2	0
O	3	4	-12	-1
H	0	2	0	-2
S	0	1	-3	0

Note that the coefficients for the products are all assigned the negative value. This is a direct consequence of the law of conservation of mass since the quantity of any element needs to add to zero on both sides of the chemical equation. This is usually the method employed when using a system of linear equations to determine the coefficients (*Gabriel et. al, 2015*).

Step 2:

From this array we then extract the matrix K :

$$K = \begin{bmatrix} 2 & 0 & -2 & 0 \\ 3 & 4 & -12 & -1 \\ 0 & 2 & 0 & -2 \\ 0 & 1 & -3 & 0 \end{bmatrix}$$

Step 3:

Construct the augmented matrix A:

$$i = j = 4$$

$$\therefore a = d = 6$$

$$A = \begin{bmatrix} 2 & 0 & -2 & 0 & 0 & 0 \\ 3 & 4 & -12 & -1 & 0 & 1 \\ 0 & 2 & 0 & -2 & 0 & 0 \\ 0 & 1 & -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

Step 4:

Extract the coefficient matrix B:

$$i = j = 4$$

$$\therefore x = y = 5$$

$$B = \begin{bmatrix} 2 & 0 & -2 & 0 & 0 \\ 3 & 4 & -12 & -1 & 0 \\ 0 & 2 & 0 & -2 & 0 \\ 0 & 1 & -3 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

Step 5:

Determine the inverse matrix B^{-1} as well as the determinant:

$$B^{-1} = \begin{bmatrix} \frac{2}{3} & -\frac{1}{9} & \frac{2}{9} & 0 & \frac{1}{3} \\ 0 & 0 & \frac{1}{2} & 0 & 1 \\ \frac{1}{6} & -\frac{1}{9} & \frac{2}{9} & 0 & \frac{1}{3} \\ 0 & 0 & 0 & 0 & 1 \\ \frac{1}{2} & -\frac{1}{3} & \frac{1}{6} & 1 & 0 \end{bmatrix} \text{ and } \det(B) = -36$$

Step 6:

Transpose the last column of B^{-1} into a new matrix D:

$$C = \begin{bmatrix} \frac{1}{3} & 1 & \frac{1}{3} & 1 & 0 \end{bmatrix}$$

Step 7:

Multiply D by the absolute value of the determinant:


$$|-36| \cdot C = D = [12 \quad 36 \quad 12 \quad 36 \quad 0]$$

Step 8:

Divide each element of E the greatest common divisor:


$$\gcd(d_{11}, \dots, d_{15}) = 12 \quad \therefore E = [1 \quad 3 \quad 1 \quad 3 \quad 0]$$

And therefore, $x_1 = x_3 = 1$ and $x_2 = x_4 = 3$ resulting in the balanced chemical equation:



Mathematical Background

To fully understand why the method above yield the stoichiometric coefficients we can explore the mathematics behind the method in greater detail. Consider the following hypothetical double displacement reaction:



Setting up the composition array at the coefficient matrix:

	$A_a B_b$	$C_c D_d$	$A_o D_p$	$C_q B_r$
A	a	0	$-o$	0
B	b	0	0	$-q$
C	0	c	0	$-r$
D	0	d	$-p$	0

 and then $K = \begin{bmatrix} a & 0 & -o & 0 \\ b & 0 & 0 & -q \\ 0 & c & 0 & -r \\ 0 & d & -p & 0 \end{bmatrix}$

Supposing that the steps to augment the matrix are not followed and the matrix-inverse method is directly applied. The inverse of K can then be found by multiplying the transposed cofactor matrix C^T with the inverse of the determinant of K:

$$\begin{aligned} K^{-1} &= \det(K)^{-1} \text{adj}(K) = \frac{1}{\det(K)} C^T \\ &= \frac{1}{acpq - bdor} C^T \end{aligned}$$

The problem encountered with chemical reactions that yield square coefficient matrices is that the determinant is always zero i.e. $\det(K) = 0$ and therefore $\det(K)^{-1} = \text{undefined}$ such that matrix K is degenerate and not invertible. This can easily be shown by using the composition matrix and the traditional method for solving the stoichiometric coefficients by using a system of equations:

$$Kx = 0 \rightarrow \begin{bmatrix} a & 0 & -o & 0 \\ b & 0 & 0 & -q \\ 0 & c & 0 & -r \\ 0 & d & -p & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

It is immediately apparent that if $Kx=0$ then the column vector x must be equal to zero as well. Furthermore, it can be conclusively demonstrated that $\det(K) = 0$ by setting up the system of linear equations from the above and solving them simultaneously:

$$\begin{aligned} ax_1 - ox_3 &= 0 & \dots(1) \\ bx_1 - qx_4 &= 0 & \dots(2) \\ cx_2 - rx_4 &= 0 & \dots(3) \\ dx_2 - px_3 &= 0 & \dots(4) \end{aligned}$$

$$\begin{array}{ccc} dx_2 = px_2 & & cx_2 = rx_4 & & bx_1 = qx_4 \\ dx_2 = \frac{pax_1}{o} & \text{and} & x_2 = \frac{rx_4}{c} & \text{and} & \frac{bx_1}{q} = x_4 \end{array}$$


$$\text{so that } d\left(\frac{rx_4}{c}\right) = \frac{pax_1}{o} \text{ becomes } \frac{dr}{c}\left(\frac{bx_1}{q}\right) = \frac{pax_1}{o}$$

$$\text{and then } \frac{drb}{cq} = \frac{pa}{o}$$

$$drbo = pacq$$

$$acpq - bdor = 0$$

Revealing that determinant of the square coefficient matrix is indeed zero. If the equation is viewed carefully and keeping in mind that the column vector x is zero, we can see why this occurs. These reaction equations can be balanced in an infinite number of ways and therefore do not have a single unique solution, for example:



The algebraic method is still a step up from solving by inspection and the coefficients can be solved by hand or using a suitable software package, such as Matlab, (Gabriel *et. al*, 2015) by finding the row echelon form of the augmented matrix K and solving the subsequent system of equations. However, as the complexity of reaction equations increases so does the effort required to solve by row echelon form.

It is therefore apparent that a better method is needed to approach these problems by making use of the tools available to modern scientific calculators which almost every chemistry student will be equipped with. The steps outlined in the first section allows us to construct an invertible matrix such that $\det(B) \neq 0$. Let us now repeat the first hypothetical reaction with this method:

$$B = \begin{bmatrix} a & 0 & -o & 0 & 0 \\ b & 0 & 0 & -q & 0 \\ 0 & c & 0 & -r & 0 \\ 0 & d & -p & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \text{ and } B^{-1} = \begin{bmatrix} 0 & \frac{1}{b} & 0 & 0 & \frac{q}{b} \\ 0 & 0 & \frac{1}{c} & 0 & \frac{r}{c} \\ -\frac{1}{o} & \frac{a}{bo} & 0 & 0 & \frac{aq}{bo} \\ 0 & 0 & 0 & 0 & 1 \\ -\frac{p}{o} & \frac{ap}{bo} & -\frac{d}{c} & 1 & \frac{acpq - bdor}{bco} \end{bmatrix}$$

Now we have that $\det(B) = bco$ and then computing matrix D and E:

$$|bco| \cdot D = E = \begin{bmatrix} coq & bor & acq & bco & acpq - bdor \end{bmatrix}$$

$$E = \begin{bmatrix} coq & bor & acq & bco & 0 \end{bmatrix}$$

Depending on the values of $e_{11} - e_{14}$ we potentially have to divide by $\gcd(e_{11}, \dots, e_{14})$ if necessary, so that the lowest stoichiometric coefficients are obtained. The validity of this result can be demonstrated by using another system of linear equations:

$$Kx = 0 \rightarrow \begin{bmatrix} a & 0 & -o & 0 & 0 \\ b & 0 & 0 & -q & 0 \\ 0 & c & 0 & -r & 0 \\ 0 & d & -p & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} w \\ x \\ y \\ z \\ k_1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ z \end{bmatrix}$$

Yielding a slightly modified version of the set of equations found in the first instance:

$$\begin{aligned} ax_4 - ox_3 &= 0 & \dots(1) \\ bx_1 - qx_4 &= 0 & \dots(2) \\ cx_2 - rx_4 &= 0 & \dots(3) \\ dx_2 - px_3 + k_1 &= 0 & \dots(4) \\ x_4 &= x_4 & \dots(5) \end{aligned}$$

It can now be seen that the reason for the augmentation with reversal matrices specifically, as per step 3 of the methodology, is that the coefficients $k_1, k_2 \dots k_i$ will be equal to zero and several trivial equations such as $x_4 = x_4$ will be formed. This way the augmentation does not change the meaning of the original equation itself and therefore is mathematically sound. Solving the system of equations:

$$\begin{aligned} k_1 &= px_3 - dx_2 \\ &= p\left(\frac{ax_1}{o}\right) - d\left(\frac{rx_4}{c}\right) \\ &= \frac{x_4(acpq - bdor)}{obc} \\ obc \cdot k_1 &= x_4(acpq - bdor) \end{aligned}$$

However, we have that $x_1, x_2, x_3, x_4 \neq k_i = 0$ and from the earlier example $\det(K) = acpq - bdor = 0$ and therefore we can conclude that:

$$\begin{aligned}(obc) \cdot k_1 &= x_4 (acpq - bdor) \\ (obc) \cdot k_1 &= x_4 (0)\end{aligned}$$

From which it follows that $x_4 = obc$ leading to:

$$\begin{aligned}x_1 &= \frac{qx_4}{b} & x_2 &= \frac{rx_4}{c} & x_3 &= \frac{ax_1}{o} \\ &= \frac{q \cdot obc}{b} & &= \frac{r \cdot obc}{c} & &= \frac{a \cdot coq}{o} \\ &= coq & &= rob & &= acq\end{aligned}$$

Therefore $x_1 = coq$; $x_2 = bow$; $x_3 = acq$; $x_4 = bco$; $k_1 = acpq - bdor$ as expected.

We will explore the general case and show why this method should work for all chemical equations. Focusing on the matrix of minors:

$$M = \begin{bmatrix} M_{11} & M_{12} & \cdots & M_{1m} \\ M_{21} & M_{22} & \cdots & M_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ M_{n1} & M_{n2} & \cdots & M_{nm} \end{bmatrix}$$

Where M_{ij} denotes at row i and column j . Since the first row of a composition matrix for any chemical reaction cannot be all zeros, in fact it must contain at least two non-zero elements, the minors of the i^{th} row will never be equal to zero. Furthermore, any appended k^{th} columns will necessarily lead to the minors of the i^{th} row containing only elements from the original composition array. These two propositions can be demonstrated by using simple Laplace expansion:

$$B = \begin{bmatrix} a_{11} & a_{12} & a_{13} & 0 \\ a_{21} & a_{22} & a_{23} & 0 \\ a_{31} & a_{32} & a_{33} & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} \quad \text{then } M_{11} = \begin{vmatrix} a_{22} & a_{23} & 0 \\ a_{32} & a_{33} & 1 \\ 0 & 1 & 0 \end{vmatrix} = -a_{22}$$

and so forth for all minors until the matrix of minors is found:

$$M = \begin{bmatrix} -a_{22} & -a_{21} & 0 & a_{21}a_{32} - a_{22}a_{31} \\ -a_{12} & -a_{11} & 0 & a_{11}a_{32} - a_{12}a_{31} \\ 0 & 0 & 0 & a_{11}a_{22} - a_{12}a_{21} \\ a_{12}a_{23} - a_{13}a_{22} & a_{11}a_{23} - a_{13}a_{21} & a_{11}a_{22} - a_{12}a_{21} & \det(A) \end{bmatrix}$$

This assumed a square composition matrix as starting point, but the above method holds for the other 2 types of matrices as per step 3 of the methodology as well. Additional examples as well as possible real-life applications are shown in Appendix A. Re-examining the previous equation and matrix K, the minor of K becomes:

$$M = \begin{bmatrix} 0 & 0 & -bd & 0 & -bdp \\ -co & 0 & -ac & 0 & -acp \\ 0 & -bo & 0 & 0 & -bdo \\ 0 & 0 & 0 & 0 & -bco \\ coq & -bor & acq & -bco & acpq - bdor \end{bmatrix}$$

Notice that the last row of minors yields the stoichiometric coefficients and indeed only contain elements from the composition array. This is a result of step 5 – 7 in the methodology, where we multiplied the previous result with the $\det(B)$:

$$B^{-1} = \frac{1}{\det(B)} C^T$$

$$\det(B) \times B^{-1} = C^T$$

Since C^T is found by the minors $C_{ij} = (-1)^{i+j} M_{ij}$ a more efficient modified method is now revealed. Follow steps 1 – 4 as before but then instead of determining the inverse of the matrix B, calculate and use the absolute values of the minors of the y^{th} row of B only:


$$D = \begin{bmatrix} coq & bor & acq & bco & \det(K) \end{bmatrix}$$

$$= \begin{bmatrix} coq & bor & acq & bco & 0 \end{bmatrix}$$

And then dividing each element by the greatest common divisor. This new method yields the same stoichiometric coefficients in far fewer steps and since the tools available to chemistry students enable them to calculate determinants with relative ease, find the minors in last row becomes a relatively trivial task.

Reduction of Chromite with Aluminium

Let us return our attention now to the original problem presented by the reduction reaction of chromite.



We will now solve the stoichiometric coefficients using the cofactors only and using the matrix-inverse method below:

	$(Mg_{0.44}Fe_{0.55})$ $(Cr_{1.36}Al_{0.55}Fe_{0.08})_2O_4$	Al	Mg	Fe	Cr	Al_2O_3
Mg	44	0	-1	0	0	0
Fe	63	0	0	-1	0	0
Cr	136	0	0	0	-1	0
Al	55	1	0	0	0	-2
O	397	0	0	0	0	-3

From this array we then extract the matrix K :

$$K = \begin{bmatrix} 44 & 0 & -1 & 0 & 0 & 0 \\ 63 & 0 & 0 & -1 & 0 & 0 \\ 136 & 0 & 0 & 0 & -1 & 0 \\ 55 & 1 & 0 & 0 & 0 & -2 \\ 397 & 0 & 0 & 0 & 0 & -3 \end{bmatrix}$$


Construct the augmented matrix A according to the rules and extract the coefficient matrix B:

$$B = \begin{bmatrix} 44 & 0 & -1 & 0 & 0 & 0 \\ 63 & 0 & 0 & -1 & 0 & 0 \\ 136 & 0 & 0 & 0 & -1 & 0 \\ 55 & 1 & 0 & 0 & 0 & -2 \\ 397 & 0 & 0 & 0 & 0 & -3 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Determine the absolute minors $M_{61} - M_{66}$:

$$D = [3 \quad 629 \quad 132 \quad 189 \quad 408 \quad 397]$$

Since $\gcd(d_{11}, \dots, d_{15}) = 1$ we have the balanced equation:



Using the full augmented matrix-inversion method as described earlier. Determine the inverse matrix B^{-1} as well as the determinant $\det(B)$:

$$B^{-1} = \begin{bmatrix} 0 & 0 & 0 & 0 & \frac{1}{397} & \frac{3}{397} \\ 0 & 0 & 0 & 1 & -\frac{55}{397} & \frac{629}{397} \\ -1 & 0 & 0 & 0 & \frac{44}{397} & \frac{132}{397} \\ 0 & -1 & 0 & 0 & \frac{63}{397} & \frac{189}{397} \\ 0 & 0 & -1 & 0 & \frac{136}{397} & \frac{408}{397} \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \text{ and } \det(B) = 397$$


Transpose the last column of B^{-1} into a new matrix C:

$$C = \begin{bmatrix} \frac{3}{397} & \frac{629}{397} & \frac{132}{397} & \frac{189}{397} & \frac{408}{397} & 1 \end{bmatrix}$$

Multiply D by the absolute value of the determinant:

$$|397| \cdot C = D = \begin{bmatrix} 3 & 629 & 132 & 189 & 408 & 397 \end{bmatrix}$$

Since $\gcd(d_{11}, \dots, d_{15}) = 1$ we again get the balanced equation:



Given the large values of the stoichiometric coefficients it would have taken a considerable amount of time to solve the reaction equation by inspection but using the new cofactor method the results was determined with ease.

Conclusion

Based on the need to quickly determine the stoichiometric coefficients of reaction equations, the proven, but slow, method of using a system of linear equations was expanded upon. The resulting procedure of setting up an augmented matrix and then determining minors results in an efficient and convenient method that can handle even large reaction equations with ease.

Acknowledgements

I would like to thank Ryan Ball of CAMICO SA for generously supplying samples of chromite from their Johannesburg stockpile at no cost as well as providing valuable and experienced insight into the workings of the chromium industry in South Africa.

References

- Lavoisier, A. *Traité élémentaire de chimie*. Paris, France: Cuchet, 1789.
- IUPAC. *Compendium of Chemical Terminology*, 2nd ed.. Compiled by A. D. McNaught and A. Wilkinson. Blackwell Scientific Publications, Oxford, 1997.
- Brown, T. LeMay H. Bursten, B. Murphy, C. Woodward, P. Stoltzfus, M. *Chemistry: The Central Science*, 13th Ed. Pearson, 2015
- Toth, Z. *Balancing chemical equations by inspection*. Journal Chemical Education, 74 (11), 1363. 1997.
- Gabriel, C. Onwuka, G. *Balancing of Chemical Equations Using Matrix Algebra*. Journal of Natural Sciences Research. 2015
- Schwab, B. *Working with electron microprobe data from a high pressure experiment – Calculating mineral formulas, unit cell content and geothermometry*. Humboldt State University, 2019.

Appendix A

Mineral Structure of Chromite

For the example above a sample of South African chromite concentrate was analysed at a SANAS accredited laboratory and found to have a composition of metal oxides by weight percent as seen in column two of table 1. The method as described by (Schwab, 2019) was then used to calculate the mineral structure.

Oxides Present	% w/w	Cations in Oxide	Anions in Oxide	Molar Mass	Mol Proportion	Oxygen Proportion	Number of Anions	Number of Cations
Al₂O₃	13.94	2	3	101.961	0.137	0.410	0.831	0.554
Fe₂O₃	3.00	2	3	159.687	0.019	0.056	0.114	0.076
Cr₂O₃	51.10	2	3	151.990	0.336	1.009	2.044	1.363
FeO	19.57	1	1	71.846	0.272	0.272	0.552	0.552
CaO	0.40	1	1	56.077	0.007	0.007	0.014	0.014
MgO	8.84	1	1	40.311	0.219	0.219	0.444	0.444
<u><u>Σ 96.85</u></u>						<u><u>Σ 1.974</u></u>	<u><u>4.000</u></u>	

Oxygen Atoms in Structure

4

2.026

Chromite is found as a spinel conforming to the general formula of AB_2O_4 , where A^{2+} and B^{3+} denote the cations that occupy the tetrahedral and octahedral sites respectively (Weller, 2014). Typical chromite has the formula $FeCr_2O_4$ but the Fe^{2+} and Cr^{3+} cations are frequently replaced by other ions such as Mg^{2+} and Al^{3+} which can be clearly seen from the percentage compositions above. From the data it was calculated that the approximate structure of this specific sample of concentrate is $(Mg_{0.44}Fe_{0.55})(Cr_{1.36}Al_{0.55}Fe_{0.08})_2O_4$. A molecular formula was found:

$$\begin{aligned}
 &\therefore (Mg_{0.44}Fe_{0.55})(Cr_{1.36}Al_{0.55}Fe_{0.08})_2O_4 \\
 &= Mg_{0.44}O_{0.44} \cdot Fe_{0.55}O_{0.55} \cdot Cr_{1.36}O_{2.04} \cdot Al_{0.55}O_{0.83} \cdot Fe_{0.08}O_{0.11} \\
 &= Mg_{44}O_{44} \cdot Fe_{55}O_{55} \cdot Cr_{136}O_{204} \cdot Al_{55}O_{83} \cdot Fe_8O_{11} \\
 &= Mg_{44}Fe_{63}Cr_{136}Al_{55}O_{397}
 \end{aligned}$$

This formula can now be used in the reduction reaction.

Additional examples of matrices of minors

Assuming different starting matrices A and the corresponding augmented matrices B:

2x2 square matrix:

$$B = \begin{bmatrix} a_{11} & a_{12} & 0 \\ a_{21} & a_{22} & 1 \\ 0 & 1 & 0 \end{bmatrix} \rightarrow M = \begin{bmatrix} -1 & 0 & a_{21} \\ 0 & 0 & a_{11} \\ a_{12} & a_{11} & a_{11}a_{22} - a_{12}a_{21} \end{bmatrix}$$

Applications: Simple synthesis reactions: $A_a + B_b \rightarrow A_o B_p$ e.g. $2H_2 + O_2 \rightarrow 2H_2O$.

2x3 non-square matrix:

$$B = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 1 & 0 \end{bmatrix} \rightarrow M = \begin{bmatrix} -a_{23} & 0 & a_{21} \\ -a_{13} & 0 & a_{11} \\ a_{12}a_{23} - a_{13}a_{22} & a_{11}a_{23} - a_{13}a_{21} & a_{11}a_{22} - a_{12}a_{21} \end{bmatrix}$$

Applications: Addition reactions $A_a B_b + C_c \rightarrow A_o B_p C_q$ e.g. $C_2H_4 + Br_2 \rightarrow C_2H_4Br_2$

3x4 non-square matrix:

$$B = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


↓

$$M = \begin{bmatrix} a_{22}a_{33} - a_{23}a_{32} & a_{21}a_{33} - a_{23}a_{31} & a_{21}a_{32} - a_{22}a_{31} & 0 \\ a_{12}a_{33} - a_{13}a_{32} & a_{11}a_{33} - a_{13}a_{31} & a_{11}a_{32} - a_{12}a_{31} & 0 \\ a_{12}a_{23} - a_{13}a_{22} & a_{11}a_{23} - a_{13}a_{21} & a_{11}a_{22} - a_{12}a_{21} & 0 \\ a_{12}(a_{23}a_{34} - a_{24}a_{33}) - a_{11}(a_{23}a_{34} - a_{24}a_{33}) - a_{11}(a_{22}a_{34} - a_{24}a_{32}) - a_{11}(a_{23}a_{33} - a_{23}a_{32}) - \\ a_{13}(a_{22}a_{34} - a_{24}a_{32}) + a_{13}(a_{21}a_{34} - a_{24}a_{31}) + a_{12}(a_{21}a_{34} - a_{24}a_{31}) + a_{12}(a_{21}a_{33} - a_{23}a_{31}) + \\ a_{14}(a_{21}a_{33} - a_{23}a_{31}) & a_{14}(a_{21}a_{33} - a_{23}a_{31}) & a_{14}(a_{21}a_{32} - a_{22}a_{31}) & a_{13}(a_{21}a_{32} - a_{22}a_{31}) \end{bmatrix}$$

Applications: Reduction of metal oxides $M_a O_b + R_c \rightarrow M + R_o C_p$ e.g. $Fe_2O_3 + 3C \rightarrow 2Fe + 3CO$

At this point we will switch over to using real and hypothetical reaction equations with numerical coefficients since the size of the matrices using only variable like the ones above, quickly span multiple pages. However, note that the last row of the minor of B still only contains elements from the original matrix A.


Hypothetical reaction:



$$B = \begin{bmatrix} 2 & 0 & -12 & -4 & 0 & 0 & 0 \\ 3 & 0 & -12 & -9 & 0 & 0 & 0 \\ 1 & 0 & -10 & 0 & 0 & 0 & 1 \\ 0 & 1 & -2 & 0 & 0 & 1 & 0 \\ 0 & 2 & -4 & 0 & 0 & 0 & 0 \\ 0 & 1 & -2 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix} \rightarrow E = [120 \quad 24 \quad 12 \quad 24 \quad 0 \quad 0 \quad 0]$$

Therefore, we have $10A_2B_3C_1 + 2DE_2F \rightarrow A_{12}B_{12}C_{10}(DE_2F)_2 + 2A_4B_9$


Combustion of octane:




$$B = \begin{bmatrix} 8 & 0 & -1 & 0 \\ 18 & 0 & 0 & -2 \\ 0 & 2 & -2 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow E = [4 \quad 50 \quad 32 \quad 36]$$

Therefore, we have $2C_8H_{18} + 25O_2 \rightarrow 16CO_2 + 18H_2O$

Redox reaction involving potassium ferrocyanide:



$$B = \begin{bmatrix} 4 & 1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & -2 & 0 & 0 & 0 & 0 \\ 6 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 6 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 4 & 4 & -4 & -12 & -4 & -3 & -2 & -1 \\ 0 & 0 & 2 & -1 & 0 & 0 & -1 & 0 & -2 \\ 0 & 0 & 1 & -1 & 1-3 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow E = [10 \quad 122 \quad 299 \quad 162 \quad 5 \quad 122 \quad 60 \quad 60 \quad 188]$$



↓

