Enantioselective fullerene functionalization *via* stereochemical information transfer from a self-assembled cage

AUTHORS

Zifei Lu¹, Tanya K. Ronson¹, Andrew W. Heard¹, Sascha Feldmann², Nicolas Vanthuyne³, Alexandre Martinez³, Jonathan R. Nitschke*¹

¹Yusuf Hamied Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, United Kingdom

²Cavendish Laboratory, University of Cambridge, J J Thomson Avenue, Cambridge, CB3 0HE, United Kingdom

³Aix Marseille University, CNRS Centrale Marseille, iSm2, 13284 Marseille, France

Metal-organic cages have served as masks to effect the regioselective functionalization of C₆₀. The enantioselective functionalization of C₆₀ remains challenging, however, requiring complex chiral tethers or challenging chromatography. We report a means of defining up to six stereocentres on C₆₀, achieving both enantioselective and regioselective fullerene functionalisation. This method involves stereochemical information transfer from a chiral formylpyridine, through the framework of an enantiopure cage incorporating it, to a series of regio- and stereochemically defined C₆₀ adducts. The surrounding cage is then opened with retention of fullerene stereochemistry. The chiral formylpyridine thus ultimately dictates the stereochemistry of these chiral fullerene adducts without being incorporated into them. The cage acts both as the substrate and “blueprint” for the chiral information imprinted into the fullerene adducts, which form stereoselectively with a distance of up to 6 carbon atoms between stereocentres. Such chiral fullerene adducts may become useful in devices requiring circularly-polarized light manipulation.

INTRODUCTION

The flow of stereochemical information is integral to the processes of life, such as the recognition of steroids and saccharides by receptors, and the stereospecific transformations of substrates by enzymes. Stereochemical communication is likewise central to elegant artificial processes¹, which include extraordinarily remote stereocontrol through foldamer chains^{2–4}, the transduction of rotatory motion from molecular motors up to the macroscopic scale⁵, and the creation of valuable pharmaceuticals through organocatalysis⁶.

Many metal-organic cages are chiral, with stereochemistry controlled through the use of enantiopure ligands prepared through multi-step organic synthesis and preparative chiral HPLC^{7–9}. Alternatively, chiral counterions, chiral amines¹⁰ or auxiliary ligands¹¹ have enabled stereoselective cage construction. In recent years several elegant examples of asymmetric recognition and catalysis within the chirotopic cavities¹² of enantiopure cages have been reported^{13–15}. However, achieving these functions remains challenging due to the pseudo-spherical nature of the voids in high-symmetry molecular cages, which makes it difficult for them

to differentiate between enantiomers of chiral guests. New strategies are thus needed, both to control the stereochemistry of cages and to put their chirality to practical use.

Aromatic panelled metal-organic cages have been shown to be excellent hosts for fullerenes^{16–19} and have been used to control their regioselective functionalization, through confinement effects²⁰ or by acting as a supramolecular mask^{21–24} to control the sites of substrate addition. Such functionalised fullerenes are of interest due to their applications in optoelectronics, medicine, and materials science^{25–27}. Their development has been hampered by the difficulty of selective multiple functionalisation of the spherical fullerene core and challenges associated with separation of the desired fullerenes by column chromatography or by high performance liquid chromatography (HPLC)^{28,29}. Less use has thus been made of enantiopure fullerenes than would otherwise have been the case, despite the promise they show in chiral induction of polymers²⁵, peptide synthesis²⁶ and the detection of circularly polarized light (CPL)²⁷. Their construction has relied on bulky, permanently-attached tethers, chiral HPLC, or lengthy synthetic procedures^{29–31}. Chiral derivatives of C₆₀ fall into two main categories: those with functionalization patterns that are not inherently chiral, but that possess stereocentres in their substituents, and those with inherently chiral functionalization patterns, having chirality induced by the geometric arrangement of addends. This second category is particularly challenging to access without the use of chiral HPLC or chiral auxiliaries to aid purification. A larger steric difference between stereoisomers may be observed, however, when chirality results from the geometric arrangement of addends, rather than remotely attached on the substituents, adding interest to this second group of fullerene adducts.³²

Here we report the synthesis of an enantiopure metal-organic cage, whose stereochemistry is defined through the incorporation of a chiral formylpyridine subcomponent. This cage reacts with the fullerenes C₆₀ and [6,6]-Phenyl-C61-butyric acid methyl ester (PCBM) *via* chemo-, regio- and enantioselective Diels-Alder cycloaddition with the anthracene panels of the cage. The resulting fullerene adducts may be readily cleaved from the cage. These fullerenes have inherently chiral functionalization patterns, but contain none of the initial chiral formylpyridine, which may be recovered following cage disassembly. Optical spectroscopy confirms the existence of an excited state with a microsecond lifetime for one of the adducts, which we assign to a triplet exciton carrying net spin. Together with the observed circular dichroism, this long-lived excited state renders the reported family of materials promising candidates for future optoelectronic applications exploiting chirally-induced spin-selectivity.³³

RESULTS AND DISCUSSION

Enantiopure formylpyridine subcomponent **A** (Fig. 1a) was synthesized through a three-step process in 81% overall yield (Supplementary Scheme S1), with the aim of using it to shape the stereochemistry of new cage complexes. Whereas chiral amines had been incorporated into cage frameworks¹⁰, chiral aldehydes have not yet been used to control the handedness of metallosupramolecular constructs prepared through subcomponent self-assembly. A chiral 1-phenylethanol unit was installed in the 3-position of the pyridine ring of **A** to ensure its proximity to the metal centre while avoiding steric clash around the coordination

sphere of the metal ion, which could destabilise subsequent assemblies. With the two enantiomers of **A** in hand, we set out to explore the stereochemistry of cages that form from it *via* subcomponent self-assembly.

Self-assembly with linear terphenyl subcomponent **B**

The reaction of linear terphenyl subcomponent **B** with 2-formylpyridine and iron(II) was shown to produce a mixture of homochiral *T* ($\Lambda\Lambda\Lambda\Lambda/\Delta\Delta\Delta\Delta$), heterochiral C_3 ($\Delta\Delta\Delta\Lambda/\Lambda\Lambda\Lambda\Delta$), and achiral S_4 ($\Lambda\Lambda\Delta\Delta$) $\text{Fe}^{\text{II}}\text{L}_6$ cage diastereomers³⁴. We thus investigated the degree of stereocontrol that chiral subcomponent **A** might engender within an analogous tetrahedral cage framework incorporating **B** residues. The reaction between (*R*)-**A**, **B** and iron(II) trifluoromethanesulfonimide (triflimide, NTf_2^-) yielded $\text{Fe}^{\text{II}}\text{L}_6$ cage **1** (Fig. 1a), as confirmed by ESI-MS (Supplementary Fig. S15). The ^1H NMR and ^{13}C NMR spectra of **1** (Supplementary Figs. S8 and S11) were complex, exhibiting clusters of peaks. All ^1H NMR peaks between 4.50 and 9.50 ppm displayed a single diffusion constant in ^1H DOSY NMR (Supplementary Fig. S14), consistent with the presence of multiple diastereomeric species having the same size.

Fig. 1 | Preparation and characterization of cages 1 and 2. **a**, Synthesis of cage **1** from (*R*)-**A** as a mixture of diastereomers; only the majority *T*-symmetric diastereomer is shown. **b**, Synthesis of enantiomeric cages $\Lambda\Lambda\Lambda\Lambda$ -**2** and $\Delta\Delta\Delta\Delta$ -**2** from (*R*)-**A** and (*S*)-**A**, respectively. **c**, Comparison of the CD spectra of cages **1** and $\Lambda\Lambda\Lambda\Lambda$ -**2** (MLCT region shown, data normalized to the UV-Vis absorption at the MLCT maxima at 585 nm for **1** and **2**). **d**, CD spectra of the two enantiomers of cage **2**. **e**, X-ray crystal structure of $\Lambda\Lambda\Lambda\Lambda$ -**2** (Fe^{II} , green; C, grey; N, blue; O, red; H, white, solvent of crystallization and anions omitted for clarity).

Examination of the imine region of the ^1H NMR spectrum of **1** revealed a major peak assigned to the *T*-symmetric diastereomer, comprising 50% of the integrated imine intensity. This observation is consistent with a 1.5-fold increase in preference for this diastereomer relative to the unsubstituted cage, where an approximately equimolar mixture of the three diastereomers had been observed³⁴, already reflecting a slight

preference for the *T*-symmetric diastereomer as compared to a statistical distribution of 12.5% *T*, 50% *C*₃ and 37.5% *S*₄.

The incomplete stereochemical induction in **1** was attributed to the limited ability of dianiline **B** to transmit stereochemical information between vertices. We had previously hypothesised that cages incorporating such linear dianilines exhibited weaker stereochemical coupling between metal-centre vertices as compared to bis(2-formylpyridine)-based cages because of the lack of offset between metal-binding sites within the subsequently formed imine ligand³⁵. Such dianiline-based cages exhibit less distortion of the ligands when mixtures of *syn*- and *anti*- ligands are incorporated into the *C*₃ and *S*₄ diastereomers, and thus a decreased preference for the *T* diastereomer where all ligands adopt the *anti*-conformation. The moderate degree of chiral induction observed with (*R*)-**A** prompted us to explore chiral induction with aniline subcomponents having a greater degree of offset.

Self-assembly with 1,5-anthracene-containing subcomponent C

Dianiline subcomponent **C**, which contains a 1,5-anthracene spacer, was previously shown to form only the homochiral *T*-symmetric cage diastereomer upon self-assembly with Fe(NTf₂)₂ and 2-formylpyridine³⁶. The self-assembly of **C** (6 equiv) and enantiopure formylpyridine (*R*)-**A** (12 equiv) with Fe(NTf₂)₂ (4 equiv) yielded Fe^{II}₄L₆ cage **2**, as confirmed by ESI-MS (Fig. 1b and Supplementary Fig. S28).

The ¹H NMR spectrum of cage **2** showed a single imine peak at 9.28 ppm (Supplementary Fig. S22), indicating the selective formation of a single cage diastereomer. The circular dichroism (CD) spectrum of $\Lambda\Lambda\Lambda\Lambda$ -**2** contains negative exciton-coupled circular dichroism (ECCD) couplets around 550 nm (Fig. 1d and Supplementary Fig. S33) which correspond to the metal-to-ligand charge-transfer (MLCT) bands for Fe^{II} pyridyl-imine complexes, consistent with the presence of metal stereocentres with predominantly Λ handedness¹⁰. The solid-state structure of cage $\Lambda\Lambda\Lambda\Lambda$ -**2** was determined by single crystal X-ray diffraction analysis (Fig. 1e). This crystal structure confirmed Λ handedness for all four metal centres.

The Λ helicity of the metal stereocentres seen in the crystal structure of **2** agree with the intense ECCD couplet from coupling of the π - π^* transitions in the imine chromophores seen in the CD spectrum indicating the selective formation of the $\Lambda\Lambda\Lambda\Lambda$ isomer in solution³⁷. The enantiomeric cage $\Delta\Delta\Delta\Delta$ -**2** was also prepared, starting from (*S*)-**A**, and the CD spectra of both cages appear with exact mirror symmetry (Fig. 1d).

The CD and UV-vis absorption spectra of cages **1** and $\Lambda\Lambda\Lambda\Lambda$ -**2** are very similar in the MLCT region (400-700 nm). We thus compared the MLCT CD signals of cages **1** and $\Lambda\Lambda\Lambda\Lambda$ -**2**, in both cases normalising the CD intensity to the corresponding maximum UV-vis absorption intensity in the MLCT region (Fig. 1c). This comparison provides insight into the degree of stereochemical coupling between the chiral aldehyde residues and the metal stereocentres within cage **1**. The proportion of ellipticity $\theta(1)/\theta(2)$ is calculated to be 0.67, indicating the presence of significant chiral induction within cage **1** despite its diastereomeric diversity. Most

of this CD signal is inferred to derive from the majority $\Delta\Delta\Delta\Delta$ *T* diastereomer, which gives rise to the single major peak (50% of the integrated intensity) in the imine region of the ^1H NMR of cage **1** noted above. Further discussion can be found in Supporting Information Section 2.2.

Thus, new formylpyridine (*R*)-**A** was shown to render cage **2** enantiopure, but to have a more limited influence over the stereochemistry of cage **1**.

Synthesis of chiral fullerenes *via* stereochemical information transfer from enantiopure cage **2**

Anthracene panels analogous to those of cage **2** have been shown³⁶ to undergo Diels–Alder reaction with C_{60} . We hypothesized that enantiopure **2** would also react with C_{60} , potentially fixing the precise stereochemistry of six carbon stereocentres in a single process.

The reaction between cage **2** and excess C_{60} (5 equiv) in acetonitrile did indeed result in the formation of a new 1:1 adduct, as confirmed by ESI-MS (Supplementary Fig. S43). ^1H NMR spectroscopy (Supplementary Fig. 35) allowed us to assign the major peaks (ca. 95%) to four magnetically distinct environments of equal intensity for each proton type, consistent with the formation of a product with C_3 symmetry.

Fig. 2 | Reaction of cage **2 with C_{60} and PCBM.** **a**, Cage **2** reacted stereoselectively with C_{60} , resulting in the formation of an adduct with six new stereocentres having defined stereochemistry. **b**, The reaction between **2** and PCBM stereoselectively formed an adduct with two new stereocentres on separate hemispheres of C_{60} . In both cases, stepwise addition of *tren* resulted first in removal of the apical metal centre, followed by complete disassembly to yield the final fullerene adducts. Cages **2** and $\text{C}_{60}\bullet\mathbf{2}$ are shown as their X-ray structures; all other structures are models produced with SCIGRESS, hydrogen atoms are omitted for clarity.

The crystal structure of $\Delta\Delta\Delta\Delta\text{-C}_{60}\bullet\mathbf{2}$ confirmed the formation of a covalent Diels–Alder adduct between $\Delta\Delta\Delta\Delta\text{-2}$ and C_{60} (Fig. 3a). The fullerene is centred within the $\text{Fe}^{\text{II}}_4\text{L}_6$ cage, having reacted with three of the

six anthracene panels. The three anthracene ligands that have undergone cycloaddition share an apical vertex of the cage, producing a C_3 -symmetric configuration consistent with solution NMR spectra. Despite the preference for a Λ metal configuration, which the (*R*)-**A** derivative was demonstrated to induce, the handedness of the apical Fe^{II} centre of $\text{C}_{60}\cdot\mathbf{2}$ was observed to invert following reaction, as was also observed in its racemic analogue.

Fig. 3 | X-ray crystal structures of cages $\Delta\Delta\Delta\Delta\text{-C}_{60}\cdot\mathbf{2}$ and $\text{PCBM}\cdot\mathbf{2}'$. **a**, Three anthracene panels per cage have reacted with C_{60} in $\Delta\Delta\Delta\Delta\text{-C}_{60}\cdot\mathbf{2}$, lending the structure C_3 symmetry. **b**, One anthracene panel per cage has reacted stereoselectively with PCBM in $\text{PCBM}\cdot\mathbf{2}'$ (carbons on the side chain of PCBM coloured yellow; both $\Delta\Delta\Delta\Delta$ and $\Lambda\Lambda\Lambda\Lambda$ enantiomers are present in the unit cell but only the $\Delta\Delta\Delta\Delta$ enantiomer of $\text{PCBM}\cdot\mathbf{2}'$ is shown here for clarity). In both cases Δ and Λ metal centres are coloured purple and green respectively and the fullerene carbon atoms are cyan. Solvents, counterions and disorder are omitted for clarity.

Encouraged by the diastereoselective reaction observed with C₆₀, we investigated the Diels–Alder reaction of cage **2** with the singly-functionalised C₆₀ derivative PCBM under identical reaction conditions. ESI-MS confirmed the formation of a 1:1 adduct (Supplementary Fig. S58). In contrast to the C₃ symmetry reflected in the ¹H NMR spectra of C₆₀•**2**, the PCBM•**2** adduct displayed complete desymmetrisation, with all protons in magnetically distinct environments (Supplementary Fig. S51).

Although numerous attempts to obtain X-ray quality crystals of PCBM•**2** were unsuccessful, we were able to obtain the crystal structure of its racemic analogue PCBM•**2'**, which was synthesized by treating PCBM with cage **2'**, an analogue of cage **2** prepared using 2-formylpyridine in place of chiral (*R*)-**A**. Solution NMR data of PCBM•**2'** also displayed complete desymmetrisation (Supplementary Figs. S65-S69).

The helicity of the apical Fe^{II} centre of PCBM•**2'** is again opposite to those of the basal Fe^{II} centres, as observed in the C₆₀ adduct. The crystal structure shows that PCBM reacted with only one of the six anthracene panels on the opposite hemisphere to the original substituent, with the resulting adduct displaying the *trans*-3 substitution pattern²⁹.

Stereoselective di-substitution of fullerenes is a challenging problem. Past efforts have enabled the regioselective di-substitution of C₆₀ through elegant masking strategies^{21,38}, but precise control over the absolute orientations of stereocentres has remained elusive, until now.

Retrieval of fullerene adducts

Taking advantage of the reversible nature of the imine bonds of the cages, we sought to retrieve the covalently trapped fullerene and PCBM adducts from the cages *via* stepwise addition of tris(2-aminoethyl)amine (*tren*) (Fig. 2).

The gradual addition of *tren* (2 equiv) to a solution of C₆₀•**2** resulted in complete transformation to a new species C₆₀•**3** after 16 h at room temperature (Fig. 2a), which was separated from the mononuclear side-product Fe^{II}(tris(ethylimino-2-pyridyl)amine) *via* precipitation with toluene. The ¹H NMR spectrum of C₆₀•**3** (Supplementary Fig. S76) was consistent with removal of the apical metal centre and three of the twelve (*R*)-**A** residues incorporated in the periphery of cage C₆₀•**2**. ESI-MS further confirmed the identity of C₆₀•**3** as a partially disassembled “open” tetrahedral cage, with three basal Fe^{II} centres and three free amines. We infer the apical Fe^{II} to have been most readily removed due to increased strain introduced following the Diels–Alder reaction, and from the “mismatch” between (*R*)-**A** and the Δ configuration of the metal centre.

Further reaction of C₆₀•**3** with excess *tren* (5 equiv) produced a yellow precipitate after 48 h at 30 °C, which was isolated *via* centrifugation. Stepwise demetallation and disassembly of C₆₀•**2** with *tren* avoided the precipitation of partially disassembled cage products that occurred during direct reaction with 5 equiv of *tren*, thus increasing product yield and purity. Product ESI-MS and ¹H NMR spectra and were consistent with C₆₀ adduct **4** (Fig. 4a) with three residues of **C** still attached.

The same procedure also allowed isolation of the other enantiomer of the C₆₀ adduct **4**, starting from $\Delta\Delta\Delta\Delta$ -**2**. The CD spectra of the enantiomers of **4** (Fig. 4b) are mirror images, as expected for mirror-image compounds. CD spectral features of the pair of enantiomers (Fig. 4b) agreed with those reported for *e,e,e* tris-adducts reported by Hirsch³² and Nierengarten³⁹. Analysis of both enantiomers of the purified **4** adduct *via* chiral stationary phase HPLC (Supplementary Fig. S81-83) revealed an *ee* of 87%.

Fig. 4 | CD spectra of retrieved fullerene adducts **4 and **5**.** **a**, PM6-minimised models⁴⁰ of both enantiomers of **4** (See Supplementary Information Section 3). **b**, CD spectra of both enantiomers of **4**. **c**, PM6 models of both enantiomers of **5**. **d**, CD spectra of both enantiomers of **5**. **e**, Transient absorption 2D pseudo-colour map of **4** excited at 400 nm with 100 fs pulses. **f**, Extracted excited-state kinetics spectrally integrated over 600-650 nm, yielding an effective lifetime of $0.24 \pm 0.05 \mu\text{s}$ indicative of a triplet state. Data at timescales longer than a ns were recorded using 355 nm excitation (see Supplementary Information Section 1 for details).

PCBM adduct **5** was also obtained from PCBM•**2** by treatment with *tren*, using the same procedure as described above for **4** (Fig. 2b). The singly-demetalated intermediate PCBM•**3** (Fig 4c) was also isolated following reaction with 2 equiv of *tren*. Analysis of purified **5** via chiral HPLC revealed an *ee* of 64%, as compared to the 87% *ee* of **4**. The CD and UV-Vis spectra of both the *e,e,e* and *trans-3* adducts (see Fig. 4 and S78-79 and S86-87) were recorded prior to HPLC purification. As expected, the enantiomers of **5** show mirror-image CD spectra (Fig 4d). Both adducts show pronounced Cotton effects similar to those observed in other fullerene derivatives with chiral functionalization patterns²⁹.

We further studied the photophysics of fullerene adduct **4** using transient absorption (TA) spectroscopy (Figs. 4e, 4f), observing a long-lived excited state centred around 600 nm with an effective lifetime of 0.24 ± 0.05 μ s. This excited state lasts much longer than the typical radiative recombination rates of $S_0 \leftarrow S_1$ transitions for conjugated organic dyes (typically nanoseconds)⁴¹. We attribute the long lifetime observed here to the presence of triplet excitons, known to possess lifetimes of up to milliseconds, which formed rapidly within our instrument response time.

Notably, these efficiently generated triplets overlap spectrally with features observed in the CD spectrum (Fig. 4b), implying that the triplet state will likely be affected by the chirality of the adduct. In contrast to singlet states ($S = 0$), triplet states possess a net spin ($S = 1$). This allows for the interaction of the triplets with the chirality of the structure via spin-orbit coupling to the chiral electrical potential, and for the resulting chiral-induced spin-selectivity (CISS) effect to be observed.^{33,42} We therefore expect **4** to be a promising candidate for molecular spintronics applications, such as spin-valves and spin qubits, where long-lived spin states controllable via the CISS effect are desirable.^{43,44}

Conclusion

Chiral formylpyridine subcomponent **A** contributes to the growing supramolecular toolbox available for the construction of enantiopure metal-organic assemblies for chiral separations and guest molecule transformations. The point chirality of **A** transferred efficiently to the metal centres of cage **2**, which in turn imprinted its stereochemistry upon the two chiral fullerene adducts **4** and **5**, which formed with high chemo-, enantio- and regio-selectivity. Chiral fullerene adducts with specific functionalization patterns were thus obtained directly in highly enantioenriched form without the need for chromatographic separation, through the transfer of stereochemical information along an unbroken chain from subcomponent **A** to cage **2** and on to **4** and **5**, surviving removal of the initial chiral aldehyde auxiliary. The chiral fullerene adduct **4** was shown not only to exhibit circular dichroism, but also long exciton lifetimes assigned to triplets, potentially allowing for manipulation of the spin states via the CISS effect. This combination of intriguing chiroptical properties renders the adducts promising building blocks for chiral organic electronics, for example as chiral hole transport layers and spin filters for spin-LEDs⁴⁵, detectors of circularly-polarised light⁴⁶, or for use in molecular quantum information science.

The authors declare that all data supporting the findings of this study are included within the article and its Supplementary Information, and are also available from the authors upon request. Crystallographic data for the structures reported in this paper have been deposited at the Cambridge Crystallographic Data Centre, under the deposition numbers 24124135 ($\Delta\Delta\Delta\Delta$ -1), 24124134 ($\Delta\Delta\Delta\Delta$ -C₆₀•2) and 24124133 (PCBM•2'). Copies of these data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif.

Acknowledgements

This work was supported by the Engineering and Physical Sciences Research Council (EPSRC, EP/P027067/1) and the European Research Council (695009). Z.L. acknowledges the Cambridge Trust and China Scholarship Council for Ph.D. funding. S.F. acknowledges funding from the Engineering and Physical Sciences Research Council (EPSRC, UK) through an EPSRC Doctoral Prize Fellowship. We thank Diamond Light Source for beamtime on Beamline I19 (CY21497). We also thank the Yusuf Hamied Department of Chemistry NMR facility for characterization data and Carles Fuertes-Espinosa for helpful discussion.

Author contributions

Z.L, T.K.R and J.R.N. conceived the study and wrote the manuscript. Z.L. performed the synthetic work with assistance from A.W.H. T.K.R. collected the X-ray data and refined the structures. N.V. and A.M. performed chiral HPLC studies on the fullerene adducts. S.F. carried out the transient absorption measurements. Z.L led the project overall. All the authors contributed to the manuscript preparation.

Additional information

Supplementary information is available in the online version of the paper. Reprints and permissions information is available online at www.nature.com/reprints. Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations. Correspondence and requests for materials should be addressed to J.R.N.

Competing financial interests

The authors declare no competing financial interests.

References

1. Percec, V. *et al.* Steric Communication of Chiral Information Observed in Dendronized Polyacetylenes. *J. Am. Chem. Soc.* **128**, 16365–16372 (2006).
2. Clayden, J., Lund, A., Vallverdú, L. & Helliwell, M. Ultra-remote stereocontrol by conformational communication of information along a carbon chain. *Nature* **431**, 966–971 (2004).
3. Yashima, E. *et al.* Supramolecular Helical Systems: Helical Assemblies of Small Molecules, Foldamers, and Polymers with Chiral Amplification and Their Functions. *Chem. Rev.* **116**, 13752–13990 (2016).
4. Tsiamantas, C. *et al.* Selective Dynamic Assembly of Disulfide Macrocyclic Helical Foldamers with Remote Communication of Handedness. *Angew. Chem. Int. Ed.* **55**, 6848–6852 (2016).
5. Eelkema, R. *et al.* Nanomotor rotates microscale objects. *Nature* **440**, 163–163 (2006).
6. Han, B. *et al.* Asymmetric organocatalysis: an enabling technology for medicinal chemistry. *Chem. Soc. Rev.* **50**, 1522–1586 (2021).
7. Zhao, C. *et al.* Chiral Amide Directed Assembly of a Diastereo- and Enantiopure Supramolecular Host and its Application to Enantioselective Catalysis of Neutral Substrates. *J. Am. Chem. Soc.* **135**, 18802–18805 (2013).
8. Pan, M., Wu, K., Zhang, J. H. & Su, C. Y. Chiral metal–organic cages/containers (MOCs): From structural and stereochemical design to applications. *Coord. Chem. Rev.* **378**, 333–349 (2019).
9. Argent, S. P., Riis-Johannessen, T., Jeffery, J. C., Harding, L. P. & Ward, M. D. Diastereoselective formation and optical activity of an M_4L_6 cage complex. *Chem. Commun.* **37**, 4647–4649 (2005).
10. Castilla, A. M. *et al.* High-Fidelity Stereochemical Memory in a $Fe^{II}_4L_4$ Tetrahedral Capsule. *J. Am. Chem. Soc.* **135**, 17999–18006 (2013).
11. Nishioka, Y., Yamaguchi, T., Kawano, M. & Fujita, M. Asymmetric [2 + 2] Olefin Cross Photoaddition in a Self-Assembled Host with Remote Chiral Auxiliaries. *J. Am. Chem. Soc.* **130**, 8160–8161 (2008).
12. Mislow, K. & Siegel, J. Stereoisomerism and local chirality. *J. Am. Chem. Soc.* **106**, 3319–3328 (1984).
13. Howlader, P., Zangrando, E. & Mukherjee, P. S. Self-Assembly of Enantiopure Pd_{12} Tetrahedral Homochiral Nanocages with Tetrazole Linkers and Chiral Recognition. *J. Am. Chem. Soc.* **142**, 9070–9078 (2020).

14. Xuan, W., Zhang, M., Liu, Y., Chen, Z. & Cui, Y. A Chiral Quadruple-Stranded Helicate Cage for Enantioselective Recognition and Separation. *J. Am. Chem. Soc.* **134**, 6904–6907 (2012).
15. Brown, C. J., Bergman, R. G. & Raymond, K. N. Enantioselective Catalysis of the Aza-Cope Rearrangement by a Chiral Supramolecular Assembly. *J. Am. Chem. Soc.* **131**, 17530–17531 (2009).
16. Mahata, K., Frischmann, P. D. & Würthner, F. Giant Electroactive M_4L_6 Tetrahedral Host Self-Assembled with Fe(II) Vertices and Perylene Bisimide Dye Edges. *J. Am. Chem. Soc.* **135**, 15656–15661 (2013).
17. Kishi, N., Li, Z., Yoza, K., Akita, M. & Yoshizawa, M. An M_2L_4 Molecular Capsule with an Anthracene Shell: Encapsulation of Large Guests up to 1 nm. *J. Am. Chem. Soc.* **133**, 11438–11441 (2011).
18. Purba, P. C., Maity, M., Bhattacharyya, S. & Mukherjee, P. S. A Self-Assembled Palladium(II) Barrel for Binding of Fullerenes and Photosensitization Ability of the Fullerene-Encapsulated Barrel. *Angew. Chem. Int. Ed.* **60**, 14109–14116 (2021).
19. Nakamura, T., Ube, H., Miyake, R. & Shionoya, M. A C_{60} -Templated Tetrameric Porphyrin Barrel Complex via Zinc-Mediated Self-Assembly Utilizing Labile Capping Ligands. *J. Am. Chem. Soc.* **135**, 18790–18793 (2013).
20. Huang, N. *et al.* Tailor-Made Pyrazolide-Based Metal–Organic Frameworks for Selective Catalysis. *J. Am. Chem. Soc.* **140**, 6383–6390 (2018).
21. Fuertes-Espinosa, C. *et al.* Supramolecular Fullerene Sponges as Catalytic Masks for Regioselective Functionalization of C_{60} . *Chem* **6**, 169–186 (2020).
22. Ubasart, E. *et al.* A three-shell supramolecular complex enables the symmetry-mismatched chemo- and regioselective bis-functionalization of C_{60} . *Nat. Chem.* **13**, 420–427 (2021).
23. Chen, B., Holstein, J. J., Horiuchi, S., Hiller, W. G. & Clever, G. H. Pd(II) Coordination Sphere Engineering: Pyridine Cages, Quinoline Bowls, and Heteroleptic Pills Binding One or Two Fullerenes. *J. Am. Chem. Soc.* **141**, 8907–8913 (2019).
24. Leonhardt, V., Fimmel, S., Krause, A. M. & Beuerle, F. A covalent organic cage compound acting as a supramolecular shadow mask for the regioselective functionalization of C_{60} . *Chem. Sci.* **11**, 8409–8415 (2020).

25. Nishimura, T. *et al.* Macromolecular Helicity Induction on a Poly(phenylacetylene) with C_2 -Symmetric Chiral [60]Fullerene-Bisadducts. *J. Am. Chem. Soc.* **126**, 11711–11717 (2004).
26. Bianco, A. *et al.* Synthesis, Chiroptical Properties, and Configurational Assignment of Fulleroproline Derivatives and Peptides. *J. Am. Chem. Soc.* **118**, 4072–4080 (1996).
27. Shi, W. *et al.* Fullerene Desymmetrization as a Means to Achieve Single-Enantiomer Electron Acceptors with Maximized Chiroptical Responsiveness. *Adv. Mater.* **33**, 2004115 (2021).
28. Fuertes-Espinosa, C., Pujals, M. & Ribas, X. Supramolecular Purification and Regioselective Functionalization of Fullerenes and Endohedral Metallofullerenes. *Chem* **6**, 3219–3262 (2020).
29. Thilgen, C. & Diederich, F. Structural Aspects of Fullerene Chemistry - A Journey through Fullerene Chirality. *Chem. Rev.* **106**, 5049–5135 (2006).
30. Riala, M. & Chronakis, N. A Facile Access to Enantiomerically Pure [60]Fullerene Bisadducts with the Inherently Chiral *Trans* -3 Addition Pattern. *Org. Lett.* **13**, 2844–2847 (2011).
31. Filippone, S., Maroto, E. E., Martín-Domenech, Á., Suarez, M. & Martín, N. An efficient approach to chiral fullerene derivatives by catalytic enantioselective 1,3-dipolar cycloadditions. *Nat. Chem.* **1**, 578–582 (2009).
32. Djojo, F. & Hirsch, A. Synthesis and Chiroptical Properties of Enantiomerically Pure Bis- and Trisadducts of C_{60} with an Inherent Chiral Addition Pattern. *Chem. – Eur. J.* **4**, 344–356 (1998).
33. Naaman, R., Paltiel, Y. & Waldeck, D. H. Chiral molecules and the electron spin. *Nat. Rev. Chem.* **3**, 250–260 (2019).
34. Meng, W., Clegg, J. K., Thoburn, J. D. & Nitschke, J. R. Controlling the Transmission of Stereochemical Information through Space in Terphenyl-Edged Fe_4L_6 Cages. *J. Am. Chem. Soc.* **133**, 13652–13660 (2011).
35. Ousaka, N. *et al.* Efficient Long-Range Stereochemical Communication and Cooperative Effects in Self-Assembled Fe_4L_6 Cages. *J. Am. Chem. Soc.* **134**, 15528–15537 (2012).
36. Ronson, T. K., Pilgrim, B. S. & Nitschke, J. R. Pathway-Dependent Post-assembly Modification of an Anthracene-Edged $M^{II}_4L_6$ Tetrahedron. *J. Am. Chem. Soc.* **138**, 10417–10420 (2016).
37. Knof, U. & von Zelewsky, A. Predetermined Chirality at Metal Centers. *Angew. Chem. Int. Ed.* **38**, 302–322 (1999).

38. Xiao, Z., Geng, X., He, D., Jia, X. & Ding, L. Development of isomer-free fullerene bisadducts for efficient polymer solar cells. *Energy Environ. Sci.* **9**, 2114–2121 (2016).
39. Guerra, S., Schillinger, F., Sigwalt, D., Holler, M. & Nierengarten, J. F. Synthesis of optically pure [60]fullerene *e,e,e*-tris adducts. *Chem. Commun.* **49**, 4752–4754 (2013).
40. Stewart, J. J. P. Optimization of parameters for semiempirical methods V: Modification of NDDO approximations and application to 70 elements. *J. Mol. Model.* **13**, 1173–1213 (2007).
41. In *Electronic Processes in Organic Semiconductors I–XIII* (John Wiley & Sons, Ltd, 2015). doi:10.1002/9783527685172.fmatter.
42. Naaman, R. & Waldeck, D. H. Spintronics and Chirality: Spin Selectivity in Electron Transport Through Chiral Molecules. *Annu. Rev. Phys. Chem.* **66**, 263–281 (2015).
43. Chappert, C., Fert, A. & Van Dau, F. N. The emergence of spin electronics in data storage. *Nat. Mater.* **6**, 813–823 (2007).
44. Wasielewski, M. R. *et al.* Exploiting chemistry and molecular systems for quantum information science. *Nat. Rev. Chem.* **4**, 490–504 (2020).
45. Kim, Y. H. *et al.* Chiral-induced spin selectivity enables a room-temperature spin light-emitting diode. *Science* **371**, 1129–1133 (2021).
46. Yang, Y., da Costa, R. C., Fuchter, M. J. & Campbell, A. J. Circularly polarized light detection by a chiral organic semiconductor transistor. *Nat. Photonics* **7**, 634–638 (2013).