


Synthesis of melazolide B, a degraded limonoid, from a natural terpene precursor

Yannan Liu,[†] Alexander W. Schuppe,[†] Yizhou Zhao, Jaehoo Lee, Timothy R. Newhouse*

Department of Chemistry, Yale University, 225 Prospect St., New Haven, Connecticut 06520-8107, United States.

*E-mail: timothy.newhouse@yale.edu



Synthesis of melazolide B, a degraded limonoid, from a natural terpene precursor

Yannan Liu^{a, 1}, Alexander W. Schuppe^{a, 1}, Yizhou Zhao^a, Jaehoo Lee^a, Timothy R. Newhouse^{a, *}

^aDepartment of Chemistry, Yale University, 225 Prospect Street, New Haven, Connecticut 06520-8107, United States

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

melazolide B

limonoid

total synthesis

natural product

terpenoid


ABSTRACT

Degraded limonoids are a subclass of limonoid natural products that derive from ring-intact or ring-rearranged limonoids. Establishment of robust synthetic routes to access them could provide valuable materials to identify the simplest active pharmacophore responsible for the observed biological activities of the parent molecules. This communication delineates the development of a divergent strategy to furnish melazolide B and several other related congeners from a common ketolactone intermediate, which was rapidly assembled from α -ionone. A chemoselective carbonyl α,β -dehydrogenation and a Wharton reduction were key strategic steps in this synthetic pathway.

1. Introduction

Limonoids are a large family of terpenoid natural products with more than a thousand members isolated to date.¹⁻³ These phytochemicals display a broad array of biological activities, ranging from anti-cancer, anti-inflammation, anti-feydant, and neurological activities.¹⁻³ Due to their diverse and intricate structures as well as interesting biological profiles, several synthetic campaigns have targeted this family of natural products.⁴⁻¹³ Over the past few years, our group has reported the total synthesis of several rearranged limonoids and pyridine-containing bislactone limonoid alkaloids,¹⁰⁻¹² such as


Scheme 1. Selected limonoids and structurally related compounds



xylogranatopyridine B (1) and PTP1B-inhibiting granatamine A (2).¹⁴ The establishment of these robust synthetic routes has enabled efficient access to them and their analogs for SAR studies.¹⁵ These investigations piqued our interest in identifying the active pharmacophore for the observed PTP1B inhibition.

Degraded limonoids, such as azedaralide (3), pyroangolensolide (4) and (-)

Scheme 2. Proposed biosynthesis of (-)-melazolide B




-melazolide B (5), are a subclass of limonoid natural products arising from ring-intact or ring-rearranged limonoids.¹⁶ Although highly speculative at this juncture, Guerriero and co-workers hypothesized, based on the co-isolation of pyroangolensolide (4) and melazolide B (5), that they may derive from an uncertain common tetranotriterpenoid precursor such as deoxyandirobin (7) (Scheme 2).¹⁶ Fragmentation along the C9-C10 bond would yield fragments resembling degraded limonoid natural products (Scheme 2).¹⁶ Mechanistic proposals have previously been suggested but have not been substantiated.^{17,18}

¹ These authors contributed equally.

* Corresponding author. Department of Chemistry, Yale University, 225 Prospect Street, New Haven, Connecticut 06520-8107, United States.
Email address: timothy.newhouse@yale.edu

Our established approach provided a synthetic pathway to access compounds related to the DE-ring fragments, such as azedaralide (**3**) and pyroangolensolide (**4**).¹² Herein we disclose the development of synthetic routes to access compounds related to the AB-ring fragments, including (–)-melazolidine B (**5**) and actinidiolide (**6**).

Scheme 3. Synthesis of C3-*epi*-melazolidine B^a



^aReagents and conditions: (1) 5 mol % (S,S)-(+)-N,N'-Bis(3,5-di-*tert*-butylsalicylidene)-1,2-cyclohexanediaminomanganese(III) chloride, 4-phenylpyridine-N-oxide (5 mol %), aq. NaOCl (1 equiv), CH₂Cl₂, 0 to 23 °C, 43%, 89:11 er, 10:1 dr; (2) 1 mol % [Rh(COD)(OH)]₂, PhMe₂SiH (1.3 equiv), THF, 23 to 60 °C, 2 h, 89%; (3) O₃, acetone, –78 °C 0.5 h, then Jones reagent (2.0 equiv), 0 to 23 °C, 2 h, 51%; (4) KHMDS (1.3 equiv), PhNTf₂ (1.3 equiv), THF, –78 to 23 °C, 1 h, 84%; (5) 5 mol % Pd(OAc)₂, 10 mol % PPh₃, formic acid (2.0 equiv), Et₃N (3.0 equiv), DMF, 60 °C, 0.5 h, 87% (6) Mn(OAc)₃ (0.7 equiv), TBHP (5.0 equiv), EtOAc, 70 °C, 3 d, 40%; (7) LiBH₄ (2.0 equiv), CeCl₃ (2.0 equiv), THF/MeOH, 0 °C to rt, 1 h, 72%, 20:1 dr; (6') Zn(TMP)₂ (1.0 equiv), diethyl allyl phosphate (1.0 equiv), 2.5 mol % [Pd(allyl)Cl]₂, PhMe, 120 °C, 3 h, 85%.

2. Results and discussion

Our initial plan focused on a bidirectional search between the known degraded limonoids, such as **5** and **6**, and our previously reported intermediate **11**. The benefit of **11** as a starting material goal is that it already contains the key ring systems, quaternary center, and two stereocenters common to these degraded limonoids.¹⁹ A cyclohexyl ketone would need to be converted to an allylic alcohol wherein the hydroxyl group has formally undergone a reductive transposition. As an added benefit, the route to **11** was robust, and involved conversion of α -ionone (**8**) by a three-step sequence involving a kinetic resolution via Jacobsen epoxidation,²⁰ 1,4-hydrosilylation, and oxidative cleavage, as shown in scheme 3A.¹² Treatment of ketone **11** with KHMDS and N-phenyl-bis(trifluoromethanesulfonylimide) resulted in the formation of an


intermediate vinyl triflate (**12**) in 84% yield, which was then reduced to alkene **13** in 87% yield (Scheme 3A). Other reductants employed in this Pd-catalyzed reduction, including Et₃SiH and Bu₃SnH, were less effective in this context.

Conversion of alkene **13** to an intermediate enone was achieved by allylic oxidation utilizing Mn(OAc)₃ and *t*-BuOOH. Employing alternative allylic oxidation conditions to furnish **5** or **15** directly, such as SeO₂ and Cr-based oxidants, were unsuccessful. A diastereoselective Luche reduction of the enone intermediate (**14**) resulted exclusively in the formation of C3-*epi*-melazolidine B (**15**).

Although **13** was not a viable intermediate to melazolidine B (**5**), considering our laboratory's lactone α,β -dehydrogenation,²¹ we reasoned that subjection of **13** to lactone α,β -dehydrogenation conditions could give rise to (–)-actinidiolide (**6**), an ionone-related compound that was proposed to be produced from kiwionoside in nature.²² Indeed, dehydrogenation of the lactone functionality in **13** with our laboratory's allyl Pd-catalyzed dehydrogenation conditions revealed that the conditions originally developed for ketone dehydrogenation were most effective (the Zn(TMP)₂ and diethyl allyl phosphate system) to produce actinidiolide (**6**), as shown in scheme 3B.²¹ Employing the conditions previously developed by our laboratory for ester²³ or amide²⁴ dehydrogenation resulted in lower conversion and diminished yield (23% and 47% yield respectively). These results suggest that the Zn(TMP)₂ system may be more general for the dehydrogenation of other basic functionalities.

In order to obtain melazolidine B (**5**), we undertook an alternative route through enone **17** (Scheme 4). Several ketone dehydrogenation conditions of **11** was first examined. A two-step sequence involving TMS enol ether formation and dehydrogenation was first attempted. Treatment of ketone **11** with KHMDS and TMSCl resulted in **16** in 36% yield (Scheme 4A). Although the original Saegusa-Ito oxidation conditions only led to full decomposition of the silyl enol ether starting material (**16**),²⁵ subjection of **16** to Tsuji's modified conditions smoothly delivered the enone product (**17**) in 91% yield (Scheme 4A).²⁶

Scheme 4. Optimization of ketone dehydrogenation^a



Entry	Conditions	% Yield (19:20) ^b
1 ^c	2.5 mol % [Pd(allyl)Cl] ₂ , LiTMP, ZnCl ₂ , allyl acetate	9:0 (30%)
2 ^c	2.5 mol % [Pd(allyl)Cl] ₂ , LDA, ZnCl ₂ , allyl acetate	1:0 (24%)
3 ^c	2.5 mol % [Pd(allyl)Cl] ₂ , Zn(TMP) ₂ , ZnCl ₂ , diethyl allyl phosphate	36:39 (95%)
4	20 mol % NiBr ₂ -glyme, Zn(TMP) ₂ , diethyl allyl phosphate	0:0 (60%)
5	NaHMDS, Mukaiyama reagent	25:0 (50%)
6	LDA, Mukaiyama reagent	6:18 (87%)
7	IBX, NMO, DMSO	19:0 (99%)
8	5 mol % Pd(TFA) ₂ , DMSO, AcOH, O ₂	64:0 (99%)
9 ^d	5 mol % Pd(TFA) ₂ , DMSO, AcOH, O ₂	92:0 (99%)

^aReagents and conditions: (1) KHMDS (1 equiv), TMSCl (1.5 equiv), THF, –78 °C to rt, 2 h, 36%; (2) 10 mol % Pd(OAc)₂, 10 mol % dppe, diallyl carbonate (1.5 equiv), MeCN, 80 °C, 4 h, 91%. ^bYield of the crude reaction mixture, using 0.05 mmol **11**, was determined by ¹H NMR using 1,2-dibromomethane as an internal standard. Conversion of **11** in parenthesis. ^cReactions conducted on 0.2 mmol scale. ^dConducted on a 3-gram scale.

Several one-step dehydrogenation conditions were surveyed (Scheme 4B). Utilizing our laboratory's allyl-Pd-catalyzed dehydrogenation conditions²¹ (entries 1–3) resulted in overoxidation (**18**), however, the Ni-catalyzed dehydrogenation conditions only gave partially recovered starting material (entry 4). Employing either Mukaiyama's reagent²⁷ (entries 5–6) or IBX (entry 7) resulted in minimal desired product.²⁸ Subjection of **11** to Stahl's Pd-

catalyzed aerobic dehydrogenation conditions afforded enone **17** with excellent selectivity (64:0, entry 8) and upon conducting on a 3-gram scale, excellent yield (92% yield, entry 9).²⁹

With enone **17** in hand, we were ready to test the proposed synthesis of melazolide B (**5**) through Wharton reduction.³⁰ Treatment of enone **17** with urea-H₂O₂ effected a nucleophilic epoxidation to furnish epoxide **19** in 67% yield as a 1:1 mixture of diastereomers (Scheme 5).¹² Reduction of the mixture of diastereomers of the α,β -epoxy ketone (**19** with Wharton's hydrazine protocol resulted in the formation of (–)-melazolide B (**5**), via the intermediacy of hydrozone **20** (Scheme 5).³⁰ Interestingly, during the reduction of the α,β -epoxy ketone (**19**) with hydrazine only the desired diastereomer of the allylic alcohol **5** was observed, whereas the undesired diastereomer was degraded. It is unclear at this juncture what those decomposition pathway or pathways are, however the presence of an electrophilic lactone was possibly a liability. This six-step sequence marks the first reported synthesis of (–)-melazolide B.

3. Conclusion

In summary, we have documented efficient synthesis of C3-*epi*-melazolide B, melazolide B and actinidiolide. Tapping into the natural terpene precursors by utilizing α -ionone as a starting material accelerated the assembly of the core bicyclic structures and led to the first reported synthesis of (–)-melazolide

1. Q.-G. Tan; X.-D. Luo. *Chemical reviews* 111 (2011) 7437–7522.
2. a. D. Taylor. In *Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products*; Springer, 1984; pp. 1–102.
3. Y. Zhang; H. Xu. *RSC advances* 7 (2017) 35191–35220.
4. D. C. Behenna; E. J. Corey. *J Am Chem Soc* 130 (2008) 6720–6721.
5. J. M. Faber; W. A. Eger; C. M. Williams. *J Org Chem* 77 (2012) 8913–8921.
6. S. M. Fu; B. Liu. *Organic Chemistry Frontiers* 7 (2020) 1903–1947.
7. B. Heasley. *European Journal of Organic Chemistry* 2011 (2011) 19–46.
8. C. Lv; X. Yan; Q. Tu; Y. Di; C. Yuan; X. Fang; Y. Ben-David; L. Xia; J. Gong; Y. Shen; Z. Yang; X. Hao. *Angew Chem Int Ed Engl* 55 (2016) 7539–7543.
9. D. M. Pinkerton; S. Chow; N. H. Eisa; K. Kainth; T. J. Vanden Berg; J. M. Burns; L. W. Guddat; G. P. Savage; A. Chadli; C. M. Williams. *Chemistry* 25 (2019) 1451–1455.
10. A. W. Schuppe; D. Huang; Y. Chen; T. R. Newhouse. *J Am Chem Soc* 140 (2018) 2062–2066.
11. A. W. Schuppe; T. R. Newhouse. *J Am Chem Soc* 139 (2017) 631–634.
12. A. W. Schuppe; Y. Zhao; Y. Liu; T. R. Newhouse. *J Am Chem Soc* 141 (2019) 9191–9196.
13. S. Yamashita; A. Naruko; Y. Nakazawa; L. Zhao; Y. Hayashi; M. Hiram. *Angew Chem Int Ed Engl* 54 (2015) 8538–8541.
14. Z. F. Zhou; H. L. Liu; W. Zhang; T. Kurtan; A. Mandi; A. Benyei; J. Li; O. Tagliatalata-Scafati; Y. W. Guo. *Tetrahedron* 70 (2014) 6444–6449.
15. T. Newhouse; A. Schuppe; Y. Liu; Y. Zhao; S. Ibarra; D. Huang; E. Wang; J. Lee; P. Loria. *Chemrxiv* (2021).
16. M. D'Ambrosio; A. Guerriero. *Phytochemistry* 60 (2002) 419–424.
17. A. Almeida; L. Dong; G. Appendino; S. Bak. *Natural Product Reports* 37 (2020) 1207–1228.
18. W.-S. Li; A. Mandi; J.-J. Liu; L. Shen; T. Kurtan; J. Wu. *The Journal of Organic Chemistry* 84 (2019) 2596–2606.
19. Z. G. Brill; M. L. Condakes; C. P. Ting; T. J. Maimone. *Chemical reviews* 117 (2017) 11753–11795.
20. E. N. Jacobsen; W. Zhang; A. R. Muci; J. R. Ecker; L. Deng. *Journal of the American Chemical Society* 113 (1991) 7063–7064.
21. Y. Chen; D. Huang; Y. Zhao; T. R. Newhouse. *Angew Chem Int Ed Engl* 56 (2017) 8258–8262.
22. T. Sakan; S. Isoe; S. B. Hyeon. *Tetrahedron Letters* 8 (1967) 1623–1627.
23. Y. Chen; J. P. Romaine; T. R. Newhouse. *Journal of the American Chemical Society* 137 (2015) 5875–5878.
24. Y. Chen; A. Turlik; T. R. Newhouse. *Journal of the American Chemical Society* 138 (2016) 1166–1169.

B. Among the reported synthesis of (–)-actinidiolide (**6**) including those from Jorgensen (7 steps),³¹ Eidman (7 steps)³², and Mori (10 steps),^{33–35} this six-step sequence represents an alternative and concise asymmetric synthesis of (–)-actinidiolide (**6**). Our robust and scalable pathway will enable future investigations into the PTP1B inhibition of these and similar compounds.

Acknowledgments

Financial support for this work was provided by Yale University, Amgen, the Dreyfus Foundation, the Sloan Foundation, Bristol-Myers Squibb (Graduate Fellowship to A.W.S.), Roberts fellowship (to Y.L.), Dox fellowship (to Y.L.), National Science Foundation (GRFP to A.W.S), NSF CAREER (1653793), and the NIH (GM118614). We gratefully acknowledge the National Science Foundation for financial support in the establishment of the Yale University High Performance Computing (HPC) Center (CNS 08-21132).

Supplementary Material

Supplementary data to this article can be found online at [place holder].

References and notes

25. Y. Ito; T. Hirao; T. Saegusa. *The Journal of Organic Chemistry* 43 (1978) 1011–1013.
26. J. Tsuji; I. Minami; I. Shimizu. *Tetrahedron letters* 24 (1983) 5635–5638.
27. T. Mukaiyama; J.-i. Matsuo; H. Kitagawa. *Chemistry letters* 29 (2000) 1250–1251.
28. K. Nicolaou; Y.-L. Zhong; P. Baran. *Journal of the American chemical society* 122 (2000) 7596–7597.
29. T. Diao; S. S. Stahl. *Journal of the American Chemical Society* 133 (2011) 14566–14569.
30. P. Wharton; D. Bohlen. *The Journal of Organic Chemistry* 26 (1961) 3615–3616.
31. S. Yao; M. Johannsen; R. G. Hazell; K. A. Jørgensen. *The Journal of organic chemistry* 63 (1998) 118–121.
32. K. F. Eidman; B. S. MacDougall. *The Journal of organic chemistry* 71 (2006) 9513–9516.
33. K. Mori; V. Khlebnikov. *Liebigs Annalen der Chemie* 1993 (1993) 77–82.
34. K. Mori; S. Aki. *Acta Chemica Scandinavica* 46 (1992) 625–629.
35. K. Mori; P. Puapoomchareon. *Liebigs Annalen der Chemie* 1991 (1991) 1053–1056.

