

Photocaged activity-based probes for spatiotemporal detection of protein S-sulfenylation in living cells

Jiahao Zhang,[†] Haiyue Peng,[†] Zi'an Chen, Guorui Li* and Jing Huang*

State Key Laboratory of Chemo/Biosensing and Chemometrics, School of Biomedical Sciences, Hunan University, Changsha 410082, China.

[†] These authors contributed equally

ABSTRACT: Protein S-sulfenylation (protein sulfenic acid), as one of the most significant oxidative post-translational modifications (OxiPTMs), plays a vital role in regulating protein function. A variety of activity-based probes have been developed to profile sulfenic acid in living cells. However, due to the transient presence and low content of sulfenic acid in living cell, high doses of probes are needed to achieve efficient labeling. More importantly, current probes have no temporal control over sulfenic acid labeling. To overcome these limitations, two caged cysteine sulfenic acid probes DYn-2-ONB and DYn-2-Cou with either an o-nitrobenzyl or coumarin protecting group were developed in this study. Both probes can be efficiently uncaged via irradiation to produce the active C-nucleophile probe DYn-2. Labeling assay in living cells demonstrated DYn-2-ONB exhibited better labeling capacity compared with DYn-2, providing it as a powerful tool to detect protein S-sulfenylation in spatio-temporally controllable manner.

Cysteine residues play a vital role in maintaining the function of many proteins since they are catalytic residues for many enzymes including transferase (desulfurase),¹ hydrolase (cysteine protease),² and isomerase (protein disulfide isomerase).³ Cysteine can undergo oxidative post-translational modifications (OxiPTMs) by reactive oxygen species (ROS) and these processes have been implicated in many cellular processes, including signal transduction,⁴ autophagy,⁵ and differentiation.⁶

Cysteine can be oxidized to different oxidation states. As shown in Fig. 1A, the thiol group (-SH) of cysteine can be first oxidized to sulfenic acid (-SOH). Sulfenic acid can be further oxidized to sulfinic acid (-SO₂H) and sulfonic acid (-SO₃H). Among these oxidation states, sulfenic acid attracts much attention due to its reversible property, which is proved to be involved in redox regulatory mechanisms and signaling pathways. Sulfenic acid is also a biomarker of cellular oxidative stress.⁷ However, sulfenic acid detection is a big challenge due to its low content, high electrophilicity and rapid turnover in cells. Moreover, it is an intermediate state, and difficult to capture.⁸ Initially protein S-sulfenylation was detected by an indirect approach which involved pre-blocking of cysteine, reduction of Cys-SOH and then labelling of nascent thiols with cysteine probes.⁹ This approach has some limitations such as extensive denaturing conditions and unmanageable reduction steps. To address these issues, different probes which directly trapped Cys-SOH were developed. Dimedone is a classical Cys-SOH probe.¹⁰ Based on this scaffold, Carroll group has generated a variety of C-nucleophile probes to achieve better reactivity and selectivity.^{7,11,12} These probes can be also attached with an alkyne or azide group, so the labelled proteins can be easily detected via CuI-catalyzed azide-alkyne click reaction (CuAAC).^{13,14} To facilitate quantitative proteomic analysis, Carroll group also developed isotope-labelled dimedone probes.¹¹ As shown in Fig. 1B, DYn-2 is a widely used Cys-SOH probe in recent years, which has been successfully applied for monitoring changes of protein sulfenylation during cell signaling¹⁵ and other cell processes. Besides dimedone-based probes, some other effective trapping reagents including strained alkynes¹⁶ and alkenes¹⁷⁻¹⁹ have also been applied for Cys-SOH detection.

So far, a variety of Cys-SOH probes have been discovered and some of them including DYn-2 have been successfully exploited for profiling protein S-sulfenylation in live cells. However, there

are still some limitations for these probes. First, high concentrations of these probes (e.g. 1 mM or even 5 mM DYn-2) were usually required for successful labeling in living cells. Sulfenic acid is short-lived and it's easy to be affected by the local microenvironment in live cell.²⁰ Although some probes have shown good selectivity towards sulfenic acid in cell-free experiments with various reactants, they might not behave the same in cells considering the complexity of cellular context. Thus a large quantity of these probes might be consumed before they reached the targets. This may account for high doses of these probes for efficient labeling of sulfenic acid in cells. Second, sulfenic acid is a transient state, which is easily to be transformed to other states. Thus an in situ activation of caged probes would be advantageous for its capture. We envision photochemistry might be a good strategy to address this issue in view of its spatial and temporal controllability.^{21,22}

To this end, here we developed photocaged probes for in situ detection of sulfenic acid in living cells. The similar approach has been applied for cysteine reactivity analysis in cells by Weerapana group previously.^{23,24} The carbonyl group in these cysteine probes can be easily modified to form protected ketal, thus abolishing their reactivity towards thiols.²⁴⁻²⁶ Inspired by these findings, we introduced photoremovable protecting groups (PPGs) to DYn-2 to obtain caged Cys-SOH probes. Among a variety of PPGs,²⁷ o-nitrobenzyl (ONB) and coumarin (Cou) derivatives are two of the most widely used due to their easy modification and removal. So, we chose them to prove our concept (Fig. 1C). As shown in Scheme. 1, both PPGs can be easily attached to DYn-2 by one step ketalization reaction to obtain the desired probes DYn-2-ONB and DYn-2-Cou. The PPG1 (4,5-dimethoxy-2-nitrophenyl) methanol is commercially available and PPG2 7-diethylamino-4-(hydroxymethyl)-coumarin can be easily synthesized by two steps using 7-diethylamino-4-methylcoumarin as the starting material (Scheme S1, ESI[†]). The detailed synthetic information and characterization data (including ¹H NMR, ¹³C NMR and HRMS) of these compounds were given in the supporting information.





Fig. 1. (A) Different oxidation states of cysteine. (B) The chemical structure of the widely used sulfenic acid probe DYn-2 and its reaction mechanism for capturing sulfenic acid. (C) This work: Photocaged activity-based probes for detection of sulfenic acid.



Scheme 1. Synthetic routes for the caged probes DYn-2-ONB (A) and DYn-2-Cou (B).

With the probes in hand, we first tested their uncaging process by HPLC analysis via irradiation with an ultraviolet (UV) lamp (365 nm, 10 W). As shown in Fig. 2B, DYn-2-ONB had a peak at retention time (RT) of 15.9 min, and DYn-2-Cou gave a peak at RT 17.2 min. Upon irradiation for 5 min, the peaks for these two probes disappeared and a major peak with RT 2.5 min which was consistent with that of DYn-2. Then we appeared monitored the conversion by irradiation with different time. As seen in Fig. 2C, the proto-cleaving processes for both probes were irradiation-time dependent. For the probe DYn-2-ONB, the deprotection efficiency reached up to 90% after 3 min UV-irradiation. Compared with DYn-2-ONB, the probe DYn-2-Cou was more photolabile. More than 90% of DYn-2 was obtained from DYn-2-Cou within 1 min irradiation. It's worth noting that besides the remaining probe and produced DYn-2, some other products peaks appeared in shorter irradiation time, which might indicate the intermediates,²⁷⁻²⁹ since eventually most of them were converted to DYn-2.

Having established that the caged Cys-SOH probes (DYn-2-ONB and DYn-2-Cou) can be efficiently uncaged by irradiation to produce the active DYn-2, we then asked whether they can be used for sulfenic acid capture. Considering the instability property of Cys-SOH, a good model would facilitate the reactivity study.³⁰ Recently Carroll group discovered dipeptide cyclic sulfenamide (CSA) as a sulfenic acid model.⁷ CSA can be readily transformed to transient sulfenic acid in situ in aqueous





Fig. 2. The uncaging process of the probes by irradiation under 365 nm. (A) Schematic illustration of the uncaging process. (B) HPLC analysis of the uncaging process. (C) Time-dependent uncaging process measured by HPLC.

solution. DYn-2 has previously been reported to react with sulfenic acid on dipeptide cyclic sulfenamide (CSA)⁷. Here we also used CSA to test the reactivity of our probes. We first tested the reaction of DYn-2 with CSA using ESI-MS. Incubation of DYn-2 with CSA (in mixed solution, v/v, PBS buffer: ACN=2: 1, pH = 7.4) at H₂O₂ for 30 min can give a major peak with m/z 544, which was consistent with that of the desired adduct (Fig. 3A and Fig. S1). We then treated the caged probes with CSA under the same condition by 5 min UV irradiation (365 nm, 10 W). As expected, we obtained the desired adduct for both probes (Fig. 3B and Fig. S2, ESI⁺).

The above results proved the reaction of our probes with a cysteine sulfenic acid. We then asked whether these probes can label Cys-SOH in purified protein. Here, USP2CD was used as a model protein to test their reaction. USP2 is a human deubiquitinase (DUB), which can remove ubiquitin from its substrate. Previous reports have shown that the function of USP2 can be regulated by ROS, via the formation of sulfenic acid of its active-site cysteine.^{31,32} We purified USP2CD, the catalytic domain of USP2 (amino acids 259-605, 42 kDa). USP2CD was first treated with or without DYn-2 in the presence of 0.5 mM H₂O₂. Then after CuI-catalyzed azide-alkyne click reaction (CuAAC), the formed adduct was analysed by western blot against streptavidin-HRP. As shown in Fig. 3C, compared with no probe treatment, DYn-2 formed adducts at 42 kDa. The caged probes were treated using the similar condition, except that the systems were irradiated by UV 365 nm for 3 min before CuAAC. The adducts were obviously observed for both probes (Fig. 3C). Coomassie blue staining showed similar amounts of samples were used for all the conditions.

Taken together, all the above data have demonstrated our probes can react with sulfenic acid either on a small molecule model or on a purified protein system. We next tested whether our probes can label protein-SOH in cell lysate or living cells. As shown in Fig. 4A, cell lysate was treated with H₂O₂ (0.5 mM) at r.t. for 20 min. After addition of probes (1 mM), the mixtures were irradiated for 3 min. The azide tag was attached by CuAAC after a further incubation for 1 h, followed by western blot analysis. Compared with the control (lane 1), all probes gave labelling bands (Fig. 4B). It was clear that among these probes, the caged probe DYn-2-ONB exhibited highest labelling



efficiency. Coomassie blue staining for the same samples indicated same amounts of proteins were loaded (Fig. S5, ESI⁺). DYn-2-Cou showed less labelling bands, maybe due to its poor solubility (data not shown). Finally, profiling protein S-sulfenylation in live cells was tested by our probes. HeLa cells were incubated with 0.5 mM H₂O₂ for 30 min, then different probes was added and incubated for 2 h. Next, after UV irradiation (or no irradiation for DYn-2) for 3 min, HeLa cells were incubated in fresh medium for 30 min. Then cells were lysed and the following procedure was same as the above. The labelling results in living cells were shown in Fig. 4C. Compared with cell lysate labelling in the presence of DYn-2, much less bands were observed in the living cell labelling. Distinctly, DYn-2-ONB still exhibited strongest labelling efficiency. Compared with that, fewer bands were observed in DYn-2-Cou treated cells. As mentioned above, the poor solubility of DYn-2-Cou can be one reason. Actually, some precipitation was observed when DYn-2-Cou was added into the medium. Phototoxicity might be another reason for its low labeling ability in living cells, since multiple studies have shown coumarin-containing reagents are toxic under irradiation.^{33,34} We evaluated cell viability with MTT assay. DYn-2-ONB showed modest phototoxicity (53% viability) in the presence of 1 mM DYn-2-ONB, 0.5 mM H₂O₂ and UV irradiation (Fig. S3A). As a comparison, DYn-2-Cou was more toxic than DYn-2-ONB. Only ~36% of cells were alive under the same treatment (Fig. S3B).

Collectively, all the data have proved DYn-2-ONB exhibited potent ability to capture protein-SOH in living cells, which was worthy of further investigation. Therefore, we carried out




Fig. 4. (A) Schematic diagram of the operation flow of in vitro and in vivo experiments (B) Comparison of the ability of DYn-2, DYn-2-ONB and DYn-2-Cou to label sulfenic acid in cell lysate. (C) Comparison of the ability of DYn-2, DYn-2-ONB and DYn-2-Cou to label sulfenic acid in living cell.

experiments with DYn-2-ONB of different concentrations. As shown in Fig. S4, the ability of DYn-2-ONB to capture sulfenic acid was concentration-dependent. It's worth noting that without UV irradiation, no labeling bands were observed in the presence of DYn-2-ONB. This result clearly substantiates our notion that the caged probe is not active in living cells, until it was activated by irradiation.

In conclusion, we developed two caged Cys-SOH probes DYn-2-ONB and DYn-2-Cou for the in situ detection of protein S-sulfenylation. The extracellular systems proved both probes can efficiently capture sulfenic acid either with a small-molecule model (CSA) or with a purified protein model (USP2CD). However, they exhibited distinct labeling property in living cells. The caged probe DYn-2-ONB showed better labeling capacity than the widely used probe DYn-2, while DYn-2-Cou labeled fewer protein-SOH, partly due to its poor solubility and its high phototoxicity towards cells. DYn-2-ONB can capture sulfenic acid in cells in a spatiotemporally controllable way, which implies that the change of sulfenic acid in cells may be monitored. The good performance of DYn-2-ONB, together with its temporal control over sulfenic acid, provides it as a powerful tool for the study of biological role of cysteine oxidation in living cells. The different labeling styles between DYn-2-ONB and the conventional probe DYn-2 also intrigue us to compare their application in Cys-SOH proteomics and this work is ongoing in our lab.

METHODS

Methods are provided in the Supporting Information.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (31871365 and 22177029) and Fundamental Research Funds for the Central Universities.

REFERENCES

- Zheng, L.; White, R. H.; Cash, V. L.; Dean, D. R., Mechanism for the Desulfurization of L-Cysteine Catalyzed by the nifS Gene Product. *Biochemistry* **1994**, *33* (15), 4714-4720.

2. Baker, E. N., Structure of actinidin, after refinement at 1.7 resolution. *J. Mol. Biol.* **1980**, *141* (4), 441-484.
3. Edman, J. C.; Ellis, L.; Blacher, R. W.; Roth, R. A., et al., Sequence of protein disulphide isomerase and implications of its relationship to thioredoxin. *Nature* **1985**, *317* (6034), 267-270.
4. Sundaresan, M.; Yu, Z.-X.; Ferrans Victor, J.; Irani, K., et al., Requirement for Generation of H₂O₂ for Platelet-Derived Growth Factor Signal Transduction. *Science* **1995**, *270* (5234), 296-299.
5. Zhang, J.; Kim, J.; Alexander, A.; Cai, S., et al., A tuberous sclerosis complex signalling node at the peroxisome regulates mTORC1 and autophagy in response to ROS. *Nat. Cell. Biol.* **2013**, *15* (10), 1186-1196.
6. Hamanaka Robert, B.; Glasauer, A.; Hoover, P.; Yang, S., et al., Mitochondrial Reactive Oxygen Species Promote Epidermal Differentiation and Hair Follicle Development. *Sci. Signal.* **2013**, *6* (261), ra8-ra8.
7. Gupta, V.; Carroll, K. S., Profiling the reactivity of cyclic C-nucleophiles towards electrophilic sulfur in cysteine sulfenic acid. *Chem. Sci.* **2016**, *7* (1), 400-415.
8. Poole, L. B.; Nelson, K. J., Discovering mechanisms of signaling-mediated cysteine oxidation. *Curr. Opin. Chem. Biol.* **2008**, *12* (1), 18-24.
9. Saurin Adrian, T.; Neubert, H.; Brennan Jonathan, P.; Eaton, P., Widespread sulfenic acid formation in tissues in response to hydrogen peroxide. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, *101* (52), 17982-17987.
10. Benitez, L. V.; Allison, W. S., The Inactivation of the Acyl Phosphatase Activity Catalyzed by the Sulfenic Acid Form of Glyceraldehyde 3-Phosphate Dehydrogenase by Dimedone and Olefins. *J. Biol. Chem.* **1974**, *249* (19), 6234-43.
11. Gupta, V.; Yang, J.; Liebler, D. C.; Carroll, K. S., Diverse Redox Reactivity Profiles of Carbon Nucleophiles. *J. Am. Chem. Soc.* **2017**, *139* (15), 5588-5595.
12. Shi, Y.; Carroll, K. S., Activity-Based Sensing for Site-Specific Proteomic Analysis of Cysteine Oxidation. *Acc. Chem. Res.* **2020**, *53* (1), 20-31.
13. Paulsen, C. E.; Truong, T. H.; Garcia, F. J.; Homann, A., et al., Peroxide-dependent sulfenylation of the EGFR catalytic site enhances kinase activity. *Nat. Chem. Biol.* **2012**, *8* (1), 57-64.
14. Seo, Y. H.; Carroll, K. S., Profiling protein thiol oxidation in tumor cells using sulfenic acid-specific antibodies. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106* (38), 16163-16168.
15. Truong, T. H.; Carroll, K. S., Redox Regulation of EGFR Signaling Through Cysteine Oxidation. *Biochemistry* **2012**, *51* (50), 9954-9965.
16. Poole, T. H.; Reisz, J. A.; Zhao, W.; Poole, L. B., et al., Strained Cycloalkynes as New Protein Sulfenic Acid Traps. *J. Am. Chem. Soc.* **2014**, *136* (17), 6167-6170.
17. Alcock, L. J.; Farrell, K. D.; Akol, M. T.; Jones, G. H., et al., Norbornene probes for the study of cysteine oxidation. *Tetrahedron* **2018**, *74* (12), 1220-1228.
18. Alcock, L. J.; Oliveira, B. L.; Deery, M. J.; Pukala, T. L., et al., Norbornene Probes for the Detection of Cysteine Sulfenic Acid in Cells. *ACS. Chem. Biol.* **2019**, *14* (4), 594-598.
19. Scinto, S. L.; Ekanayake, O.; Seneviratne, U.; Pigga, J. E., et al., Dual-Reactivity trans-Cyclooctenol Probes for Sulfenylation in Live Cells Enable Temporal Control via Bioorthogonal Quenching. *J. Am. Chem. Soc.* **2019**, *141* (28), 10932-10937.
20. Gupta, V.; Carroll, K. S., Sulfenic acid chemistry, detection and cellular lifetime. *Biochim. Biophys. Acta* **2014**, *1840* (2), 847-875.
21. Jia, S.; Sletten, E. M., Spatiotemporal Control of Biology: Synthetic Photochemistry Toolbox with Far-Red and Near-Infrared Light. *ACS. Chem. Biol.* **2021**. DOI: 10.1021/acscchembio.1c00518.
22. McKenna, S. M.; Fay, E. M.; McGouran, J. F., Flipping the Switch: Innovations in Inducible Probes for Protein Profiling. *ACS. Chem. Biol.* **2021**, *16* (12), 2719-2730.
23. Xu, P.; Kelly, M.; Vann, W. F.; Qadri, F., et al., Conjugate Vaccines from Bacterial Antigens by Squaric Acid Chemistry: A Closer Look. *Chembiochem.* **2017**, *18* (8), 799-815.
24. Abo, M.; Weerapana, E., A Caged Electrophilic Probe for Global Analysis of Cysteine Reactivity in Living Cells. *J. Am. Chem. Soc.* **2015**, *137* (22), 7087-7090.
25. Li, G.; Liang, Q.; Gong, P.; Tencer, A. H., et al., Activity-based diubiquitin probes for elucidating the linkage specificity of deubiquitinating enzymes. *Chem. Commun.* **2014**, *50* (2), 216-218.
26. Liu, J.; Li, Y.; Deol, K. K.; Strieter, E. R., Synthesis of Branched Triubiquitin Active-Site Directed Probes. *Org. Lett.* **2019**, *21* (17), 6790-6794.
27. Klán, P.; Šolomek, T.; Bochet, C. G.; Blanc, A., et al., Photoremovable Protecting Groups in Chemistry and Biology: Reaction Mechanisms and Efficacy. *Chem. Rev.* **2013**, *113* (1), 119-191.
28. Shen, W.; Zheng, J.; Zhou, Z.; Zhang, D., Approaches for the synthesis of o-nitrobenzyl and coumarin linkers for use in photocleavable biomaterials and bioconjugates and their biomedical applications. *Acta. Biomater.* **2020**, *115*, 75-91.
29. Hellrung, B.; Kamdzhilov, Y.; Schwörer, M.; Wirz, J., Photorelease of Alcohols from 2-Nitrobenzyl Ethers Proceeds via Hemiacetals and May Be Further Retarded by Buffers Intercepting the Primary aci-Nitro Intermediates. *J. Am. Chem. Soc.* **2005**, *127* (25), 8934-8935.
30. Wu, Y.; Zhang, C.; Wang, L.; Duan, H., A graph-convolutional neural network for addressing small-scale reaction prediction. *Chem. Commun.* **2021**, *57* (34), 4114-4117.
31. Ohayon, S.; Refua, M.; Hendler, A.; Aharoni, A., et al., Harnessing the Oxidation Susceptibility of Deubiquitinases for Inhibition with Small Molecules. *Angew. Chem. Int. Ed.* **2015**, *54* (2), 599-603.
32. Gopinath, P.; Mahammed, A.; Ohayon, S.; Gross, Z., et al., Understanding and predicting the potency of ROS-based enzyme inhibitors, exemplified by naphthoquinones and ubiquitin specific protease-2. *Chem. Sci.* **2016**, *7* (12), 7079-7086.
33. Zou, Q.; Fang, Y.; Zhao, Y.; Zhao, H., et al., Synthesis and in Vitro Photocytotoxicity of Coumarin Derivatives for One- and Two-Photon Excited Photodynamic Therapy. *J. Med. Chem.* **2013**, *56* (13), 5288-5294.
34. Ortega-Forte, E.; Rovira, A.; Gandioso, A.; Bonelli, J., et al., COUPY Coumarins as Novel Mitochondria-Targeted Photodynamic Therapy Anticancer Agents. *J. Med. Chem.* **2021**, *64* (23), 17209-17220.