

Cu-Catalysed Coupling of Aliphatic Amines with Alkylboronic Esters

Francesca M. Dennis,^a Antonio Romero Arenas,^a George Rodgers,^a Jonathan A. Andrews,^a Samantha L. Peralta-Arriaga,^a Benjamin M. Partridge^{a,*}

^a Department of Chemistry, University of Sheffield, Dainton Building, Sheffield, S3 7HF, United Kingdom

Email: b.m.partridge@sheffield.ac.uk


Abstract

We report a Cu-catalysed oxidative coupling of aliphatic amines with alkylboronic esters to give high value alkyl amines, products found widely in applications from medicinal chemistry to materials science. This operationally simple reaction, which can be performed on gram scale, runs under mild conditions and exhibits excellent functional group tolerance. The terminal oxidant of the reaction is O₂ from the air, avoiding the need for additional chemical oxidants. Investigation into the reaction mechanism suggests the intermediacy of an alkyl radical, generated from the boronic ester upon activation through an amino radical transfer process. To demonstrate its utility and potential for late-stage functionalization, we showcase the method as the final step in the total synthesis of a TRPV 1 antagonist.

Keywords: amination, amines, boron, Chan-Lam, copper

Methods to form C-N bonds are of high synthetic value due to the wide prevalence of N-based functional groups in biologically active compounds, materials, and catalysts. Advances in transition metal catalysis have made methods to form of Csp²-N bonds become synthetically reliable and routinely used.¹⁻⁹ However, perhaps the preparation of alkyl amines is now a greater synthetic challenge.¹⁰ Traditional methods to make alkyl amines, such as alkylation reactions and reductive amination^{11,12} often suffer from poor selectivity and the need for protecting group strategies. Other methods to form alkyl amines, such as hydrogenation¹³ and biocatalysis¹⁴⁻¹⁶ strategies are excellent particularly for process chemistry applications, but are less easily applied in discovery chemistry settings. A practical and reliable cross coupling method for the formation of alkyl amines is therefore desirable, comparable to versatility of the Buchwald-Hartwig reaction for the formation of aryl amines.¹⁷


Our approach to this challenge has been to develop an alkyl variant of the Chan-Lam amination, the oxidative coupling of an organoboron reagent with an N-based nucleophile. We and others have made some progress in this field (scheme 1),¹⁸⁻²⁴ but there is room for improvement particularly in the scope of reaction to both amine and boron coupling partners. Herein, we describe conditions for the Chan-Lam amination of alkylboronic esters with aliphatic amines, coupling partners that have previously been found challenging in alkyl Chan-Lam reactions. Importantly, our method uses O₂ from air (at atmospheric pressure) as the terminal oxidant, rather than a peroxide-based oxidant, reducing the risk and making the reaction easy to perform.²⁵ <insert reference to video in SI>




Scheme 1: Previous examples of Chan-Lam amination with alkylboron reagents.

Discussion:

We initially found that by heating boronic ester **1** with CuBr₂ in neat morpholine, amine **2** was formed in quantitative yield (Table 1, entry 1). While this provided proof of concept, it was clear that we needed to reduce the loading of amine and identify a suitable solvent to generate a more general procedure. After investigating a range of Cu-salts and ancillary ligands, we identified that conducting the reaction under air using a combination of CuBr₂ and N,O-ligand **L1** in toluene gave amine **2** in high yield (entry 2).²⁶ While ketone **3** and alcohol **4** side products were also observed, unlike our previous findings^{18,27} these were only formed in modest amounts. A survey of ligands, including chiral ligands **L2** and **L3**, gave little improvement (entry 3-4). Investigation of the reaction solvent led to improved results, with a 1:1 mixture of toluene/isopropanol giving high yield of amine **2** with minimal oxidation side products (entry 5). Under these conditions, the loading of CuBr₂ could be reduced to 10 mol% without loss of yield. However, a compromise is needed between loading of amine and yield of **2** (entries 6-7). A control experiment with CuBr₂ but without ligand suggested that the presence of **L1** had minimal effect on the reaction (entry 8). We therefore believe the likely effect of ligands, such as **L1-L3**, is to aid solubilisation of CuBr₂, rather than generate discrete Cu complexes that act as a catalyst. This perhaps why all reactions we have conducted using chiral ligands have generated **2** as a racemate. Other control experiments showed that CuBr₂ is required for amination to occur (entry 9). Under an inert atmosphere only trace amounts of amine **2** was formed (entry 10), suggesting that O₂ is the terminal oxidant in the reaction. By switching the limiting reagent to morpholine (entry 11), and using an excess of boronic ester **1**, amine **2** was still formed albeit in moderate yield. Finally, our best conditions were successfully performed on gram scale, with amine **2** isolated in excellent yield (scheme 2).


Table 1: Evaluation of reaction conditions on the yield of amine **2**.^a


Entry	CuBr ₂ mol%	L (mol%)	Amine equiv.	solvent	Yield			
					1	2	3	4
1 ^b	100	-	40	-	<5%	>95%	<5%	-
2 ^c	25	L1 (25)	4	toluene	<5%	81%	9%	10%
3 ^d	25	L2 (25)	40	-	<5%	95% ^e	-	-
4 ^d	25	L3 (25)	40	-	<5%	67% ^e	-	-
5	25	L1 (25)	4	toluene/IPA (1:1)	<5%	95%	<5%	2%
6	10	L1 (10)	4	toluene/IPA (1:1)	<5%	95%	3%	-
7	10	L1 (10)	3	toluene/IPA (1:1)	<5%	79%	6%	-
8	10	-	3.5	toluene/IPA (1:1)	<5%	95%	0%	-
9	0	-	3.5	toluene/IPA (1:1)	<5%	<5%	4%	58%
10 ^b	10	-	3.5	toluene/IPA (1:1)	93%	3%	0%	0%
11 ^f	10	-	1	toluene/IPA (1:1)	20% ^g	46%	35% ^g	12% ^g



a) Reactions performed using 0.5 mmol of boronic ester **1** unless otherwise stated. Yields determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. b) reaction carried out under an inert atmosphere. c) reaction carried out at 60 °C. d) reaction carried out at 40 °C. e) formed as a racemate. f) reaction using morpholine as the limiting reagent and 3.5 equivalents of boronic ester **1**. g) yield based on amount of boronic ester **1** instead of the limiting reagent. IPA = isopropyl alcohol.


We next looked to explore the scope and limitations of our reaction conditions (scheme 2). Pleasingly, a broad range of secondary amines could be coupled in good to excellent yield with boronic ester **1**. This includes simple cyclic amines such as pyrrolidine, piperidine and azepane (**5a-5c**). Substituted piperidines (**5d-5e**) and mono-protected piperazines (**5i-5k**) were also tolerated well. The coupling of amines of particular relevance to medicinal chemistry including fluoropiperidine (**5d**)²⁸ and bicyclic amine (**5h**),²⁹ albeit in low yield, was successful. Acyclic secondary amines could also be reacted successfully (**5l-5p**), though the yield is typically a slightly lower. Pleasingly, the reaction conditions are also successful when coupling primary aliphatic amines (**5q-5s**) – the first time these have been reported in an alkyl Chan-Lam reaction with boronic esters. However, the coupling of benzylamine was not successful. The yield of reactions with these amines are generally lower, which correlates with the lower nucleophilicity of primary versus secondary amines. Anilines can also be successfully coupled (**5t-5v**), however the yield is typically lower than with our previously reported conditions.³⁰ Overall, the process shows excellent functional group tolerance, including groups such as acetals, alcohols, nitriles, and sulfonamides.



Scheme 2: Scope of amines in the amination of boronic ester **1**. Reactions were conducted on a 0.5 mmol scale. Yields reported are of isolated material of the corresponding product.

We were keen to test the stereoselectivity of the reaction. However, attempts to use enantiomerically enriched boronic ester (**S**)-**1** only resulted in the formation of racemic product. To investigate this further, we stopped the reaction early allowing the isolation of unreacted boronic ester **1**. Chiral HPLC analysis confirmed that the boronic ester had not undergone racemisation under the reaction conditions. Furthermore, we explored whether the reaction is diastereoselective when coupling chiral amines (**5w**-**5z**). In each case, essentially a 1:1 d.r. of the respective product was observed. Attempts to use amines with larger substituents at the 2-position were unsuccessful, presumably due reaching the reaction's limit of steric congestion.

The stereoablative nature of the amination reaction is consistent with the formation of a radical intermediate. However, we cannot rule out at this stage the formation of an alkyl copper intermediate, which are thought to be configurationally unstable above $-50\text{ }^{\circ}\text{C}$.³¹ To probe the possibility of a radical intermediate, we performed the amination of cyclopropane **6** in a radical clock experiment (eq 1). This resulted in a mixture of two amines, amine **7** with the cyclopropane intact and amine **8** in which the cyclopropane ring has opened. This suggests a radical intermediate is formed, with recombination occurring on a similar timescale to cyclopropane ring opening.³²



While formation of alkyl radicals from alkylboronic esters has been previously reported, this typically requires prior formation of a boron ‘ate’ complex through coordination of an alkoxide, pyridine or aryl lithium.^{33–37} However, such an ‘ate’ complex was not observed by ¹H or ¹¹B NMR spectroscopy when analysing mixtures of boronic ester **1** with morpholine in the presence of isopropanol.²⁶ <See SI>

To investigate this further, we conducted a series of cyclic voltammetry experiments. Boronic ester **1** showed a high oxidation potential ($E_{on} > 1.44$ V) in MeCN (figure 1a).³⁸ <insert footnote: MeCN was used as a solvent to enable solubility of the electrolyte. When the standard reaction was performed using MeCN as solvent, amination was successful with amine **2** formed in 78% yield. See the supporting information for details.> The voltammogram of CuBr₂ does not change in the presence of the boronic ester, suggesting there is not a direct interaction between **1** and the Cu salt. However, cyclic voltammetry of a mixture of morpholine and **1** does show the formation of a new peak (oxidation potential of $E_{pc} = 1.25$ V; figure 1b). Given that this is greater than the oxidation potential of morpholine ($E_{pc} = 0.81$ V), we suggest this could arise from a new species generated upon interaction of **1** and a morpholine radical, which can be oxidised at a lower potential than the boronic ester without activation. This type of interaction was recently proposed by Speckmeier and Maier³⁹ as a way to generate alkyl radicals from alkyl boronic esters. While the mechanism of this oxidation is currently unclear, given the reduction potential of O₂ has been estimated as +1.21 V (in MeCN)⁴⁰ it should be feasible under aerobic conditions our reaction. Further investigations into elucidating this complex reaction mechanism are ongoing.






Figure 1: Cyclic voltammetry experiments, conducted in MeCN under N₂, of a) boronic ester **1** and CuBr₂, and b) boronic ester **1** and morpholine.


We have also explored the scope of reaction with respect to the boronic ester (scheme 3). A range of secondary boronic esters were successfully reacted with morpholine in good to excellent yield. Varying the electronics of the substitutes on the aryl ring of the boronic ester did not lead to great variation in terms of yield. However, the more electron donating MeO- (**10a**) and electron withdrawing CF₃- (**10f**) groups showed a small reduction in yield, in part to a small increase in corresponding alcohol and ketone side products observed in the crude reaction mixture. Importantly, aryl bromides and chlorides (**10e**, **10h**, **10i**) were tolerated, with no products from Ullmann-Goldberg amination observed. In addition, functional groups including azides, ethers and indole were also tolerated. Pleasingly, an alkyl chloride was also tolerated, with no products of amination through nucleophilic substitution observed in the reaction mixture (**10n**). A primary benzylic boronic ester was also successfully coupled in high yield (**10q**).




Benzylic Boronic Esters




Aliphatic Boronic Esters



Tertiary Boronic Esters



Synthesis of TRPV1 Antagonist 13:



Scheme 3: Scope of amination with respect to the boronic ester substrate. Reactions were conducted on a 0.5 mmol scale. Yields reported are of isolated material of the corresponding product.

We could extend the amination reaction to aliphatic boronic esters. In particular, primary boronic esters could be coupled successfully (**10r-10u**), though the efficiency of the reaction is slightly reduced compared with benzylic boronic esters.

Subjecting tertiary boronic esters **9v** and **9w** to our reaction conditions did lead to formation of a C-tertiary amines **10v** and **10w**, albeit in lower yield compared to our previous results when coupling anilines.³⁰ In reaction of **9v**, the corresponding alkene **11** was formed as the major product from the mixture.

Finally, we wanted to apply our amination reaction as the final step in the synthesis of a complex molecule, to simulate how the method could be used in a discovery chemistry scenario. We targeted amine **13**, a TRVP 1 inhibitor which is a potential treatment for chronic pain.⁴¹ Due to the poor solubility of **12** in most organic solvents, we required the use of DMSO as reaction solvent in which **12** was partially soluble. Despite this, and without further optimisation, amine **13** could be isolated successfully, and in a yield that would allow the material to be tested for its biological activity. We therefore believe our method could find application in settings such as medicinal chemistry, and allow chemists to develop structure-activity relationships through the coupling of a common advanced amine or boronic ester intermediates.

Conclusions:

In conclusion, we have developed conditions for the Cu-catalysed Chan-Lam amination of alkylboronic esters with aliphatic amines. This method expands the scope to the coupling of both primary and secondary amines, and uses oxygen from air as the terminal oxidant for the reaction. The reaction conditions tolerate a broad range of functional groups, and have been applied successfully to the synthesis of a TRVP 1 inhibitor. Mechanistic studies suggest that the reaction proceeds via an alkyl radical intermediate, generated from the boronic ester upon activation through an amino radical transfer process. The investigation of related Cu-catalysed oxidative couplings of alkylboron reagents is ongoing, and will be reported in due course.

Acknowledgements:

We thank the EPSRC (Grant EP/T009292/1), the Royal Society (Grant RSG\R1\180065) and the University of Sheffield for financial support.

References:

- (1) Heravi, M. M.; Kheilkordi, Z.; Zadsirjan, V.; Heydari, M.; Malmir, M. Buchwald-Hartwig Reaction: An Overview. *J. Organomet. Chem.* **2018**, *861*, 17–104. <https://doi.org/https://doi.org/10.1016/j.jorganchem.2018.02.023>.
- (2) Ruiz-Castillo, P.; Buchwald, S. L. Applications of Palladium-Catalyzed C–N Cross-Coupling Reactions. *Chem. Rev.* **2016**, *116* (19), 12564–12649. <https://doi.org/10.1021/acs.chemrev.6b00512>.
- (3) Beletskaya, I. P.; Cheprakov, A. V. The Complementary Competitors: Palladium and Copper in C–N Cross-Coupling Reactions. *Organometallics* **2012**, *31* (22), 7753–7808. <https://doi.org/10.1021/om300683c>.
- (4) Qiao, J. X.; Lam, P. Y. S. Copper-Promoted Carbon-Heteroatom Bond Cross-Coupling with Boronic Acids and Derivatives. *Synthesis (Stuttg)*. **2011**, No. 6, 829–856. <https://doi.org/10.1055/s-0030-1258379>.
- (5) Monnier, F.; Taillefer, M. Copper-Catalyzed C(Aryl)–N Bond Formation. In *Topics in Organometallic Chemistry*; Taillefer, M., Ma, D., Eds.; Springer Berlin Heidelberg: Berlin, Heidelberg, 2013; pp 173–204. https://doi.org/10.1007/3418_2013_69.
- (6) Munir, I.; Zahoor, A. F.; Rasool, N.; Naqvi, S. A. R.; Zia, K. M.; Ahmad, R. Synthetic Applications and

Methodology Development of Chan–Lam Coupling: A Review. *Mol. Divers.* **2019**, *23* (1), 215–259. <https://doi.org/10.1007/s11030-018-9870-z>.

- (7) West, M. J.; Fyfe, J. W. B.; Vantourout, J. C.; Watson, A. J. B. Mechanistic Development and Recent Applications of the Chan–Lam Amination. *Chem. Rev.* **2019**, *119* (24), 12491–12523. <https://doi.org/10.1021/acs.chemrev.9b00491>.
- (8) Sambigiato, C.; Marsden, S. P.; Blacker, A. J.; McGowan, P. C. Copper Catalysed Ullmann Type Chemistry: From Mechanistic Aspects to Modern Development. *Chem. Soc. Rev.* **2014**, *43* (10), 3525–3550. <https://doi.org/10.1039/C3CS60289C>.
- (9) Monnier, F.; Taillefer, M. Catalytic C–C, C–N, and C–O Ullmann-Type Coupling Reactions. *Angew. Chemie Int. Ed.* **2009**, *48* (38), 6954–6971. <https://doi.org/10.1002/anie.200804497>.
- (10) Trowbridge, A.; Walton, S. M.; Gaunt, M. J. New Strategies for the Transition-Metal Catalyzed Synthesis of Aliphatic Amines. *Chem. Rev.* **2020**, *120* (5), 2613–2692. <https://doi.org/10.1021/acs.chemrev.9b00462>.
- (11) Abdel-Magid, A. F.; Mehrman, S. J. A Review on the Use of Sodium Triacetoxyborohydride in the Reductive Amination of Ketones and Aldehydes. *Org. Process Res. Dev.* **2006**, *10* (5), 971–1031. <https://doi.org/10.1021/op0601013>.
- (12) Afanasyev, O. I.; Kuchuk, E.; Usanov, D. L.; Chusov, D. Reductive Amination in the Synthesis of Pharmaceuticals. *Chem. Rev.* **2019**, *119* (23), 11857–11911. <https://doi.org/10.1021/acs.chemrev.9b00383>.
- (13) Cabré, A.; Verdaguer, X.; Riera, A. Recent Advances in the Enantioselective Synthesis of Chiral Amines via Transition Metal-Catalyzed Asymmetric Hydrogenation. *Chem. Rev.* **2022**, *122* (1), 269–339. <https://doi.org/10.1021/acs.chemrev.1c00496>.
- (14) Patil, M. D.; Grogan, G.; Bommarius, A.; Yun, H. Oxidoreductase-Catalyzed Synthesis of Chiral Amines. *ACS Catal.* **2018**, *8* (12), 10985–11015. <https://doi.org/10.1021/acscatal.8b02924>.
- (15) Simić, S.; Zukić, E.; Schmermund, L.; Faber, K.; Winkler, C. K.; Kroutil, W. Shortening Synthetic Routes to Small Molecule Active Pharmaceutical Ingredients Employing Biocatalytic Methods. *Chem. Rev.* **2022**, *122* (1), 1052–1126. <https://doi.org/10.1021/acs.chemrev.1c00574>.
- (16) Sangster, J. J.; Marshall, J. R.; Turner, N. J.; Mangas-Sanchez, J. New Trends and Future Opportunities in the Enzymatic Formation of C–C, C–N, and C–O Bonds. *ChemBioChem* **2022**, *23* (6), e202100464. <https://doi.org/https://doi.org/10.1002/cbic.202100464>.
- (17) Campos, K. R.; Coleman, P. J.; Alvarez, J. C.; Dreher, S. D.; Garbaccio, R. M.; Terrett, N. K.; Tillyer, R. D.; Truppo, M. D.; Parmee, E. R. The Importance of Synthetic Chemistry in the Pharmaceutical Industry. *Science (80-)*. **2019**, *363* (6424), eaat0805. <https://doi.org/10.1126/science.aat0805>.
- (18) Grayson, J. D.; Dennis, F. M.; Robertson, C. C.; Partridge, B. M. Chan–Lam Amination of Secondary and Tertiary Benzylic Boronic Esters. *J. Org. Chem.* **2021**, *86* (14), 9883–9897. <https://doi.org/10.1021/acs.joc.1c00976>.
- (19) Mori-Quiroz, L. M.; Shimkin, K. W.; Rezazadeh, S.; Kozłowski, R. A.; Watson, D. A. Copper-Catalyzed Amidation of Primary and Secondary Alkyl Boronic Esters. *Chem. - A Eur. J.* **2016**, *22* (44), 15654–15658. <https://doi.org/10.1002/chem.201603677>.
- (20) Racine, E.; Monnier, F.; Vors, J.-P.; Taillefer, M. Direct N-Cyclopropylation of Secondary Acyclic Amides Promoted by Copper. *Chem. Commun.* **2013**, *49* (67), 7412–7414. <https://doi.org/10.1039/C3CC42575D>.
- (21) Sueki, S.; Kuninobu, Y. Copper-Catalyzed N - and O -Alkylation of Amines and Phenols Using Alkylborane Reagents. *Org. Lett.* **2013**, *15* (7), 1544–1547. <https://doi.org/10.1021/ol400323z>.

- (22) Kim, D. S.; Lee, H. G. Formation of the Tertiary Sulfonamide C(Sp³)–N Bond Using Alkyl Boronic Ester via Intramolecular and Intermolecular Copper-Catalyzed Oxidative Cross-Coupling. *J. Org. Chem.* **2021**, *86* (23), 17380–17394. <https://doi.org/10.1021/acs.joc.1c01759>.
- (23) Bénard, S.; Neuville, L.; Zhu, J. Copper-Promoted N-Cyclopropylation of Anilines and Amines by Cyclopropylboronic Acid. *Chem. Commun.* **2010**, *46* (19), 3393–3395. <https://doi.org/10.1039/B925499D>.
- (24) Singh, K.; Kumar, M.; Pavadai, E.; Naran, K.; Warner, D. F.; Ruminski, P. G.; Chibale, K. Synthesis of New Verapamil Analogues and Their Evaluation in Combination with Rifampicin against Mycobacterium Tuberculosis and Molecular Docking Studies in the Binding Site of Efflux Protein Rv1258c. *Bioorg. Med. Chem. Lett.* **2014**, *24* (14), 2985–2990. <https://doi.org/https://doi.org/10.1016/j.bmcl.2014.05.022>.
- (25) <add Tex>.
- (26) Footnote 1.
- (27) Grayson, J. D.; M. Partridge, B. Mild Cu-Catalyzed Oxidation of Benzylic Boronic Esters to Ketones. *ACS Catal.* **2019**, *9* (5), 4296–4301. <https://doi.org/10.1021/acscatal.9b00992>.
- (28) Meanwell, N. A. Fluorine and Fluorinated Motifs in the Design and Application of Bioisosteres for Drug Design. *J. Med. Chem.* **2018**, *61* (14), 5822–5880. <https://doi.org/10.1021/acs.jmedchem.7b01788>.
- (29) Zask, A.; Kaplan, J.; Verheijen, J. C.; Richard, D. J.; Curran, K.; Brooijmans, N.; Bennett, E. M.; Toral-Barza, L.; Hollander, I.; Ayrál-Kaloustian, S.; Yu, K. Morpholine Derivatives Greatly Enhance the Selectivity of Mammalian Target of Rapamycin (MTOR) Inhibitors. *J. Med. Chem.* **2009**, *52* (24), 7942–7945. <https://doi.org/10.1021/jm901415x>.
- (30) Grayson, J. D.; Dennis, F. M.; Robertson, C. C.; Partridge, B. M. Chan–Lam Amination of Secondary and Tertiary Benzylic Boronic Esters. *J. Org. Chem.* **2021**, *86* (14), 9883–9897. <https://doi.org/10.1021/acs.joc.1c00976>.
- (31) Skotnitzki, J.; Morozova, V.; Knochel, P. Diastereoselective Copper-Mediated Cross-Couplings between Stereodefined Secondary Alkylcoppers with Bromoalkynes. *Org. Lett.* **2018**, *20* (8), 2365–2368. <https://doi.org/10.1021/acs.orglett.8b00699>.
- (32) Halgren, T. A.; Roberts, J. D.; Horner, J. H.; Martinez, F. N.; Tronche, C.; Newcomb, M. Kinetics and Equilibrium Constants for Reactions of α -Phenyl-Substituted Cyclopropylcarbinyl Radicals. *J. Am. Chem. Soc.* **2000**, *122* (13), 2988–2994. <https://doi.org/10.1021/ja991114h>.
- (33) Lima, F.; Kabeshov, M. A.; Tran, D. N.; Battilocchio, C.; Sedelmeier, J.; Sedelmeier, G.; Schenkel, B.; Ley, S. V. Visible Light Activation of Boronic Esters Enables Efficient Photoredox C(Sp²)–C(Sp³) Cross-Couplings in Flow. *Angew. Chemie Int. Ed.* **2016**, *55* (45), 14085–14089. <https://doi.org/10.1002/anie.201605548>.
- (34) Lennox, A. J. J.; Nutting, J. E.; Stahl, S. S. Selective Electrochemical Generation of Benzylic Radicals Enabled by Ferrocene-Based Electron-Transfer Mediators. *Chem. Sci.* **2018**, *9* (2), 356–361. <https://doi.org/10.1039/C7SC04032F>.
- (35) Corcé, V.; Ollivier, C.; Fensterbank, L. Boron{ } Silicon{ } Nitrogen and Sulfur-Based Contemporary Precursors for the Generation of Alkyl Radicals by Single Electron Transfer and Their Synthetic Utilization. *Chem. Soc. Rev.* **2022**, *51* (4), 1470–1510. <https://doi.org/10.1039/D1CS01084K>.
- (36) Shu, C.; Noble, A.; Aggarwal, V. K. Photoredox-Catalyzed Cyclobutane Synthesis by a Deboronative Radical Addition–Polar Cyclization Cascade. *Angew. Chemie Int. Ed.* **2019**, *58* (12), 3870–3874. <https://doi.org/https://doi.org/10.1002/anie.201813917>.
- (37) Pillitteri, S.; Ranjan, P.; Van der Eycken, E. V.; Sharma, U. K. Uncovering the Potential of Boronic Acid and Derivatives as Radical Source in Photo(Electro)Chemical Reactions. *Adv. Synth. Catal.* **2022**, *n/a*

(n/a). <https://doi.org/https://doi.org/10.1002/adsc.202200204>.

- (38) Footnote 2.
- (39) Speckmeier, E.; Maier, T. C. ART—An Amino Radical Transfer Strategy for C(Sp²)–C(Sp³) Coupling Reactions, Enabled by Dual Photo/Nickel Catalysis. *J. Am. Chem. Soc.* **2022**. <https://doi.org/10.1021/jacs.2c03220>.
- (40) Pegis, M. L.; Roberts, J. A. S.; Wasylenko, D. J.; Mader, E. A.; Appel, A. M.; Mayer, J. M. Standard Reduction Potentials for Oxygen and Carbon Dioxide Couples in Acetonitrile and N,N-Dimethylformamide. *Inorg. Chem.* **2015**, *54* (24), 11883–11888. <https://doi.org/10.1021/acs.inorgchem.5b02136>.
- (41) Wang, H.-L.; Katon, J.; Balan, C.; Bannon, A. W.; Bernard, C.; Doherty, E. M.; Dominguez, C.; Gavva, N. R.; Gore, V.; Ma, V.; Nishimura, N.; Surapaneni, S.; Tang, P.; Tamir, R.; Thiel, O.; Treanor, J. J. S.; Norman, M. H. Novel Vanilloid Receptor-1 Antagonists: 3. The Identification of a Second-Generation Clinical Candidate with Improved Physicochemical and Pharmacokinetic Properties. *J. Med. Chem.* **2007**, *50* (15), 3528–3539. <https://doi.org/10.1021/jm070191h>.