

# Mechanochemical Protocol Facilitates the Generation of Arylmanganese Nucleophiles from Unactivated Manganese Metal

Rina Takahashi,<sup>[a]</sup> Pan Gao,<sup>[a,b]</sup> Koji Kubota,<sup>[a,b]\*</sup> and Hajime Ito<sup>[a,b]\*</sup>

**Abstract:** We report a mechanochemical ball-milling protocol facilitates the generation of various arylmanganese nucleophiles from aryl halides and commercially available manganese metal without the need for complicated pre-activation processes. These manganese-based carbon nucleophiles can be used directly for one-pot addition reactions with various electrophiles and palladium-catalyzed cross-coupling reactions under bulk-solvent-free mechanochemical conditions. Importantly, all experimental operations can be conducted under atmospheric conditions.


Organomanganese reagents play a unique role in organic synthetic chemistry because of their mild reactivity, distinctive chemoselectivity, and low toxicity.<sup>[1]</sup> Most organomanganese reagents are typically prepared via the following methods: (1) the insertion of manganese metal into a carbon–halogen bond (direct synthesis);<sup>[2–4]</sup> (2) metathesis between an organolithium or organomagnesium reagent and a manganese halide;<sup>[5]</sup> or (3) a directed deprotonation using  $\text{TMP}_2\text{Mn}\cdot 2\text{LiCl}$ .<sup>[6]</sup> Given the ready availability of manganese metal, and the mild reaction conditions and high atom efficiency associated with direct synthesis, this method can be expected to afford convenient access to versatile organomanganese reagents. Although manganese metal has a stronger reduction potential than zinc, direct synthesis is only applicable to highly reactive organic halides, such as allylic halides or  $\alpha$ -haloesters when commercial manganese metal is used.<sup>[2,3]</sup> One of the main reasons for this low reactivity is the presence of a usually tightly bound oxide layer on its surface.<sup>[3a]</sup> Therefore, the preparation of activated manganese (Rieke manganese) using strong reducing reagents, such as alkali metals, is often required when aryl halides are used as substrates (Scheme 1a).<sup>[4]</sup> Unfortunately, this greatly reduces the practical utility of arylmanganese nucleophiles. Recently, Knochel et. al. have reported that the direct synthesis of arylmanganese reagents from unactivated manganese metal can be achieved using  $\text{LiCl}$ ,  $\text{PbCl}_2$ , and  $\text{InCl}_3$  additives (Scheme 1b).<sup>[3b]</sup> Although this achievement is remarkable, this solution-based approach still requires multiple metal additives, long reaction times (24 h), the use of dry organic solvents, and inert-gas-line techniques. All of which represent considerable drawbacks from both an environmental and an economic perspective. Therefore, the development of an operationally simple protocol that does not

require pre-activation of the reagents would be highly desirable as it can be expected to advance the widespread application of arylmanganese nucleophiles in organic synthesis.


Recently, a mechanochemical synthetic technique that uses ball milling has attracted significant attention as a new tool for organic reactions.<sup>[7]</sup> The advantages of mechanochemical synthesis include avoiding the use of potentially harmful organic solvents, shorter reaction times, and operational simplicity. Moreover, recent studies have revealed that strong mechanical agitation during the ball-milling process can activate zero-valent metals in the bulk state, facilitating surface reactions with organic halides.<sup>[8]</sup> For example, Browne et. al. have reported the direct synthesis of organozinc reagents from metallic zinc facilitated by ball milling.<sup>[9]</sup> The groups of Harrowfield, Birke, Hanusa, Bolm, and our own group have independently reported the mechanochemical synthesis of Grignard reagents via the direct insertion of magnesium metal into organic halides.<sup>[10,11]</sup> More recently, we have demonstrated that a ball-milling approach allows the generation of heavy Grignard reagents based on calcium from aryl halides and unactivated calcium metal.<sup>[12]</sup> Notably, these protocols do not involve the pre-activation of the metals and are applicable to a broad range of substrates.

**Scheme 1.** Direct insertion of manganese metal into carbon–halogen bonds.


(a) Preparation of Rieke Mn via reaction with strong reducing reagents




(b) Use of  $\text{LiCl}$ ,  $\text{PbCl}_2$  and  $\text{InCl}_3$  as activating reagents



(c) **This work:** mechanochemical approach using ball milling



(d) Related study: manganese-mediated dimerization of arylidene malonates



These achievements led us to consider the mechanochemical generation of organomanganese reagents via the direct synthesis from unactivated manganese metal, which is

[a] R. Takahashi, Dr. P. Gao, Prof. Dr. K. Kubota, Prof. Dr. H. Ito  
Division of Applied Chemistry and Frontier Chemistry Center,  
Faculty of Engineering, Hokkaido University, Sapporo,  
Hokkaido 060-8628 (Japan)

[b] Dr. P. Gao, Prof. Dr. K. Kubota, Prof. Dr. H. Ito  
Institute for Chemical Reaction Design and Discovery  
WPI-ICReDD, Hokkaido University, Sapporo, Hokkaido,  
060-8628 (Japan)  
E-mail: hajito@eng.hokudai.ac.jp

Supporting information for this article is given via a link at the end of the document.


barely reactive in solution (Scheme 1c). We envisaged that the mechanical impact of ball milling could remove the unreactive oxide layer covering the surface of commercially available manganese metal, thus facilitating the direct insertion of manganese metal into organic halide bonds without the need for a complicated pre-activation process.<sup>[4]</sup> Whilst our study was already underway, Browne et. al. reported that commercial manganese metal can be activated using ball-milling to realize the reductive dimerization of arylidene malonates (Scheme 1d).<sup>[13]</sup> However, in that report, the direct synthesis of organomanganese nucleophiles from alkyl halides and the subsequent nucleophilic addition to electrophiles was unsuccessful whilst other organic halides were not explored.<sup>[13]</sup> Herein, we report the first successful example of the direct synthesis of arylmanganese nucleophiles from unactivated manganese metal and aryl halides using mechanochemistry. These reactions are highly efficient and went to completion within 3 hours to form the desired arylmanganese nucleophiles. These nucleophiles were then reacted with various electrophiles under mechanochemical conditions. The direct synthesis of alkylmanganese nucleophiles was also achieved using this protocol, although there is room to improve the efficiency. Notably, the entire synthetic procedure can be conducted under atmospheric conditions without using any special precautions or synthetic techniques.

We began by attempting the preparation of various arylmanganese species from different aryl iodides (**1**) (1.0 equiv) and commercial manganese metal (Wako, powder, >98% purity) (3.0 equiv), using a Retch MM400 mixer mill (5 mL stainless-steel milling jar with a 10-mm-diameter stainless-steel ball), followed by the subsequent nucleophilic addition to an aldehyde **2a** (1.05 equiv) (Table 1). Based on a previous study on the isolation of THF-coordinated arylmanganese species by Westerhausen et. al.,<sup>[14]</sup> 2.0 equivalents of tetrahydrofuran (THF) were used to facilitate the formation of the organomanganese nucleophile. Unfortunately, aryl iodides **1a–1f**, which bear electron-donating and -withdrawing groups, did not react under the applied conditions. However, the substrate bearing a fluorine atom at the *ortho* position (**1g**) afforded the desired product **3g** in 9% yield. No reaction occurred when the corresponding aryl bromide (**1h**) was used. We found that the reaction of pentafluorophenyl iodide (**1i**) underwent direct metalation to form the corresponding addition product (**3i**) in 35% yield. These results suggest that only *ortho*-halogenated aryl iodides are reactive in the direct manganese metalation under mechanochemical conditions, a finding which is consistent with the results of Knochel's previously reported solution-based method.<sup>[3b]</sup> The use of activated organic halides such as benzyl bromide only provided the homocoupling products. This is probably due to a facile radical-radical coupling or the nucleophilic substitution of the substrate by an *in-situ*-generated benzylmanganese nucleophile.

Next, we proceeded to optimize the reaction conditions using pentafluorophenyl iodide (**1i**) as the substrate and 2.0 equivalent of **2a** as a trapping reagent (Table 2). The reaction without the addition of an ether additive provided a trace amount of **3i** (2% yield; entry 1). The addition of THF significantly promoted the generation of the arylmanganese nucleophile and **3i** was obtained in 54% yield (entry 2). After screening further ether additives (entries 3–8), we found that the reaction with methyl *tert*-butyl ether (MTBE) gave the highest yield (75% yield; entry 3). Next, the mechanochemical conditions for the synthesis of **3g** from **1g** were optimized (entries 9–11). While **1g** showed much lower reactivity toward the metalation than **1i**, the reaction in the presence of THF gave the corresponding addition product


(**3g**) in 25% yield (entry 9). Prolonging the reaction time for the metalation step led to a slightly improved yield (30% yield; entry 10). Interestingly, the addition of MTBE, which was effective for the reaction of **1i**, did not produce **3g** (<1%; entry 11). Next, 1,3-difluoro-2-iodobenzene (**1j**) was used as a substrate (entries 12–14). While **1j** also showed lower reactivity than **1i**, the reaction with manganese metal in the presence of THF for 3 hours afforded the desired product (**3j**) in 69% yield (entry 13). Again, the use of MTBE did not lead to the formation of **3j**, indicating that MTBE is specifically effective for the highly electron-deficient pentafluorophenyl iodide **1i**.

**Table 1.** Initial attempts to generate arylmanganese nucleophiles under mechanochemical conditions.<sup>[a]</sup>



[a] Conditions: **1** (1.0 mmol), Mn powder (3.0 mmol), THF (2.0 mmol), and **2a** (1.05 mmol) in a stainless-steel milling jar (5 mL) with a stainless-steel ball (10 mm).

**Table 2.** Optimization of the reaction conditions.<sup>[a]</sup>




Entry	Substrate	Additive	Time [h]	Yield [%] <sup>[b]</sup>
1	<b>1i</b>	-	1.5	2
2	<b>1i</b>	THF	1.5	54
3	<b>1i</b>	MTBE	1.5	75 (54)
4	<b>1i</b>	anisole	1.5	66
5	<b>1i</b>	Et <sub>2</sub> O	1.5	64
6	<b>1i</b>	CPME	1.5	57
7	<b>1i</b>	1,2-DME	1.5	40
8	<b>1i</b>	1,4-dioxane	1.5	10
9	<b>1g</b>	THF	1.5	25

10	<b>1g</b>	THF	3	30 (25)
11	<b>1g</b>	MTBE	3	<1
12	<b>1j</b>	THF	1.5	25
13	<b>1j</b>	THF	3	69 (68)
14	<b>1j</b>	MTBE	1.5	<1

[a] Conditions: **1** (1.0 mmol), Mn powder (3.0 mmol), THF (2.0 mmol), and **2a** (2.0 mmol) in a stainless-steel milling jar (5 mL) with a stainless-steel ball (10 mm). [b] Determined via  $^{19}\text{F}$  NMR analysis with an internal standard. Isolated yields are shown in parenthesis.

With the optimized conditions for the generation of the arylmanganese reagents in hand, a series of one-pot mechanochemical reactions with various electrophiles were evaluated (Scheme 2). Following the reaction between **1i** and the unactivated manganese metal, benzaldehyde (**2b**) was added to the mixture and subsequent ball milling afforded the desired addition product (**3k**) in 60% yield. The reaction with cyclohexanone (**2c**) also produced the corresponding product (**4**) in 45% yield. In addition, we found that phenyl isocyanate (**5a**) efficiently reacted with the mechanochemically generated arylmanganese nucleophile to give the desired amide (**6a**) in 69% yield. Furthermore, a mechanochemical palladium-catalyzed cross-coupling reaction between aryl iodide and the arylmanganese nucleophile in the presence of a Pd-PEPPSI-*i*Pr catalyst<sup>[15,16]</sup> furnished the corresponding biaryl compound (**7**) in 71% yield under high-temperature ball-milling conditions using a heat gun (for details, see the Supporting Information).<sup>[17]</sup>

**Scheme 2.** The use of mechanochemically generated arylmanganese nucleophiles for reactions with various electrophiles.<sup>[a,b]</sup>




[a] For the reaction conditions, please see the Supporting Information. [b] Isolated yields are shown. NMR yields are shown in parenthesis.

Next, the scope of aryl halides that can be used for the mechanochemical generation of arylmanganese nucleophiles was investigated (Table 3). Cyclohexyl isocyanate (**5b**) was used as a trapping reagent and the reactions of several fluorinated aryl iodides (**1g**, **1i–1k**) under the optimized conditions proceeded

smoothly to give the corresponding amides (**6b–6e**) in moderate to good yields (38–79%). Polyfluorinated substrates bearing methoxy, dimethylamino, or siloxy groups (**1l–1n**) were successfully converted into the desired products (**6f–6h**) in 28–70% yield. The iodide moiety of 5-bromo-1,3-difluoro-2-iodobenzene (**1o**) selectively reacted with manganese metal to furnish the desired product (**6i**) in 75% yield. We found that 1,3-dichloro-2-iodobenzene (**1p**) also reacted under the applied conditions to furnish **6j** in 55% yield. The reaction of 1-bromo-3-fluoro-2-iodobenzene (**1q**) produced the desired amide (**6k**) in 70% yield. In the case of 1,3-dibromo-2-iodobenzene (**1r**), the desired product (**6m**) was obtained, albeit that the bromide moiety also reacted to form side product **6n**. In addition to aryl iodides, electron-deficient polyfluorinated aryl bromides (**1s–1v**) could be used as substrates to afford the corresponding amides 39–69% yield. Even though the aryl-halide substrate scope of our reaction is limited compared to that of the Rieke method,<sup>[4]</sup> it is comparable to that of Knochel's conditions.<sup>[3b]</sup>


**Table 3.** Scope of aryl iodides and bromides.<sup>[a,b]</sup>





[a] For the reaction conditions, please see the Supporting Information. [b] Isolated yields are shown.

The direct formation of alkylmanganese nucleophiles from unactivated alkyl halides can be accomplished using Rieke manganese.<sup>[4d]</sup> We found that the mechanochemical reaction of alkyl iodide **8** with commercial manganese metal, followed by reaction with **5b** formed the desired product (**9**) in 17% yield (Scheme 3). Although the yield needs to be improved, this is the first example of the direct generation of alkylmanganese nucleophiles from unactivated alkyl halides and commercial, unactivated manganese metal.

**Scheme 3.** Reaction of alkyl iodide **8** and manganese metal under mechanochemical conditions.



We conducted a number of nucleophilic addition reactions with **5b** after exposing the mechanochemically generated arylmanganese nucleophile **1i** to air for different lengths of time (10–60 min; Figure 1). The results showed that, even though there were no obvious changes in its appearance, the yield of **6b** decreased when the generated arylmanganese species were exposed to air for 10 min or more (Figure 1). Although pentafluorobenzene was detected in the <sup>1</sup>H NMR spectrum after the exposure of the manganese nucleophile to air, byproducts derived from oxygen or carbon dioxide, such as phenol and benzoic acid, were not detected. These results suggest that the decreased yield is most likely caused by protonation resulting from exposure to moisture in the air.




**Figure 1.** Stability of organomanganese reagents in air and the appearance of the crude mixture.

In order to confirm that the arylmanganese nucleophiles are indeed generated under our mechanochemical conditions, preliminary mechanistic studies were performed (Scheme 4). First, we conducted a deuteration experiment (Scheme 4a) wherein the mechanochemically formed arylmanganese nucleophile was quenched by the addition of CD<sub>3</sub>CO<sub>2</sub>D (10 equiv). The

corresponding deuterated product (**10**) was obtained in a high deuteration ratio (95% D), suggesting that an arylmanganese nucleophile was most likely generated under mechanochemical conditions. Next, in order to test the hypothesis that transition-metals, such as iron and chromium, which could potentially leach from the stainless-steel milling jar and balls, catalyze the direct metalation or subsequent nucleophilic addition, we investigated the reaction using jars made of different materials such as tungsten carbide (WC) or zirconium oxide (Scheme 4b). Although the use of a 5 mL jar is optimal for this scale, we compared the reactions with 10 mL jars and 10 mm balls without changing the reaction scale because 10 mL is the minimum volume for the commercially available jars made of these materials. The results confirmed that the direct metalation and subsequent nucleophilic addition to **5b** proceeded regardless of the material of the milling jars and balls. Thus, we excluded the possibility that the reaction is catalyzed by metals leached from the ball milling equipment (Scheme 4b). Finally, we assessed the use of unactivated manganese metal in solution (Scheme 4c). When we conducted the reaction between unactivated manganese metal and **1i** in THF (1.0 M) at room temperature, the corresponding amide (**6b**) was not detected and a large amount of unreacted **1i** was recovered (93%). This result indicates that mechanical activation of the manganese metal via ball milling is essential for the generation of arylmanganese nucleophiles. When Knochel's method was applied to the direct metalation of **1i**, **6b** was obtained in 83% yield.<sup>[3b]</sup> This result suggests that manganese-based carbon nucleophiles (Ar–MnX) similar to those formed under Knochel's conditions are also most likely formed under the mechanochemical conditions.


#### Scheme 4. Mechanistic studies.<sup>[a]</sup>




[a] For the reaction conditions, please see the Supporting Information.

To validate our hypothesis that the mechanical impact provided by ball milling can reduce or remove the unreactive oxide layer covering the surface of the commercial manganese metal, we analyzed its surface using X-ray photoelectron spectroscopy (XPS) (Figure 2). When commercial manganese metal used in this study was subjected to XPS analysis we found that a signal derived from the Mn-O bonds was detected on its surface (Figure 2a). The manganese oxide content can be estimated based on the Mn/O atom ratio.<sup>[3a]</sup> The XPS analysis of the ball-milled sample revealed that the Mn ratio on the surface was increased after ball milling (Figure 2b) (for details, see the Supporting Information). This result suggests that ball milling can increase the reactive surface area of manganese metal via mechanical removal of the oxide layer, thus supporting our hypothesis.

(a) Commercial manganese metal (Wako, >98%)



(b) Ball-milled manganese metal



**Figure 2.** X-Ray photoelectron spectra of commercial manganese metal (a) before ball milling and (b) after ball milling.

In summary, we have demonstrated that a mechanochemical protocol using ball milling, that does not require complicated pre-activation processes, allows the generation of arylmanganese nucleophiles via the reaction between an aryl halide and commercial manganese metal. The mechanochemically generated arylmanganese reagents can be used as carbon nucleophiles for the one-pot addition to various electrophiles and palladium-catalyzed cross-coupling reactions under bulk-solvent-free ball-milling conditions. Notably, all synthetic operations can be conducted under atmospheric conditions. We therefore anticipate that this work will provide a new platform for the rapid discovery of novel organic transformations mediated by organomanganese species.

#### Acknowledgements

This work was financially supported by the Japan Society for the Promotion of Science (JSPS) via KAKENHI grants 22H0031802 and 21H01926, and 22H05328, by the JST via CREST grant JPMJCR19R1, by FOREST grant JPMJFR2011, and by the Institute for Chemical Reaction Design and Discovery (ICReDD) established by the World Premier International Research Initiative (WPI), MEXT, Japan.


**Keywords:** Manganese • Ball milling • Mechanochemistry • Solid state • Synthetic methods

- [1] For selected reviews and examples on the use of organomanganese reagents in organic synthesis, see: a) G. Cahiez, C. Duplais, J. Buendia, *Chem. Rev.* **2009**, *109*, 1434–1476; b) G. Cahiez, E. Métails, *Tetrahedron Asymmetry* **1997**, *8*, 1373–1376; c) G. Cahiez, E. Métails, *Tetrahedron Lett.* **1995**, *36*, 6449–6452.
- [2] For the direct synthesis of organomanganese reagents from commercial manganese metal and reactive organic halides without pre-activation, see: G. Cahiez, P.-Y. Chavant, *Tetrahedron Lett.* **1989**, *30*, 7373–7376.
- [3] For examples on the direct synthesis of organomanganese reagents from commercial manganese metal with activating reagents, see: a) K. Takai, T. Ueda, T. Hayashi, T. Moriwake, *Tetrahedron Lett.* **1996**, *37*, 7049–7052; b) Z. Peng, P. Knochel, *Org. Lett.* **2011**, *13*, 3198–3201.
- [4] For examples of the direct synthesis of organomanganese reagents from activated manganese metal, see: a) T. Hiyama, M. Obayashi, A. Nakamura, *Organometallics* **1982**, *1*, 1249–1251; b) R. D. Rieke, S.-H. Kim, X. Wu, *J. Org. Chem.* **1997**, *62*, 6921–6927; c) A. Fürstner, H. Brunner, *Tetrahedron Lett.* **1996**, *37*, 7009–7012; d) S.-H. Kim, M. V. Hanson, R. D. Rieke, *Tetrahedron Lett.* **1996**, *37*, 2197–2200; e) H. Kakiya, S. Nishimae, H. Shinokubo, K. Oshima, *Tetrahedron* **2001**, *57*, 8807–8815; f) J. Tang, H. Shinokubo, K. Oshima, *Synlett* **1998**, 1075–1076.
- [5] For examples of the synthesis of organomanganese reagents via metathesis with carbon nucleophiles, see: a) G. Cahiez, D. Bernard, J. F. Normant, *Synthesis* **1977**, 130–133; b) G. Cahiez, B. Laboue, *Tetrahedron Lett.* **1989**, *30*, 3545–3546; c) G. Friour, G. Cahiez, J. F. Normant, *Synthesis* **1984**, 37–40; d) G. Cahiez, D. Luart, F. Lecomte, *Org. Lett.* **2004**, *6*, 4395–4398; e) Y. Ahn, W. W. Doubleday, T. Cohen, *Synth. Commun.* **1995**, *25*, 33–41; f) I. Klement, H. Stadtmüller, P. Knochel, G. Cahiez, *Tetrahedron Lett.* **1997**, *38*, 1927–1930; g) K. Ritter, M. Hanack, *Tetrahedron Lett.* **1985**, *26*, 1285–1288; h) G. Cahiez, B. Figadere, *Tetrahedron Lett.* **1986**, *26*, 4445–4448; i) M. T. Reetz, K. Rölfing, N. Griebenow, *Tetrahedron Lett.* **1994**, *35*, 1969–1972; j) G. Cahiez, L. Razafintsalama, B. Laboue, F. Chau, *Tetrahedron Lett.* **1998**, *39*, 849–852; k) G. Cahiez, B. Laboue, *Tetrahedron Lett.* **1989**, *30*, 7369–7372; l) C. Boucley, G. Cahiez, S. Carini, V. Cerè, M. Comes-Franchini, P. Knochel, S. Pollicino, A. Ricci, *J. Organomet. Chem.* **2001**, *624*, 223–228.
- [6] S. H. Wunderlich, M. Kienle, P. Knochel, *Angew. Chem., Int. Ed.* **2009**, *48*, 7256–7260; *Angew. Chem.* **2009**, *121*, 7392–7396.
- [7] For selected reviews on the use of ball milling for organic synthesis, see: a) S. L. James, C. J. Adams, C. Bolm, D. Braga, P. Collier, T. Friščić, F. Grepioni, K. D. M. Harris, G. Hyett, W. Jones, A. Krebs, J. Mack, L. Maini, A. G. Orpen, I. P. Parkin, W. C. Shearouse, J. W. Steed, D. C. Waddell, *Chem. Soc. Rev.* **2012**, *41*, 413–447; b) G.-W. Wang, *Chem. Soc. Rev.* **2013**, *42*, 7668–7700; c) J.-L. Do, T. Friščić, *ACS Cent. Sci.* **2017**, *3*, 13–19; d) J. G. Hernández, C. Bolm, *J. Org. Chem.* **2017**, *82*, 4007–4019; e) T.-X. Métro, J. Martinez, F. Lamaty, *ACS Sustainable Chem. Eng.* **2017**, *5*, 9599–9602; f) T. K. Achar, A. Bose, P. Mal, *Beilstein J. Org. Chem.* **2017**, *13*, 1907–1931; g) O. Eguagie, J. S. Vyle, P. F. Conlon, M. A. Gilea, Y. Liang, *Beilstein J. Org. Chem.* **2018**, *14*, 955–970; h) J. L. Howard, Q. Cao, D. L. Browne, *Chem. Sci.* **2018**, *9*, 3080–3094; i) J. Andersen, J. Mack, *Green Chem.* **2018**, *20*, 1435–1443; j) C. Bolm, J. G. Hernández, *Angew. Chem., Int. Ed.* **2019**, *58*, 3285–3299; *Angew. Chem.* **2019**, *131*, 3320–3335; k) T. Friščić, C. Mottillo, H. M. Titi, *Angew. Chem., Int. Ed.* **2020**, *59*, 1018–1029; *Angew. Chem.* **2020**, *132*, 1030–1041; l) K. Kubota, H. Ito, *Trends Chem.* **2020**, *2*, 1066–1081; j) A. Porcheddu, E. Colacino, L. De Luca, F. Delogu, *ACS Catal.* **2020**, *10*, 8344–8394; k) J. A. Leitch, D. L. Browne, *Chem.-Eur. J.* **2021**, *27*, 9721–9726.
- [8] A. C. Jones, J. A. Leitch, S. E. Raby-Buck, D. L. Browne, *Nat. Synth.* **2022**, in press. DOI: 10.1038/s44160-022-00106-4
- [9] a) Q. Cao, J. L. Howard, E. Wheatley, D. L. Browne, *Angew. Chem., Int. Ed.* **2018**, *57*, 11339–11343; *Angew. Chem.* **2018**, *130*, 11509–11513; b) Q. Cao, R. T. Stark, I. A. Fallis, D. L. Browne, *ChemSusChem* **2019**, *12*, 2554–2557; c) J. Yin, R. T. Stark, I. A. Fallis, D. L. Browne, *J. Org. Chem.* **2020**, *85*, 2347–2354.
- [10] a) J. M. Harrowfield, R. J. Hart, C. R. Whitaker, *Aust. J. Chem.* **2001**, *54*, 423–425; b) V. Birke, C. Schütt, H. Burmeier, W. K. L. Ruck, W. K. L. Fresenius *Environ. Bull.* **2011**, *20*, 2794–2805; c) I. R. Speight, T. P. Hanusa, *Molecules* **2020**, *25*, 570–578; d) R. Takahashi, A. Hu, P. Gao, Y. Gao, Y. Pang, T. Seo, J. Jiang, S. Maeda, H. Takaya, K. Kubota, H. Ito, *Nat. Commun.* **2021**, *12*, 6691; e) V. S. Pfennig, R. C. Villella, J. Nikodemus, C. Bolm, *Angew. Chem., Int. Ed.* **2022**, *61*, e202116514; *Angew. Chem.* **2022**, *134*, e202116514.
- [11] Yang, Dai, and co-workers have reported magnesium-mediated reductive radical homo-coupling reactions of polyhaloarenes using mechanochemistry; for details, see: H. Chen, J. Fan, Y. Fu, C.-L. Do-Thanh, X. Suo, T. Wang, I. Popovs, D.-E. Jiang, Y. Yuan, Z. Yang, S. Dai, *Adv. Mater.* **2021**, *33*, 2008685.
- [12] P. Gao, J. Jiang, S. Maeda, K. Kubota, H. Ito, *Angew. Chem., Int. Ed.* **2022**, in press. DOI: 10.1002/anie.202207118
- [13] W. I. Nicholson, J. L. Howard, G. Magri, A. C. Seastram, A. Khan, R. R. A. Bolt, L. C. Morrill, E. Richards, D. L. Browne, *Angew. Chem., Int. Ed.* **2021**, *60*, 23128–23133; *Angew. Chem.* **2021**, *133*, 23312–23317.
- [14] R. Fischer, H. Görls, M. Friedrich, M. Westerhausen, *J. Organomet. Chem.* **2009**, *694*, 1107–1111.
- [15] C. J. O'Brien, E. A. B. Kantchev, C. Valente, N. Hadei, G. A. Chass, A. Lough, A. C. Hopkinson, M. G. Organ, *Chem.-Eur. J.* **2006**, *12*, 4743–4748.
- [16] a) G. Cahiez, S. Marquis, *Tetrahedron Lett.* **1996**, *37*, 1773–1776; b) A. Leleu, Y. Fort, R. Schneider, *Adv. Synth. Catal.* **2006**, *348*, 1086–1092.
- [17] T. Seo, N. Toyoshima, K. Kubota, H. Ito, *J. Am. Chem. Soc.* **2021**, *143*, 6165–6175.

---

## Entry for the Table of Contents

---



Herein, we report a mechanochemical ball-milling protocol that facilitates the generation of various arylmanganese nucleophiles from aryl halides and commercially available manganese metal without the need for complicated pre-activation processes. Importantly, all experimental operations can be conducted under atmospheric conditions. These manganese-based carbon nucleophiles can be used directly for one-pot addition reactions with various electrophiles and palladium-catalyzed cross-coupling reactions under bulk-solvent-free mechanochemical conditions.

*Rina Takahashi, Pan Gao, Koji Kubota\*,  
and Hajime Ito\**

**Mechanochemical Protocol  
Facilitates the Generation of  
Arylmanganese Nucleophiles from  
Unactivated Manganese Metal**

---