

Polarizing agents beyond pentacene for efficient triplet dynamic nuclear polarization in glass matrices

Keita Sakamoto,^{†,§} Tomoyuki Hamachi,^{†,§} Katsuki Miyokawa,^{‡,§} Kenichiro Tateishi,[#] Tomohiro Uesaka,[#] Yuki Kurashige,^{*,‡} Nobuhiro Yanai^{*,†,||}

[†]Department of Applied Chemistry, Graduate School of Engineering, Kyushu University, 744 Moto-oka, Nishi-ku, Fukuoka 819-0395, Japan

[‡] Department of Chemistry, Graduate School of Science, Kyoto University, Oiwake-Cho, Kitashirakawa, Sakyo-Ku, Kyoto 606-8502, Japan

[#]Cluster for Pioneering Research, RIKEN, RIKEN Nishina Center for Accelerator-Based Science, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

^{||} JST-FOREST, Honcho 4-1-8, Kawaguchi, Saitama 332-0012, Japan

ABSTRACT: Triplet dynamic nuclear polarization (triplet-DNP) is a technique that can obtain high nuclear polarization under moderate conditions. However, in order to obtain practically useful polarization, large single crystals doped with a polarizing agent must be strictly oriented with respect to the magnetic field to sharpen the electron spin resonance (ESR) spectra, which is a fatal problem that prevents its application to truly useful biomolecular targets. Instead of this conventional physical approach of controlling crystal orientation, here, we propose a chemical approach, i.e., molecular design of polarizing agents; pentacene molecules, the most typical triplet-DNP polarizing agent, are modified so as to make the triplet electron distribution wider and more isotropic without loss of the triplet polarization. In fact, the modification of pentacene with thiophene moieties makes the ESR spectrum sharper and stronger than that of pentacene. To elucidate the effect of the substitutions on spin polarization and zero-field splitting parameters, which determine ESR spectrum, state-of-the-art quantum chemical calculations were performed and revealed that the direction of the spin polarization is altered by the modification with thiophene moieties and the size of D and E parameters are reduced from parent pentacene due to the partial delocalization of spin densities on the thiophene moieties. The triplet-DNP with the new polarizing agent successfully exceeds the previous highest ^1H polarization of glassy materials by a factor of 5. This demonstrates the feasibility of a polarizing agent that can surpass pentacene, the best polarizing agent for more than 30 years since triplet-DNP was first reported, in the unoriented state. This work provides a pathway toward practically useful high nuclear polarization of various biomolecules by triplet-DNP.

INTRODUCTION

Nuclear magnetic resonance (NMR) and magnetic resonance imaging (MRI) are powerful analytical techniques that provide useful information in the chemical and clinical fields, but poor polarization of nuclear spins limits their sensitivity and hinders their versatility. Dynamic nuclear polarization (DNP) is groundbreaking technology that boosts the sensitivity of NMR and MRI by transferring the polarization of electron spins to nuclear spins.¹⁻⁷ However, sufficiently high electron spin polarization in thermal equilibrium requires harsh conditions, such as high-field (~ 7 T) and cryogenic temperatures (~ 1 K), preventing a wide range of practical applications.⁸⁻¹¹

DNP using photo-excited triplet electrons (triplet-DNP) can achieve nuclear hyperpolarization even at room temperature since high electron spin polarization between triplet sub-levels is achieved regardless of temperature by the spin-selective intersystem crossing (ISC) (Figure 1a).¹²⁻¹⁸ The polarization of triplet electron spins can be efficiently transferred to nuclear spins by integrated solid effect (ISE)

sequence with microwave irradiation and magnetic field sweep (Figure 1b). Since the electron spin resonance (ESR) linewidth of triplet is broadened over 100 mT at X-band by hyperfine coupling and dipolar interactions between triplet electrons, only limited part of spin packets simultaneously satisfies the resonance condition for polarization transfer, so-called Hartmann-Hahn condition. By sweeping the magnetic field during microwave irradiation, spin packets in the sweep range can be used for polarization transfer. However, the magnetic field sweep width is practically limited to about 10 mT, since the field sweep must be slow enough to maintain the adiabatic condition and within the lifetime of triplet polarization.

In previous studies of triplet-DNP, the strategy has been to suppress the ESR linewidth broadening due to dipole interactions by strictly controlling the orientation of the single crystal doped with the polarizing agent relative to the magnetic field.^{14,15,19,20} This strategy has enabled most of the electron spin packets to be utilized for polarization transfer, and nuclear spin polarization of several tens of percent has been achieved. However, this strategy is technically

demanding because it requires growing large single crystals on the order of centimeters, and the orientation of the single crystals must be very strictly controlled. Furthermore, the matrix single crystals have been limited to aromatic compounds that can be doped with polarizing agents, which is a serious problem for application to important non-aromatic MRI probes such as pyruvate. Although the transfer of nuclear spin polarization to various molecules by dissolving hyperpolarized naphthalene single crystals at high concentrations has been reported, its medical applications are not easy because it has been limited to DNP in organic solvents.²¹ Considering these difficult challenges faced by the physical strategy of controlling crystal orientation, new breakthroughs are needed to extend the application of triplet-DNP to a variety of biomolecules.




Figure 1. a) Scheme of triplet-DNP. After photoexcitation of polarizing agents, spin-selective ISC produces spin-polarized triplet electrons. Triplet electron spin polarization is transferred to nuclear spin by the integrated solid effect (ISE). b) ISE sequence. The pulsed laser excitation is followed by microwave irradiation and magnetic field sweep. c) ESR simulation with various D and E values while other parameters are fixed ($g = 2$, $B = 8.98$ GHz, $p_x = 0.4$, $p_y = 0.2$, $p_z = 0.2$).

Here, we propose a solution to the essential problem of triplet-DNP by a chemical approach of molecular design of polarizing agents instead of a physical approach of orientation control. In order to achieve triplet-DNP of various biomolecules, it is necessary to achieve high nuclear polarization in randomly oriented samples, free from the constraints of single crystals. The key parameters are the zero-field splitting (ZFS) parameters D and E , which represent the energy gaps between triplet sublevels in the absence of magnetic field and determine the linewidth of the ESR spectrum of the triplet state (Figure 1b).^{22–24} Simulations with different values of D and E while keeping other parameters fixed show that the ESR spectrum becomes sharper as D and E become smaller in the examined range, and that there are more spin packets within 10 mT, the magnetic field sweep width of the ISE sequence (Figure 1c). D and E values are described as follows,

$$D = \frac{3}{4} \frac{\mu_0}{4\pi h} (g_e \mu_B)^2 \left\langle \frac{r^2 - 3z^2}{r^5} \right\rangle$$

$$E = \frac{3}{4} \frac{\mu_0}{4\pi h} (g_e \mu_B)^2 \left\langle \frac{y^2 - x^2}{r^5} \right\rangle$$

where r is the average distance between electron spins, x, y, z are dipole axes, g_e is the g -factor, μ_B is the Bohr magneton, h is plank constant and μ_0 is Magnetic permeability in vacuum. From the above equations, the D and E values are related to the average distance between spins and the rhombicity of the zero-field splitting tensor, respectively. This implies that the molecular design with smaller D and E should make the triplet electron distribution wider and more isotropic.

As a proof of concept, based on pentacene, the most typical polarizing agent of triplet-DNP, we designed novel polarizing agents by modifying pentacene with 5-membered rings, 6,13-di(benzofuran-2-yl)pentacene (DBFP), 6,13-bis(benzo[b]thiophen-2-yl)pentacene (DBTP) and 6,13-di(thiophen-2-yl)pentacene (DTP) (Figure 2a). The relatively less sterically hindered furan and thiophene ring is not orthogonal to pentacene and expected to extend the conjugation, contributing to an extended and more isotropic triplet electron distribution. These new polarizing agents actually exhibit a sharper and stronger ESR signal than pentacene, as well as a higher polarization ratio. The state-of-the-art quantum chemical calculations were performed to predict the ZFS parameters and the ISC rate. The calculations of the D tensor based on molecular structures allows us to predict direction of the principal axis, i.e., the sign of the D and E parameters. It was confirmed that the modifications to parent pentacene enhance broadening the triplet electron distribution, providing theoretical support for the rationality of the present design guidelines, i.e., reducing the size of D and E values. Interestingly, the spin selectivity during the ISC, which is found to be governed by the vibronic effects, was proved to be altered by the modification with thiophene moieties. Triplet-DNP with novel polarizing agents exceeded the ^1H polarization obtained so far in amorphous matrices by a factor of 5, demonstrating that they rationally outperform pentacene, which has been the best performance to date.

RESULTS AND DISCUSSION

Time-resolved ESR measurements. The new polarizing agents DBFP, DBTP and DTP were synthesized following the reported method,²⁵ and their purity was confirmed by NMR and elemental analysis (see Supporting Information for details). For time-resolved ESR measurements, the polarizing agents were dispersed in β -estradiol glass. β -estradiol forms a colorless glass that is stable at room temperature, making it a convenient matrix for ESR measurements.²⁶ Parent pentacene and phenyl-modified pentacene, 6,13-diphenylpentacene (DPP) were used in addition to DBFP, DBTP, and DTP for comparison (Figure 2a).

Each polarizing agent was mixed with β -estradiol at a concentration of 0.01 mol% and dispersed in β -estradiol by heating to 453 K, which is above the melting point of β -estradiol, followed by rapid cooling with liquid nitrogen. Molecularly dispersed state of polarizing agents in β -estradiol was confirmed by absorption and fluorescence spectra. The peak positions and shapes of the absorption spectra of each




Figure 2. a) Chemical structures of polarizing agents, pentacene, 6,13-diphenylpentacene (DPP), 6,13-di(benzofuran-2-yl)pentacene (DBFP), 6,13-bis(benzo[b]thiophen-2-yl)pentacene (DBTP) and 6,13-di(thiophen-2-yl)pentacene (DTP). b) Time-resolve ESR spectra of pentacene, DPP, DBFP, DBTP and DTP doped in β -estradiol at room temperature. Microwave frequencies for each spectral measurement are 8.988, 8.959, 8.983, 8.960 and 8.963 GHz, respectively. Results simulated with EasySpin toolbox in Matlab are shown as red lines. c) Principal axes of zero-field splitting interaction of pentacene triplet electrons.

polarizing agent in the β -estradiol glass were similar to those in the chloroform solution, and were completely different from those of the neat solid, which had a large red shift and broadening (Figure S1). When the absorption spectra were measured by changing the concentration of DBTP in the β -estradiol glass, the spectral shape changed above 0.1 mol%, while it was maintained below 0.01 mol% (Figure S2). Pentacene and its derivatives do not emit

fluorescence due to efficient singlet fission when aggregated,²⁷ but the fluorescence spectra in the β -estradiol glass of each polarizing agent were similar to those in chloroform (Figure S3). From these results, we confirmed that each polarizing agent was dispersed in β -estradiol glass without aggregation at the concentration of 0.01 mol%.

Under pulsed laser irradiation at 527 nm, ESR spectra and signal decays of each polarizing agents in β -estradiol glass were measured at X-band (0.24-0.40 T) and room temperature (Figure 2b, S4). Laser intensity was set to 8.4 A for pentacene, DBFP, DBTP and DTP and 8.8 A for DPP. ESR spectra were simulated by EasySpin toolbox in Matlab²⁸ and obtained parameters were summarized in Table 1. The principal axes of ZFS interactions were same as pentacene and all signs of D and E values were assumed to be $D > 0$ and $E < 0$ (Figure 2c).^{29,30} Compared to pentacene, DBTP and DTP exhibited much stronger and sharper ESR spectra, which is reflected in their smaller values of $|D|$ and $|E|$. On the other hand, DPP showed similar ESR spectral shape and intensity and similar values of $|D|$ and $|E|$ as pentacene. The $|D|$ value was even smaller in DBFP than in DBTP. The reason for these differences in ZFS parameters will be discussed later based on theoretical calculations. The ESR intensity of DBFP was smaller than that of DBTP. This may be due to the lower ISC efficiency of DBFP in addition to the smaller population difference (Table 1, S1). Interestingly, pentacene, DPP, and DBFP were polarized along the x-axis, whereas DBTP and DTP were y-polarized, indicating a clear substituent effect. Decays of the ESR signal showed that both DBTP and DTP exhibited spin-lattice relaxation times T_{1e} comparable to that of pentacene and long enough to apply the ISE sequence (Figure S4, Table S2).

Table 1. $|D|$ and $|E|$ values and relative population ratio of p_x , p_y , p_z of each polarizing agent in the zero-field spin state.

	$ D $ (MHz)*	$ E $ (MHz)*	p_x	p_y	p_z
Pentacene	1360.1	47.2	0.56	0.31	0.13
DPP	1368.3	39.9	0.54	0.34	0.12
DBFP	1190.9	45.1	0.60	0.26	0.14
DBTP	1330.0	26.9	0.20	0.80	0.00
DTP	1311.5	29.2	0.20	0.78	0.02

*Sign of D and E value are assumed to be $D > 0$ and $E < 0$.

Calculations of spin selective ISC rates. To elucidate a detailed mechanism of the spin polarization experimentally observed for the pentacene derivatives, we performed quantum chemical calculations. It was assumed that the spin polarization is caused by a spin selective ISC between the lowest singlet excited state (S_1) and an energetically close triplet state (T_n) of the molecules. To characterize the final state of the ISC, energy levels of the low-lying triplet states of an unmodified pentacene molecule were calculated at the LC-BLYP($\mu=0.15$)/def2-TZVP level of theory. The 0-0 energy differences from S_1 state were predicted as

-1.33, -0.19, 0.77 eV for T_1 (1^3B_{2u}), T_2 (1^3B_{1g}), and T_3 (1^3B_{3u}) states of the pentacene, and therefore the T_2 state that is characterized by the HOMO \rightarrow LUMO+1 and HOMO-1 \rightarrow

LUMO singly excited state is likely to be the final state of the spin selective ISC. The energy differences between the initial (S_1) and final (T_2) at each minimum were predicted to be -193, -85.1, 3.96 meV for pentacene, DBTP, DBFP molecules, respectively. The prediction that the energy level of T_2 state is about the same or higher than that of S_1 state for the DBFP molecule is consistent with the observation in our experiments that the fluorescence quantum yield of DBFP is higher than that of the other molecules, i.e., the ISC transition is not efficient in DBFP.

The ISC transition rate from the S_1 to T_2 spin sublevels T_2^X , T_2^Y , and T_2^Z states are expressed approximately

$$k_{\text{ISC}}^{\mu(=X,Y,Z)} = \frac{2\pi}{\hbar} \sum_{i,f} |\langle \Psi_i | \hat{H}_{\text{SOC}} | \Psi_f^\mu \rangle|^2 \delta(E_i - E_f) \quad (1)$$

applying the Fermi's Golden rule where Ψ_i and Ψ_f^μ indicate vibronic states associated respectively with S_1 and T_2^μ states, E_i and E_f are their energy levels, and \hat{H}_{SOC} is the spin-orbit coupling operator. Since the potential energy surfaces of the spin sublevels are virtually degenerated, i.e., the energy differences are negligible in the prediction of the rates, the difference in the ISC rate for $\mu=X, Y, Z$ should be caused solely by the difference in the magnitude of the SOC matrix elements (SOCMEs) $\langle \Psi_i | \hat{H}_{\text{SOC}} | \Psi_f^\mu \rangle$. Because the SOCMEs of the planar molecules vanish for all the components $\mu = X, Y, Z$ at the equilibrium geometries due to the point group symmetry, and the Heltzberg-Teller like effects, e.g., vibronic spin-orbit coupling, caused by molecular vibrations that break the planar symmetry are dominant,^{31,32} and the SOCMEs should be corrected by adding the first-order vibronic spin-orbit coupling term as

$$\begin{aligned} V^\mu &= \langle \Psi_i | \hat{H}_{\text{SOC}} | \Psi_f^\mu \rangle \\ &\approx \langle \Phi_i | \hat{H}_{\text{SOC}} | \Phi_f^\mu \rangle \Big|_{q=0} \langle v_i | \bar{v}_f \rangle \\ &+ \sum_k \partial_{q_k} \langle \Phi_i | \hat{H}_{\text{SOC}} | \Phi_f^\mu \rangle \Big|_{q=0} \langle v_i | q_k | \bar{v}_f \rangle + O(q^2) \quad (2) \end{aligned}$$

where v_i and \bar{v}_f denotes the vibrational states in the initial (Φ_i) and final (Φ_f^μ) electronic states. Hereafter, we denote the derivative $\partial_{q_k} \langle \Phi_i | \hat{H}_{\text{SOC}} | \Phi_f^\mu \rangle$ at the reference structure ($q=0$) as $\partial_{q_k} V_0^\mu$. There are many ways to evaluate Eq.(1). In this study, the time-dependent formulation³³⁻³⁷ implemented in ORCA^{538,39} was used,^{37,40} where the ISC rates are evaluated by the Fourier transform of the time-correlation function. The approximations adopted in the method, e.g., the potential surfaces are modeled by simple harmonic oscillators, can be insufficient to make quantitative predictions for the ISC rates, but should be sufficient to predict the size relation between the spin sublevels. The calculated ISC rates in the pentacene, DBFP and DBTP molecules are shown in Table 2. The most efficient ISC transitions were predicted for $S_1 \rightarrow T_2^X$ in the pentacene and DBFP molecules and for $S_1 \rightarrow T_2^Y$ in the DBTP molecule. It is consistent with the spin polarization in T_1 state observed in our ESR experiments.

To analyze what vibrational modes contribute to enhancing the vibronic spin-orbit coupling, the first-order derivative of SOCME along each mode was calculated by finite difference techniques. The largest derivative of the SOCME of

Table 2. Predicted ISC rates between S_1 and T_2 sublevels T_2^μ ($\mu = X, Y, Z$) based on the Fermi Golden rule. Relative rate ratios are shown in parentheses.

ns ⁻¹	T_2^X	T_2^Y	T_2^Z
Pentacene	4.4×10^{-2} (0.61)	2.8×10^{-2} (0.38)	1.1×10^{-5} (0.00)
DBFP	1.8×10^{-3} (0.97)	3.1×10^{-5} (0.02)	1.1×10^{-5} (0.01)
DBTP	1.0×10^{-4} (0.00)	5.1×10^{-2} (0.89)	5.9×10^{-3} (0.10)

each sublevel are summarized in Table 3 and the associated normal modes are shown in Figure 3. For the pentacene molecule, it was found that the out-of-plane modes consisting of C-H and C-C bending contribute to increasing the SOCME of the T_2^X and T_2^Y components. Those out-of-plane distortions should mix the carbon π orbitals with σ bonding orbitals, i.e., the transition between carbon $2p_z$ and $2p_y$ or $2p_x$ orbitals are involved in the ISC, and thus the SOCME of the T_2^X and T_2^Y should be enhanced. None of the modes contribute to increasing that of the T_2^Z components. It implies that transitions between carbon $2p_x$ and $2p_y$ orbitals are negligible in the ISC even with the out-of-plane distortions. Similar observation was made for the DBFP molecule.

For the DBTP molecule, the largest derivatives were found along the modes involving displacements in the benzothio- phene substituents, and those modes contribute only to T_2^Y and T_2^Z components. The derivative values along those modes are much larger than that along out-of-plane distortions of the pentacene backbone, which increase the SOCME of the T_2^X and T_2^Y components. For the DBFP molecule, by contrast, the increase in the SOCME caused by the modes involving displacements of the benzofuran substituents is less than those by the out-of-plane modes of the pentacene backbone, and it results in that the transition to the T_2^X component is the most efficient.

The discrepancy between the experimental and theoretical predictions of the order of T_2^X and T_2^Z transition rates for the DBTP molecule, i.e., the transition rate to the T_2^Z was largely overestimated in the FGR calculations, should be attributed to the approximations adopted in the calculations. It is quite possible that the modes that significantly increase the SOCME of the T_2^Z are coupled with the rotation of the benzothio- phene substituents and cannot be described by the harmonic oscillator model nor the first-order vibronic spin-orbit coupling model Eq.(2). In fact, the time-dependent method for the ISC rate could not converged in the case of DTP, which has similar structure to DBTP. In contrast, those approximations should work well for the pentacene molecule because all the modes that contribute to enhancing the ISC are out-of-plane modes and the transition should occur in the vicinity of the minimum along the modes for S_1 and T_2 states. In fact, the ratio is in good agreement with the experiment.

Calculations of ZFS parameters. To investigate the trends of D and E values observed in the experiments, the CASSCF calculations for the T_1 state were performed. Because the D and E values are parameters adopted in the phenomenolog

Table 3. Derivatives of the SOCME along normal modes. Normal modes that give the large derivatives are selected for each sublevel T_2^X , T_2^Y , and T_2^Z .

$\omega_k / \text{cm}^{-1}$	$\frac{\partial_{q_k} V_0^X}{/i \text{ cm}^{-1}}$	$\frac{\partial_{q_k} V_0^Y}{/i \text{ cm}^{-1}}$	$\frac{\partial_{q_k} V_0^Z}{/i \text{ cm}^{-1}}$
Pentacene			
902	0.358	0.002	0.000
866	0.000	0.249	0.000
1508	0.000	0.002	0.007
DBFP			
906	0.249	0.004	0.005
834	0.009	0.127	0.002
1579	0.042	0.003	0.029
DBTP			
901	0.175	0.000	0.000
1364*	0.002	0.281	0.169

*The mode ($\omega = 1364 \text{ cm}^{-1}$) provide the largest derivative for both Y and Z

ical model Hamiltonian, there are many ways to define those values from the *ab initio* quantum chemical calculations. In the case of triplet states, it is straightforward to determine the D and E parameters such that energy gaps of the spin Hamiltonian reproduce the *ab initio* calculations. Because the contributions of the SOC term to the ZFS is negligible as compared with that of the SSC term for aromatic hydrocarbons, only the latter was considered in the calculations. The active space consisting of all the π orbitals of the pentacene backbones, i.e., as much as CAS(22e, 22o), were adopted in the CASSCF calculations. It exceeds the capability of the computer available for the foreseeable future, and the DMRG-CASSCF theory⁴¹⁻⁴³ implemented in the BLOCK2 program⁴⁴ was used with a sufficient bond dimension ($m=1024$) to circumvent the limitation. The D -tensor were calculated as the matrix elements of the SSC between the non-relativistic triplet states with the different spin projections⁴⁵ and the D and E parameters and the orientation of the principal magnetic axis were obtained by the diagonalization. The SSC two-electron integrals were evaluated by using the RI approximation; the two- and three-center integrals were evaluated with Libcint and PySCF⁴⁶⁻⁴⁸ package by using the translational invariance of the two-electron field gradient integrals.

Table 4 shows the calculated D and E parameters for the pentacene derivatives. One of the advantages of predicting the D -tensor by *ab initio* calculations is that the direction of the principal axes in the molecular axis system, i.e., sign of the D and E parameters, can also be directly determined. The signs were predicted to be $D > 0$ and $E < 0$, i.e., the energy ordering is $T_2^X > T_2^Y > T_2^Z$. The signs of the parameters were identical to those reported by the previous experimental study for pentacene molecules in a *p*-terphenyl host crystal,⁴⁹ while those have not been reported for the other




Figure 3. Normalized displacement vector of the modes shown in Table 3. The modes along which the derivatives of the SOCME are less than 0.1, i.e., not significant, are omitted.

molecules as far as we know, the signs predicted are necessary for reproducing the EAEAEA pattern of the experimental ESR spectra for DBTP and DTP in Figure 2b. The DBFP, DBTP, DTP, DPP, and pentacene molecules have larger D parameters in that order, which is in agreement with the experiments except for DPP of which $|D|$ is larger by 8.2 MHz than pentacene in the experiment. It was found that the DBFP, which exhibits the smallest D value, has the smallest dihedral angle between the pentacene backbone and the substituents planes, and the spin density is more delocalized on the substituents than the DPP, DTP, and DBTP (see Figure S14). Although the DPP, DTP, and DBTP have similar size of the dihedral angle, D value of the DTP and DBTP are clearly smaller than that of the DPP, which could reflect the effect of the M-shell electrons of sulfur atoms. Note that the spin-spin interaction is a two-electron interaction and it is not necessary explained solely by the one-electron property, spin density. In fact, it was reported that the mean-field approximation to the spin-spin interaction causes a non-negligible error for ZFS parameters.⁵⁰

As observed in the experiments, the $|E|$ values for the pentacene derivatives were predicted to be very small as less than 50 MHz, though the size relationship between the molecules were not identical between the experiments and *ab*

initio calculation. The energy gap between T_2^X and T_2^Y sublevels is so small that there remains the possibility that the accuracy of the prediction is not fully converged within that range, e.g., the size of the active space and bond dimension etc. Furthermore, the fitting made for experimental spectrum can involve errors when the signal is not enough clear.

Table 4. Calculated D and E parameters obtained by the DMRG-CASSCF theory and dihedral angles between the pentacene backbones and the planar moieties.

	D (MHz)*	E (MHz)*	Dihedral angle
Pentacene	1171.4	-44.5	-
DPP	1143.1	-28.7	69 °
DBFP	1101.3	-17.8	52 °
DBTP	1129.0	-21.1	71 °
DTP	1125.2	-19.9	69 °

*The signs $D > 0$ and $E < 0$ indicate that the energy ordering of the spin sublevels is $T_2^X > T_2^Y > T_2^Z$ where the molecular in-plane long and short axes and the out-of-plane axis correspond to the X, Y, Z principal axes

The theoretical analysis indicates that molecular vibrations that break the planar symmetry and cause a mixing of carbon σ bonding orbitals with π orbitals determine the spin-selectivity in the ISC of the pentacene molecule. The spin-selectivity is largely changed by the substituent modifications; the substituent effect is so large that the direction of the polarization has altered between pentacene (long-axis) and DBTP (short-axis) molecules, though it seems that we should go beyond the harmonic oscillator model or the first-order vibronic spin-orbit coupling model for predicting the rate quantitatively for DBTP molecule. The ZFS parameters D and E are also changed by the modification with substituents. Because the spin-selectivity in the ISC and the ZFS parameters are determined by the SOC and SSC terms, respectively, those properties are expected to be controlled separately by substituent modifications.

Triplet-DNP experiments. To evaluate the potential of the new polarizing agent, the ^1H NMR signal intensity after triplet-DNP was compared between DTP and pentacene. Since β -estradiol has a short ^1H spin-lattice relaxation time T_1 of only a few seconds, *o*-terphenyl (OTP) was employed instead. OTP has been used as a model glass matrix for triplet-DNP because of its long T_1 of about 60 s (Figure S7).⁵¹ The polarizing agents, DTP or pentacene, were doped in OTP using the same procedure except that the heating temperature was changed to 333 K. Triplet-DNP experiments were performed with OTP in the glassy state by ISE sequence under a magnetic field of 0.66 T for pentacene and 0.65 T for DTP at 120 K. After photoexcitation by irradiation of 527 nm pulsed laser, polarization was transferred from electron spins to nuclear spins by microwave irradiation (17.6 GHz) under magnetic field sweep (10 mT). After repeating this

ISE sequence at a frequency of 500 Hz for a certain time, the sample was shuttled into an NMR coil within 1 s and ^1H NMR spectra were measured. (See Supporting Information for details). The ^1H polarization and enhancement factor was calculated by comparing the ^1H NMR signal intensity of ethanol at 296 K and hyperpolarized signal after triplet-DNP at 120 K (Figure S8). Figure 4 shows the buildup curves of ^1H spin polarization by varying the time of triplet-DNP. Experimental parameters such as concentration of the polarizing agent and the magnetic field were optimized based on the intensity of the enhanced NMR signal (Figure S9, S10). ^1H polarization and enhancement factor of OTP after 5 min of triplet-DNP reached 0.85% and 1,540 times, respectively, with DTP, which was about 4 times larger than that with pentacene (0.21% and 370 times). While the enhancement factor obtained by triplet-DNP also depends on other parameters of the polarizing agent, such as electron spin-lattice relaxation time T_{1e} , triplet lifetime τ_T and triplet quantum yield Φ_T but these parameters did not show significant differences for the examined polarizing agents (Figure S4, S5, S6, Table S1, S2). Hence, the novel polarizing agent DTP gave significantly larger ^1H polarization than the best performing pentacene of the past mainly because DTP shows the stronger and sharper ESR than pentacene and can generate more spin packets within the magnetic field sweep range. This was achieved by the rational molecular design of polarizing agents that simultaneously achieves spin-spin interaction control to reduce $|D|$ and $|E|$ values by making the triplet distribution more extended and more isotropic, and spin-orbit coupling control with high sublevel selectivity in ISC.




Figure 4. Build up curve of ^1H signal intensity of OTP doped with pentacene (green line) and DTP (blue line), and partially deuterated OTP (OTP : [D14]OTP = 10 : 90 wt%) doped with DTP (red line).

Final polarization can be improved by reducing the density of ^1H spins to increase the spin-lattice relaxation time T_1 since the accumulation of polarization by triplet-DNP competes with the spin-lattice relaxation. By changing the glass matrix from OTP to a mixture of OTP and fully-deuterated [D14]OTP (OTP : [D14]OTP = 10 : 90 wt%), the ^1H T_1 value was elongated from 1 min to 3 min at 0.65 T and 120 K (Figure S7). Remarkably, this elongated T_1 resulted in the further increase of ^1H polarization and enhancement factor to

7.8% and 14,000 times, respectively (Figure 4), which is 5 times higher than the previous highest polarization of 1.5% in amorphous system by triplet-DNP.⁵¹

CONCLUSION

In conclusion, we have proposed a rational design guideline for polarizing agent as a chemical approach to overcome the intrinsic problem of conventional triplet-DNP, which requires the use of big single crystals with strictly controlled orientation in order to obtain high polarization enhancement. Based on pentacene, the most representative polarizing agent of triplet-DNP, we have succeeded in reducing the ZFS parameters $|D|$ and $|E|$ while maintaining the high polarization of triplet by extending the conjugation through introducing the thiophene ring. This means that it is possible to simultaneously control the spin-spin interaction of the triplet that determines $|D|$ and $|E|$ and the spin-orbit coupling related to spin polarization. Theoretical calculations support for the rationality of the present design guidelines, i.e., by extending the π conjugation to make the distribution of spin density wider and more isotropic, the size of D and E values are reduced and the electron spin polarization is increased at the same time. The power of this chemical approach was demonstrated by exceeding by a factor of 5 the highest ^1H polarization ever achieved by triplet-DNP in amorphous system. This indicates that a polarizing agent that exceeds the performance of pentacene, which has been considered the best over the last 30 years since triplet-DNP was first proposed, is feasible. We believe that further molecular design improvements based on the current discovery will lead to the practical application of triplet-DNP, achieving nuclear polarization of more than 10% for a wide variety of biomolecules under moderate conditions independent of molecular orientation.

ASSOCIATED CONTENT

Supporting Information.

Experimental details including materials, synthesis and measurement setup, UV-vis absorption and fluorescence spectra, time-resolved absorption decays, fluorescence quantum yields, time-resolved ESR and triplet-DNP measurements, calculation details, calculated normal modes and spin density distributions.

AUTHOR INFORMATION

Corresponding Author

*kura@kuchem.kyoto-u.ac.jp

*yanai@mail.cstm.kyushu-u.ac.jp

Author Contributions

§K. S., T. H. and K. M. contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was partly supported by the JST-FOREST Program (JPMJFR201Y), JSPS KAKENHI (JP20H02713, JP22K19051, JP22J21293, and JP19H02675), The Murata Science Foundation, Research Foundation for Opto-Science and Technology, Kyushu University Platform of Inter-/Transdisciplinary Energy Research (Q-PIT) through its "Module-Research Program", the RIKEN-Kyushu University of Science and Technology Hub

Collaborative Research Program, and the RIKEN Cluster for Science, Technology and Innovation Hub (RCSTI).

REFERENCES

- (1) Carver, T. R.; Slichter, C. P. Polarization of Nuclear Spins in Metals. *Phys. Rev.* **1953**, *92* (1), 212–213.
- (2) Overhauser, A. W. Polarization of Nuclei in Metals. *Phys. Rev.* **1953**, *92* (2), 411–415.
- (3) Hall, D. A.; Maus, D. C.; Gerfen, G. J.; Inati, S. J.; Becerra, L. R.; Dahlquist, F. W.; Griffin, R. G. Polarization-Enhanced NMR Spectroscopy of Biomolecules in Frozen Solution. *Science*. **1997**, *276* (5314), 930–932.
- (4) Franck, J. M.; Pavlova, A.; Scott, J. A.; Han, S. Quantitative cw Overhauser effect dynamic nuclear polarization for the analysis of local water dynamics. *Prog. Nucl. Magn. Reson. Spectrosc.* **2013**, *74*, 33–56.
- (5) Ajoy, A.; Liu, K.; Nazaryan, R.; Lv, X.; Zangara, P. R.; Safvati, B.; Wang, G.; Arnold, D.; Li, G.; Lin, A.; Raghavan, P.; Druga, E.; Dhomkar, S.; Pagliero, D.; Reimer, J. A.; Suter, D.; Meriles, C. A.; Pines, A. Orientation-independent room temperature optical ^{13}C hyperpolarization in powdered diamond. *Sci. Adv.* **2018**, *4* (5), eaar5492.
- (6) Dai, D.; Wang, X.; Liu, Y.; Yang, X.-L.; Glaubitz, C.; Denysenkov, V.; He, X.; Prisner, T.; Mao, J. Room-temperature dynamic nuclear polarization enhanced NMR spectroscopy of small biological molecules in water. *Nat. Commun.* **2021**, *12* (1), 6880.
- (7) Armstrong, B. D.; Han, S. Overhauser Dynamic Nuclear Polarization To Study Local Water Dynamics. *J. Am. Chem. Soc.* **2009**, *131* (13), 4641–4647.
- (8) Ardenkjær-Larsen, J. H.; Fridlund, B.; Gram, A.; Hansson, G.; Hansson, L.; Lerche, M. H.; Servin, R.; Thaning, M.; Golman, K. Increase in signal-to-noise ratio of $> 10,000$ times in liquid-state NMR. *Proc. Natl. Acad. Sci.* **2003**, *100* (18), 10158–10163.
- (9) Jannin, S.; Dumez, J.-N.; Giraudeau, P.; Kurzbach, D. Application and methodology of dissolution dynamic nuclear polarization in physical, chemical and biological contexts. *J. Magn. Reson.* **2019**, *305*, 41–50.
- (10) Nelson, S. J.; Kurhanewicz, J.; Vigneron, D. B.; Larson, P. E. Z.; Harzstark, A. L.; Ferrone, M.; van Criekinge, M.; Chang, J. W.; Bok, R.; Park, I.; Reed, G.; Carvajal, L.; Small, E. J.; Munster, P.; Weinberg, V. K.; Ardenkjær-Larsen, J. H.; Chen, A. P.; Hurd, R. E.; Odegardstuen, L.-I.; Robb, F. J.; Tropp, J.; Murray, J. A. Metabolic imaging of patients with prostate cancer using hyperpolarized [^{13}C]pyruvate. *Sci. Transl. Med.* **2013**, *5* (198).
- (11) Golman, K.; in 't Zandt, R.; Thaning, M. Real-time metabolic imaging. *Proc. Natl. Acad. Sci.* **2006**, *103* (30), 11270–11275.
- (12) Henstra, A.; Dirksen, P.; Wenckebach, W. Th. Enhanced dynamic nuclear polarization by the integrated solid effect. *Phys. Lett. A*. **1988**, *134* (2), 134–136.
- (13) Henstra, A.; Lin, T.-S.; Schmidt, J.; Wenckebach, W. Th. High dynamic nuclear polarization at room temperature. *Chem. Phys. Lett.* **1990**, *165* (1), 6–10.
- (14) Iinuma, M.; Takahashi, Y.; Shaké, I.; Oda, M.; Masaike, A.; Yabuzaki, T.; Shimizu, H. M. Proton polarization with *p*-terphenyl crystal by integrated solid effect on photoexcited triplet state. *J. Magn. Reson.* **2005**, *175* (2), 235–241.
- (15) Tateishi, K.; Negoro, M.; Nishida, S.; Kagawa, A.; Morita, Y.; Kitagawa, M. Room temperature hyperpolarization of nuclear spins in bulk. *Proc. Natl. Acad. Sci.* **2014**, *111* (21), 7527–7530.
- (16) Nishimura, K.; Kouno, H.; Kawashima, Y.; Orihashi, K.; Fujiwara, S.; Tateishi, K.; Uesaka, T.; Kimizuka, N.; Yanai, N. Materials chemistry of triplet dynamic nuclear polarization. *Chem. Commun.* **2020**, *56* (53), 7217–7232.
- (17) K. Takeda, Triplet State Dynamic Nuclear Polarization, VDM Verlag, Saarbrücken, **2009**.
- (18) Matsumoto, N.; Nishimura, K.; Kimizuka, N.; Nishiyama, Y.; Tateishi, K.; Uesaka, T.; Yanai, N. Proton Hyperpolarization Relay from Nanocrystals to Liquid Water. *J. Am. Chem. Soc.* **2022**, *144* (39), 18023–18029.

- (19) Takeda, K.; Takegoshi, K.; Terao, T. Dynamic nuclear polarization by electron spins in the photoexcited triplet state: II. High polarization of the residual protons in deuterated naphthalene. *J. Phys. Soc. Jpn.* **2004**, *73* (8), 2319–2322.
- (20) Negoro, M.; Kagawa, A.; Tateishi, K.; Tanaka, Y.; Yuasa, T.; Takahashi, K.; Kitagawa, M. Dissolution Dynamic Nuclear Polarization at Room Temperature Using Photoexcited Triplet Electrons. *J. Phys. Chem. A* **2018**, *122* (17), 4294–4297.
- (21) Eichhorn, T. R.; Parker, A. J.; Josten, F.; Müller, C.; Scheuer, J.; Steiner, J. M.; Gierse, M.; Handwerker, J.; Keim, M.; Lucas, S.; Qureshi, M. U.; Marshall, A.; Salhov, A.; Quan, Y.; Binder, J.; Jahnke, K. D.; Neumann, P.; Knecht, S.; Blanchard, J. W.; Plenio, M. B.; Jelezko, F.; Emsley, L.; Vassiliou, C. C.; Hautle, P.; Schwartz, I. Hyperpolarized Solution-State NMR Spectroscopy with Optically Polarized Crystals. *J. Am. Chem. Soc.* **2022**, *144* (6), 2511–2519.
- (22) Yang, T.-C.; Sloop, D. J.; Weissman, S. I.; Lin, T.-S. Zero-field magnetic resonance of the photo-excited triplet state of pentacene at room temperature. *J. Chem. Phys.* **2000**, *113* (24), 11194–11201.
- (23) Richert, S.; Tait, C. E.; Timmel, C. R. Delocalisation of photoexcited triplet states probed by transient EPR and hyperfine spectroscopy. *J. Magn. Reson.* **2017**, *280*, 103–116.
- (24) Hintze, C.; Steiner, U. E.; Drescher, M. Photoexcited Triplet State Kinetics Studied by Electron Paramagnetic Resonance Spectroscopy. *ChemPhysChem* **2017**, *18* (1), 6–16.
- (25) Vets, N.; Smet, M.; Dehaen, W. Reduction versus Rearrangement of 6,13-Dihydro-6,13-Diarylpentacene-6,13-Diols Affording 6,13- and 13,13'-Substituted Pentacene Derivatives. *Synlett* **2005**, 217–222.
- (26) Yamauchi, A.; Fujiwara, S.; Nishimura, K.; Sasaki, Y.; Tateishi, K.; Uesaka, T.; Kimizuka, N.; Yanai, N. Design Guidelines to Elongate Spin–Lattice Relaxation Times of Porphyrins with Large Triplet Electron Polarization. *J. Phys. Chem. A* **2021**, *125* (20), 4334–4340.
- (27) Yost, S. R.; Lee, J.; Wilson, M. W. B.; Wu, T.; McMahon, D. P.; Parkhurst, R. R.; Thompson, N. J.; Congreve, D. N.; Rao, A.; Johnson, K.; Sfeir, M. Y.; Bawendi, M. G.; Swager, T. M.; Friend, R. H.; Baldo, M. A.; Van Voorhis, T. A transferable model for singlet-fission kinetics. *Nat. Chem.* **2014**, *6* (6), 492–497.
- (28) Stoll, S.; Schweiger, A. EasySpin, a comprehensive software package for spectral simulation and analysis in EPR. *J. Magn. Reson.* **2006**, *178* (1), 42–55.
- (29) Loboda, O.; Minaev, B.; Vahtras, O.; Schimmelpfennig, B.; Ågren, H.; Ruud, K.; Jonsson, D. Ab initio calculations of zero-field splitting parameters in linear polyacenes. *Chem. Phys.* **2003**, *286* (1), 127–137.
- (30) Ganyushin, D.; Gilka, N.; Taylor, P. R.; Marian, C. M.; Neese, F. The resolution of the identity approximation for calculations of spin-spin contribution to zero-field splitting parameters. *J. Chem. Phys.* **2010**, *132* (14), 144111.
- (31) Henry, B. R.; Siebrand, W. Spin–Orbit Coupling in Aromatic Hydrocarbons. Analysis of Nonradiative Transitions between Singlet and Triplet States in Benzene and Naphthalene. *J. Chem. Phys.* **1971**, *54* (3), 1072–1085.
- (32) Tatchen, J.; Gilka, N.; Marian, C. M. Intersystem crossing driven by vibronic spin–orbit coupling: a case study on psoralen. *Phys. Chem. Chem. Phys.* **2007**, *9* (38), 5209.
- (33) Borrelli, R.; Capobianco, A.; Peluso, A. Generating Function Approach to the Calculation of Spectral Band Shapes of Free-Base Chlorin Including Duschinsky and Herzberg–Teller Effects. *J. Phys. Chem. A* **2012**, *116* (40), 9934–9940.
- (34) Peng, Q.; Yi, Y.; Shuai, Z.; Shao, J. Toward Quantitative Prediction of Molecular Fluorescence Quantum Efficiency: Role of Duschinsky Rotation. *J. Am. Chem. Soc.* **2007**, *129* (30), 9333–9339.
- (35) Niu, Y.; Peng, Q.; Deng, C.; Gao, X.; Shuai, Z. Theory of Excited State Decays and Optical Spectra: Application to Polyatomic Molecules. *J. Phys. Chem. A* **2010**, *114* (30), 7817–7831.
- (36) Etinski, M.; Rai-Constapel, V.; Marian, C. M. Time-dependent approach to spin-vibronic coupling: Implementation and assessment. *J. Chem. Phys.* **2014**, *15*.
- (37) de Souza, B.; Neese, F.; Izsák, R. On the theoretical prediction of fluorescence rates from first principles using the path integral approach. *J. Chem. Phys.* **2018**, *148* (3), 034104.
- (38) Neese, F. The ORCA program system. *WIREs Comput. Mol. Sci.* **2012**, *2* (1), 73–78.
- (39) Neese, F.; Wennmohs, F.; Becker, U.; Riplinger, C. The ORCA quantum chemistry program package. *J. Chem. Phys.* **2020**, *152* (22), 224108.
- (40) de Souza, B.; Farias, G.; Neese, F.; Izsák, R. Predicting Phosphorescence Rates of Light Organic Molecules Using Time-Dependent Density Functional Theory and the Path Integral Approach to Dynamics. *J. Chem. Theory Comput.* **2019**, *15* (3), 1896–1904.
- (41) Ghosh, D.; Hachmann, J.; Yanai, T.; Chan, G. K.-L. Orbital optimization in the density matrix renormalization group, with applications to polyenes and β -Carotene. *J. Chem. Phys.* **2008**, *128* (14), 144117.
- (42) Zgid, D.; Nooijen, M. The density matrix renormalization group self-consistent field method: Orbital optimization with the density matrix renormalization group method in the active space. *J. Chem. Phys.* **2008**, *128* (14), 144116.
- (43) Yanai, T.; Kurashige, Y.; Ghosh, D.; Chan, G. K.-L. Accelerating convergence in iterative solution for large-scale complete active space self-consistent-field calculations. *Int. J. Quantum Chem.* **2009**, *109* (10), 2178–2190.
- (44) Zhai, H.; Chan, G. K.-L. Low communication high performance *ab initio* density matrix renormalization group algorithms. *J. Chem. Phys.* **2021**, *154* (22), 224116.
- (45) Ganyushin, D.; Gilka, N.; Taylor, P. R.; Marian, C. M.; Neese, F. The resolution of the identity approximation for calculations of spin-spin contribution to zero-field splitting parameters. *J. Chem. Phys.* **2010**, *132* (14), 144111.
- (46) Sun, Q.; Berkelbach, T. C.; Blunt, N. S.; Booth, G. H.; Guo, S.; Li, Z.; Liu, J.; McClain, J. D.; Sayfutyarova, E. R.; Sharma, S.; Wouters, S.; Chan, G. K. The Python-based Simulations of Chemistry Framework (PYSCF). *WIREs Comput. Mol. Sci.* **2018**, *8* (1).
- (47) Sun, Q.; Zhang, X.; Banerjee, S.; Bao, P.; Barbry, M.; Blunt, N. S.; Bogdanov, N. A.; Booth, G. H.; Chen, J.; Cui, Z.-H.; Eriksen, J. J.; Gao, Y.; Guo, S.; Hermann, J.; Hermes, M. R.; Koh, K.; Koval, P.; Lehtola, S.; Li, Z.; Liu, J.; Mardirossian, N.; McClain, J. D.; Motta, M.; Mussard, B.; Pham, H. Q.; Pulkin, A.; Purwanto, W.; Robinson, P. J.; Ronca, E.; Sayfutyarova, E. R.; Scheurer, M.; Schurkus, H. F.; Smith, J. E. T.; Sun, C.; Sun, S.-N.; Upadhyay, S.; Wagner, L. K.; Wang, X.; White, A.; Whitfield, J. D.; Williamson, M. J.; Wouters, S.; Yang, J.; Yu, J. M.; Zhu, T.; Berkelbach, T. C.; Sharma, S.; Sokolov, A. Yu.; Chan, G. K.-L. Recent developments in the PYSCF program package. *J. Chem. Phys.* **2020**, *153* (2), 024109.
- (48) Sun, Q. Libcint: An efficient general integral library for gaussian basis functions. *J. Comput. Chem.* **2015**, *36* (22), 1664–1671.
- (49) Sloop, D. J.; Yu, H.; Lin, T.; Weissman, S. I. Electron spin echoes of a photoexcited triplet: Pentacene in *p*-terphenyl crystals. *J. Chem. Phys.* **1981**, *75* (8), 3746–3757.
- (50) Sinnecker, S.; Neese, F. Spin-Spin Contributions to the Zero-Field Splitting Tensor in Organic Triplets, Carbenes and Biradicals A Density Functional and Ab Initio Study. *J. Phys. Chem. A* **2006**, *110* (44), 12267–12275.
- (51) Tateishi, K.; Negoro, M.; Kagawa, A.; Kitagawa, M. Dynamic nuclear polarization with photoexcited triplet electrons in a glassy matrix. *Angew. Chem. Int. Ed.* **2013**, *52* (50), 13307–13310.