

# Synthesis of Bicyclo[1.1.0]butanes from Bicyclo[1.1.1]pentanes

Michael D. Mandler\*<sup>1</sup>, James Mignone<sup>1</sup>, Elizabeth A. Jurica<sup>1</sup>, Maximilian D. Palkowitz<sup>2</sup>, Darpandeep Aulakh<sup>3</sup>, Anthony N. Cauley<sup>1</sup>, Christopher A. Farley<sup>1</sup>, Shasha Zhang<sup>3</sup>, Sarah C. Traeger<sup>1</sup>, Amy Sarjeant<sup>3</sup>, Anthony Paiva<sup>1</sup>, Heidi L. Perez<sup>1</sup>, Bruce A. Ellsworth<sup>1</sup>, Alicia Regueiro-Ren<sup>1</sup>

<sup>1</sup>Bristol Myers Squibb Research & Early Development, P.O. Box 4000, Princeton, NJ 08543, United States

<sup>2</sup>Bristol Myers Squibb Research & Early Development, 100 Binney Street, Cambridge, MA 02142, United States

<sup>3</sup>Bristol Myers Squibb Chemical & Synthetic Development, 1 Squibb Drive, New Brunswick, NJ 08901, United States

**KEYWORDS:** Phenyl bioisosteres, Photo-Hunsdiecker, Bicyclo[1.1.1]pentanes, Bicyclo[1.1.0]butanes, Skeletal Editing


**ABSTRACT:** We describe a two step process for the synthesis of substituted bicyclo[1.1.0]butanes. A photo-Hunsdiecker reaction generates iodo-bicyclo[1.1.1]pentanes under metal-free conditions at room temperature. These intermediates react with amine and thiol nucleophiles to afford substituted bicyclo[1.1.0]butane products.

Bicyclo[1.1.1]pentanes (BCPs) are saturated phenyl bioisosteres<sup>1</sup> that have been recently popularized within drug discovery (Scheme 1A).<sup>2</sup> Over the past decade, there have been significant advances in synthetic methods<sup>3</sup> for producing and functionalizing this ring system. Bicyclo[1.1.0]butanes (BCBs), despite being isolated and characterized many decades ago<sup>4</sup>, are far less commonly represented in reported bioactive compounds. Much of the literature describing the synthesis of BCBs concerns intramolecular substitution reactions of cyclobutyl and cyclopropyl carbanions<sup>4B,5</sup> or intramolecular [2+1] reactions with carbenes.<sup>6</sup> Reports of bioactive compounds containing BCBs include one natural product<sup>7</sup>, but primarily include bridgehead-substituted BCBs in which the BCB serves as a pseudo-Michael acceptor for covalent modification by a biological target (Scheme 1B).<sup>8</sup> Curiously, BCPs generally have greater ring strain (67-68 kcal/mol) than BCBs (64-66 kcal/mol).<sup>9</sup> The high *s*-character of the bridgehead carbons allows BCBs to react with electrophiles, nucleophiles, and radicals.<sup>10</sup> As such, they are beginning to be employed more broadly in drug discovery as covalent modulators (Scheme 1B)<sup>11</sup> and as building blocks for the construction of sp<sup>3</sup>-rich molecules.<sup>3A,12</sup>


Adcock found that the reaction of pyridine with 1,3-diiodo-BCP formed a substituted BCP pyridinium salt (Scheme 2A)<sup>13</sup> and Wiberg found that C-I bond heterolysis of 1-iodo-BCPs in the presence of NaN<sub>3</sub> yielded a BCB observed by NMR spectroscopy (Scheme 2B).<sup>14</sup> Given the growing commercial availability of BCPs, we wondered if we could expand upon Wiberg's observation and develop a more general intermolecular reaction between iodo-BCPs and nucleophiles to make substituted BCBs.

First, we searched for the simplest way to make iodo-BCPs. We aimed to directly convert commercially-available BCP-1-carboxylic acids to the corresponding iodides as an alternative to strain-release reactions of [1.1.1]propellane<sup>3B,15</sup> or decarboxylation of redox-active esters.<sup>16</sup> Chemists at Enamine solved the longstanding challenge of preparing BCP-1,3-dicarboxylic acid on scale<sup>17</sup>, leading to increased commercial availability of **1a**, an intermediate that costs \$6.8/mmol.<sup>18</sup>

**Scheme 1.** Examples of bioactive BCPs<sup>2</sup> and BCBs<sup>11A,11B</sup>.




**Scheme 2.** Substituted BCBs can be prepared in two steps from commercially available BCP carboxylic acids.



The iodo-BCP **2a** costs \$383/mmol (Scheme 2C).<sup>19</sup> **2a** has previously been prepared via stepwise Barton ester formation and decarboxylation<sup>20</sup> or via the Kochi reaction<sup>21</sup>, which employs a toxic lead salt. Gandelman reported a mild modification of the photo-Hunsdiecker reaction using diiodohydantoin (DIH) as an iodinating reagent.<sup>22</sup> It was unknown whether these conditions could be extended to BCPs, which were not included in the original substrate scope. When **1a** was treated with 1.0 equiv DIH and light in refluxing 1,2-dichloroethane, evidence of **2a** formation was detected, concomitant with significant side reactions occurring as evidenced by <sup>1</sup>H NMR analysis of the crude reaction mixture. Cooling the reaction to room temperature and switching solvents to dichloromethane suppressed the deleterious formation of side products and smoothly afforded the iodinated product **2a** in 79% yield. We characterized the thermal stability of **2a** by differential scanning calorimetry (DSC) due to a recent report<sup>20</sup> on the instability of related compounds (See Supporting Information).<sup>23</sup>

The diphenylamide analog **1b** was selected as a model substrate with a chromophore to aid reaction monitoring for optimization. When **1b** was treated with 0.5 equiv DIH and irradiated for four hours, **2b** was observed to form with 72% conversion. With additional DIH, the conversion increased to >90% (Table 1, Entries 2, 3).

**Table 1.** Optimization of photo-Hunsdiecker reaction.<sup>a</sup>



entry	DIH (equiv)	solvent	T (°C)	CFL hv	conversion (%)
1	0.5	DCM	23	+	72
2	1.0	DCM	23	+	92
3	1.5	DCM	23	+	94
4	2.0	DCM	23	+	90
5	1.0	DCM	23	-	0
6	1.0	MeCN	23	+	32
7	1.0	acetone	23	+	21
8	1.0	DMF	23	+	5
9	1.0	THF	23	+	0
10	1.0	dioxane	23	+	8
11	1.0	DCE	23	+	55

<sup>a</sup>**1b** and 1,3-diiido-5,5-dimethylhydantoin (DIH) were added to a 4 mL vial equipped with magnetic stirbar and septum. Solvent was added and the reaction was irradiated using a white CFL bulb at 23 °C and monitored by LC/MS. See Supporting Information for experimental details.


With a mild method to access iodo-BCPs **2a-c** in hand, we were intrigued by C-N bond functionalizations based on the results of Adcock<sup>13</sup> and Wiberg (Scheme 2A-B).<sup>24</sup> We hypothesized that we could potentially form amino-substituted BCP products when **2b** is treated with an amine in a polar aprotic solvent. Alternatively, if the rearrangement described by Wiberg<sup>14, 24A</sup> occurred, we could expect a substituted BCB product. The <sup>1</sup>H NMR and <sup>13</sup>C HSQC spectrum of **3** displayed two singlets corresponding to a pair of magnetically inequivalent protons attached to the same carbon, suggesting the BCB was the product. To confirm, an X-ray crystal structure of **3** was acquired that proved a rearrangement had occurred to the BCB.

We thought this substitution reaction could offer a method for the construction of a diverse array of molecules containing the

BCB motif. Optimization of the reaction between morpholine and the iodo-BCP **2b** is described in Table 2. While the reaction tolerated different morpholine loadings (Entries 1-5), we found that the reaction was highly sensitive to solvent and reaction temperature.

Deviating from 80 °C in DMSO led to decreased yields (Entries 6-9). Many solvents were inferior to DMSO (Entries 10-17). Hexafluoroisopropanol (HFIP)<sup>25</sup> and 2,2,2-trifluoroethanol (TFE), which are low-nucleophilic, carbocation-stabilizing solvents, decomposed the starting material and only provided trace yields of the product (Entries 10-11). In contrast, the reaction performed in NMP at 80 °C was sluggish, and the remainder of the mass balance was mostly leftover starting material (Entry 13).


**Table 2.** Optimization of the iodo-BCP substitution reaction.<sup>a</sup>



entry	morpholine (equiv)	solvent	T (°C)	yield (%) <sup>b</sup>
1	1.0	DMSO	80	17
2	1.5	DMSO	80	18
3	2.0	DMSO	80	20 (21)
4	5.0	DMSO	80	19
5	10.0	DMSO	80	17
6	2.0	DMSO	100	18
7	2.0	DMSO	60	5
8	2.0	DMSO	40	4
9	2.0	DMSO	23	2
10	2.0	TFE	80	13
11	2.0	HFIP	80	6
12	2.0	DMF	80	13
13	2.0	NMP	80	4
14	2.0	AcOH	80	0
15	2.0	MeOH	80	13
16	2.0	acetone	80	0
17	2.0	DMPU	80	6
18	2.0	sulfolane	60	3
19	2.0	sulfolane	80	32 (37)
20	2.0	sulfolane	100	24
21	2.0	sulfolane	120	0
22	2.0	sulfolane	150	0
23	2.0	sulfolane	80	5 <sup>c</sup>
24	2.0	sulfolane	80	21 <sup>d</sup>

<sup>a</sup>Stock solutions of **2b** and morpholine were combined in a 4 mL vial equipped with a magnetic stir bar. The vials were sealed and heated for 24 h and then a stock solution of 5,6-dibromobenzo[d][1,3]dioxole (0.1 M, CDCl<sub>3</sub>) was added as an internal standard. An aliquot was removed and then analyzed by <sup>1</sup>H NMR spectroscopy. <sup>b</sup>NMR yields. Isolated yields in parentheses. <sup>c</sup>10 equiv. Ag<sub>2</sub>O added. <sup>d</sup>10 equiv. ZnO added.

Sulfolane<sup>26</sup> was found to be an improvement over DMSO (Entry 19); however, temperatures higher than 80 °C for 24 h led to decomposition (Entries 20-22). Decreasing the temperature to 60 °C led to a ten-fold decrease in NMR yield (Entry 18). Metal additives known to promote carbocation formation from alkyl halides,<sup>27</sup> such as silver oxide and zinc oxide<sup>28</sup> were tested and resulted in lower analytical yields than did additive-free conditions (Entries 23-24).

**Scheme 3.** Substrate scope of BCP to BCB substitution reaction.<sup>a</sup>

<sup>a</sup>All yields reported are isolated yields after chromatography. <sup>b</sup>Microwave reaction. <sup>c</sup>4.0 equiv DIPEA added. <sup>d</sup>Purified by reverse-phase HPLC.


We further examined the substrate scope of the BCP to BCB reaction in sulfolane (Scheme 3). Secondary amine nucleophiles commonly employed in medicinal chemistry were tolerated such as morpholine (**3**), piperidine (**4**), and a ketal-protected piperidone (**5**). Imidazole delivered **6** in 35% isolated yield after 3 days of heating in sulfolane at 80 °C. 1,2,3,4-tetrahydroisoquinoline afforded **7** in 41% yield after a short microwave reaction. We found that microwave heating can quickly probe whether the reaction will be successful, and in many cases isolated yields were comparable to those of conventional heating (See Table S1-S2). Either normal- or reverse-phase chromatography can be used to purify the products. In the former case, sulfolane must first be removed by aqueous workup (e.g. with methyl tert-butyl ether<sup>26</sup>). In the latter case, the sulfolane solution can be directly injected into a reverse-phase HPLC column.

Substitution with pyrrolidine gave **8** in 20% yield. The bulky primary neopentylamine and 1-adamantylamine afforded **9** and **10** respectively, albeit in lower yields. Aliphatic alcohols did not give isolable products, but the more nucleophilic thiol 2-mercaptoethanol afforded **11** in 28% isolated yield. The methyl iodo-BCP carboxylate **1b** was also a competent electrophile (**13-17**).

Limitations included four-membered ring amine nucleophiles that did not lead to product formation (**18-19**); instead, the starting material decomposed. Unfortunately, the tert-butyl ester **2c** did not yield substituted BCB product **20** and instead decomposed upon heating. Aniline did not afford **21** and cyclopentanol gave only trace amounts of **22**.

In all substitution reactions with **2b**, the desired product formed concomitantly with an elimination product that was isolated and characterized as the 3-methylene-1-cyclobutene **23** by X-ray crystallography (Scheme 4). We obtained higher yields of **23** and lower yields of **3** when microwaving at higher temperatures for shorter times. We hypothesize the BCP to BCB substitution reaction occurs via an S<sub>N</sub>1/E1-type mechanism involving the intermediacy of the bicyclo[1.1.0]butyl-1-carbinyl carbocation proposed by Wiberg.<sup>14, 24A</sup> We cannot rule out a concerted mechanism as proposed by Della and coworkers.<sup>29</sup> In our hands, sulfolane was the optimal solvent for this reaction, and while aprotic solvents are not usually associated with promoting S<sub>N</sub>1/E1 reactions, it is reported that sulfolane is capable of strongly solvating cations due to its high dipole moment (4.7 D).<sup>26</sup>

#### Scheme 4. Mechanistic proposal for BCP to BCB conversion.



Since monosubstituted BCBs have been employed as covalent modifiers (Scheme 1B), we subjected compound **3** to a glutathione (GSH) stability assay. We found no GSH conjugation upon incubation at 37 °C with NADPH (see Supporting Information).<sup>30</sup> This aligns with our finding that microwaving two equivalents of 2-mercaptoethanol with **2b** at 130 °C for 40 minutes gave **11** without opening the BCB ring.

We have demonstrated a convenient two-step procedure for the synthesis of substituted BCBs from commercially available starting materials. First, 1-iodo-3-substituted-BCPs are generated by a mild decarboxylative iodination reaction promoted by simple CFL irradiation at ambient temperature. The corresponding iodo-BCPs can then be transformed to substituted BCBs upon reaction with primary and secondary amines or aliphatic thiols in sulfolane. Microwave heating at high temperatures (130–140 °C) for less than 1 hour affords substituted BCB products in similar yields as heat block reactions at 80 °C for several days. We hope that this transformation can be used to expand the collection of BCB derivatives as sp<sup>3</sup>-rich carbocyclic cores.

#### ASSOCIATED CONTENT

##### Supporting Information

General experimental procedures and spectroscopic data for all new compounds. X-ray crystal structures of **3** and **23**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

#### ACCESSION CODES

CCDC 2259736 and 2259735 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

#### AUTHOR INFORMATION

##### Corresponding Author

\*[michael.mandler@bms.com](mailto:michael.mandler@bms.com)

#### ACKNOWLEDGMENTS

The authors would like to thank Ming Gu and Robert Langish (BMS Research and Early Development) for HRMS data. We thank R. Michael Lawrence and Phil S. Baran for helpful discussions.

#### REFERENCES

1. Subbaiah, M. A. M.; Meanwell, N. A., Bioisosteres of the Phenyl Ring: Recent Strategic Applications in Lead Optimization and Drug Design. *J. Med. Chem.* **2021**, *64*, 14046-14128.
2. (A) Stepan, A. F.; Subramanyam, C.; Efremov, I. V.; Dutra, J. K.; O'Sullivan, T. J.; DiRico, K. J.; McDonald, W. S.; Won, A.; Dorff, P. H.; Nolan, C. E.; Becker, S. L.; Pustilnik, L. R.; Riddell,

D. R.; Kauffman, G. W.; Kormos, B. L.; Zhang, L.; Lu, Y.; Capetta, S. H.; Green, M. E.; Karki, K.; Sibley, E.; Atchison, K. P.; Hallgren, A. J.; Oborski, C. E.; Robshaw, A. E.; Sneed, B.; O'Donnell, C. J., Application of the bicyclo[1.1.1]pentane motif as a nonclassical phenyl ring bioisostere in the design of a potent and orally active gamma-secretase inhibitor. *J. Med. Chem.* **2012**, *55*, 3414-3424; (B) Y. Yu, C. Fischer, S. L. Bogen, M. L. Childers, F. X. F. Llinas, J. M. Ellis, S. Esposito, D. M. Hoffman, C. Huang, S. Kattar, A. J. Kim, J. W. Lampe, M. R. Machacek, D. R. McMasters, D. L. Parker, M. Reutershan, N. Sciammetta, P. P. Shao, D. L. Sloman, W. Sun, F. Ujjainwalla, Z. Wu, and C. Gibeau, International Patent, WO 2016089830 A1, 2016.

3. (A) Lopchuk, J. M.; Fjelbye, K.; Kawamata, Y.; Malins, L. R.; Pan, C. M.; Gianatassio, R.; Wang, J.; Prieto, L.; Bradow, J.; Brandt, T. A.; Collins, M. R.; Elleraas, J.; Ewanicki, J.; Farrell, W.; Fadeyi, O. O.; Gallego, G. M.; Mousseau, J. J.; Oliver, R.; Sach, N. W.; Smith, J. K.; Spangler, J. E.; Zhu, H. C.; Zhu, J. J.; Baran, P. S., Strain-Release Heteroatom Functionalization: Development, Scope, and Stereospecificity. *J. Am. Chem. Soc.* **2017**, *139*, 3209-3226; (B) Gianatassio, R.; Lopchuk, J. M.; Wang, J.; Pan, C. M.; Malins, L. R.; Prieto, L.; Brandt, T. A.; Collins, M. R.; Gallego, G. M.; Sach, N. W.; Spangler, J. E.; Zhu, H. C.; Zhu, J. J.; Baran, P. S., Strain-release amination. *Science* **2016**, *351*, 241-246; (C) Alvarez, E. M.; Bai, Z.; Pandit, S.; Frank, N.; Torkowski, L.; Ritter, T., O-, N- and C-bicyclopentylation using thianthrenium reagents. *Nat. Synth.* **2023**; (D) Wright, B. A.; Matviitsuk, A.; Black, M. J.; Garcia-Reynaga, P.; Hanna, L. E.; Herrmann, A. T.; Ameriks, M. K.; Sarpong, R.; Lebold, T. P., Skeletal Editing Approach to Bridge-Functionalized Bicyclo[1.1.1]pentanes from Aza-Bicyclo[2.1.1]hexanes. *ChemRxiv* **2023**.

4. (A) Wiberg, K. B.; Lampman, G. M.; Ciula, R. P.; Connor, D. S.; Schertler, P.; Lavanish, J., Bicyclo[1.1.0]butane. *Tetrahedron* **1965**, *21*, 2749-2769; (B) Wiberg, K. B.; Ciula, R. P., Ethyl Bicyclo[1.1.0]Butane-1-Carboxylate. *J. Am. Chem. Soc.* **1959**, *81*, 5261-5262.

5. (A) Measom, N. D.; Down, K. D.; Hirst, D. J.; Jamieson, C.; Manas, E. S.; Patel, V. K.; Somers, D. O., Investigation of a Bicyclo[1.1.1]pentane as a Phenyl Replacement within an LpPLA(2) Inhibitor. *ACS Med. Chem. Lett.* **2017**, *8*, 43-48; (B) Applequist, D. E.; Renken, T. L.; Wheeler, J. W., Polar Substituent Effects in 1,3-Disubstituted Bicyclo[1.1.1]Pentanes. *J. Org. Chem.* **1982**, *47*, 4985-4995; (C) Hall, H. K., Jr.; Smith, C. D.; Blanchard, E. P., Jr.; Cherkofsky, S. C.; Sieja, J. B., Synthesis and polymerization of bridgehead-substituted bicyclobutanes. *J. Am. Chem. Soc.* **1971**, *93*, 121-130; (D) Hoz, S.; Livneh, M., Cyclobutane-Bicyclobutane System .14. Conversion of Cyclobutane to Bicyclobutane by Base-Catalyzed 1,3-Dehydrohalogenation Reaction - a Mechanistic Study. *J. Am. Chem. Soc.* **1987**, *109*, 7483-7488; (E) McNamee, R. E.; Haugland, M. M.; Nugent, J.; Chan, R.; Christensen, K. E.; Anderson, E. A., Synthesis of 1,3-disubstituted bicyclo[1.1.0]butanes via directed bridgehead functionalization. *Chem. Sci.* **2021**, *12*, 7480-7485.

6. (A) Wipf, P.; Stephenson, C. R. J.; Okumura, K., Transition-metal-mediated cascade reactions: C,C-dicyclopropylmethylamines by way of double C,C-sigma-bond insertion into bicyclobutanes. *J. Am. Chem. Soc.* **2003**, *125*, 14694-14695; (B) Ikota, N.; Takamura, N.; Young, S. D.; Ganem, B., Catalyzed insertion reactions of substituted  $\alpha$ -diazoesters. A new synthesis of cis-enoates. *Tetrahedron Lett.* **1981**, *22*, 4163-4166; (C) Baird, M. S.; Hussain, H. H., The Preparation and Decomposition of Alkyl 2-Diazopent-4-Enoates and 1-Trimethylsilyl-1-Diazobut-3-Enes. *Tetrahedron* **1987**, *43*, 215-224; (D) Qin, C. M.; Davies, H. M. L., Enantioselective Synthesis of 2-Arylbicyclo[1.1.0]butane Carboxylates. *Org. Lett.* **2013**, *15*, 310-313; (E) Panish, R. A.; Chintala, S. R.; Fox, J. M., A Mixed-Ligand Chiral Rhodium(II) Catalyst Enables the Enantioselective Total Synthesis of Piperarborene B. *Angew. Chem. Int. Ed.* **2016**, *55*, 4983-4987; (F) Chen, K.; Huang, X. Y.; Kan, S. B. J.; Zhang, R. K.; Arnold, F. H., Enzymatic construction of highly strained carbocycles. *Science* **2018**, *360*, 71-75.

7. (A) Schneider, C.; Niisuke, K.; Boeglin, W. E.; Voehler, M.; Stec, D. F.; Porter, N. A.; Brash, A. R., Enzymatic synthesis of a



bicyclobutane fatty acid by a hemoprotein-lipoxygenase fusion protein from the cyanobacterium *Anabaena* PCC 7120. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 18941-18945; (B) DeGuire, S. M.; Ma, S. T.; Sulikowski, G. A., Synthesis of a Bicyclobutane Fatty Acid Identified from the Cyanobacterium *Anabaena* PCC 7120. *Angew. Chem. Int. Ed.* **2011**, *50*, 9940-9942.

8. Kelly, C. B.; Milligan, J. A.; Tilley, L. J.; Sodano, T. M., Bicyclobutanes: from curiosities to versatile reagents and covalent warheads. *Chem. Sci.* **2022**, *13*, 11721-11737.

9. (A) Khoury, P. R.; Goddard, J. D.; Tam, W., Ring strain energies: substituted rings, norbornanes, norbornenes and norbornadienes. *Tetrahedron* **2004**, *60*, 8103-8112; (B) Wiberg, K. B., The Concept of Strain in Organic-Chemistry. *Angew. Chem. Int. Ed.* **1986**, *25*, 312-322.

10. (A) Fawcett, A., Recent advances in the chemistry of bicyclo- and 1-azabicyclo[1.1.0]butanes. *Pure Appl. Chem.* **2020**, *92*, 751-765; (B) Walczak, M. A. A.; Krainz, T.; Wipf, P., Ring-Strain-Enabled Reaction Discovery: New Heterocycles from Bicyclo[1.1.0]butanes. *Acc. Chem. Res.* **2015**, *48*, 1149-1158.

11. (A) T. Mukaiyama, P. C. Chua and J.-M. Vernier, International Patent, WO 2021173923 A1, 2021; (B) B. Lefker, K. Gibson, M. Spendiff, P. Humphries, S. Bucknell, W. Zawodny and R. Porter, International Patent, WO 2022051596 A1, 2022; (C) Tokunaga, K.; Sato, M.; Kuwata, K.; Miura, C.; Fuchida, H.; Matsunaga, N.; Koyanagi, S.; Ohdo, S.; Shindo, N.; Ojida, A., Bicyclobutane Carboxylic Amide as a Cysteine-Directed Strained Electrophile for Selective Targeting of Proteins. *J. Am. Chem. Soc.* **2020**, *142*, 18522-18531.

12. (A) Golfmann, M.; Walker, J. C. L., Bicyclobutanes as unusual building blocks for complexity generation in organic synthesis. *Commun Chem* **2023**, *6*; (B) Jung, M.; Lindsay, V. N. G., One-Pot Synthesis of Strain-Release Reagents from Methyl Sulfoxes. *J. Am. Chem. Soc.* **2022**, *144*, 4764-4769.

13. Adcock, J. L.; Gakh, A. A., Synthesis of Novel Cage Quaternary-Salts Via Nucleophilic-Substitution of 1,3-Diiodobicyclo[1.1.1]Pentane - Further Evidence for a Stable 3-Iodo-1-Bicyclo[1.1.1]Pentyl Cation. *Tetrahedron Lett.* **1992**, *33*, 4875-4878.

14. Wiberg, K. B.; McMurdie, N., Mechanism of the Solvolysis of Bicyclo[1.1.1]Pentyl-1 Derivatives. *J. Org. Chem.* **1993**, *58*, 5603-5604.

15. (A) Wiberg, K. B.; Waddell, S. T.; Laidig, K., [1.1.1]Propellane - Reaction with Free-Radicals. *Tetrahedron Lett.* **1986**, *27*, 1553-1556; (B) Wiberg, K. B.; Walker, F. H., [1.1.1]Propellane. *J. Am. Chem. Soc.* **1982**, *104*, 5239-5240; (C) Nugent, J.; Arroniz, C.; Shire, B. R.; Sterling, A. J.; Pickford, H. D.; Wong, M. L. J.; Mansfield, S. J.; Caputo, D. F. J.; Owen, B.; Mousseau, J. J.; Duart, F.; Anderson, E. A., A General Route to Bicyclo[1.1.1]pentanes through Photoredox Catalysis. *ACS Catal.* **2019**, *9*, 9568-9574; (D) Wong, M. L. J.; Mousseau, J. J.; Mansfield, S. J.; Anderson, E. A., Synthesis of Enantioenriched alpha-Chiral Bicyclo[1.1.1]pentanes. *Org. Lett.* **2019**, *21*, 2408-2411; (E) Dilmac, A. M.; Spuling, E.; de Meijere, A.; Brase, S., Propellanes-From a Chemical Curiosity to "Explosive" Materials and Natural Products. *Angew. Chem. Int. Ed.* **2017**, *56*, 5684-5718; (F) Makarov, I. S.; Brocklehurst, C. E.; Karaghiosoff, K.; Koch, G.; Knochel, P., Synthesis of Bicyclo[1.1.1]pentane Bioisosteres of Internal Alkynes and para-Disubstituted Benzenes from [1.1.1]Propellane. *Angew.*

*Chem. Int. Ed.* **2017**, *56*, 12774-12777; (G) Della, E. W.; Taylor, D. K., Synthesis of Some Bridgehead Bridgehead-Disubstituted Bicyclo[1.1.1]Pentanes. *J. Org. Chem.* **1994**, *59*, 2986-2996; (H) Kanazawa, J.; Maeda, K.; Uchiyama, M., Radical Multicomponent Carboamination of [1.1.1]Propellane. *J. Am. Chem. Soc.* **2017**, *139*, 17791-17794.

16. Fu, M. C.; Wang, J. X.; Shang, R., Triphenylphosphine-Catalyzed Alkylative Iododecarboxylation with Lithium Iodide under Visible Light. *Org. Lett.* **2020**, *22*, 8572-8577.

17. Ripenko, V.; Vysochyn, D.; Klymov, I.; Zherish, S.; Mykhailiuk, P. K., Large-Scale Synthesis and Modifications of Bicyclo[1.1.1]pentane-1,3-dicarboxylic Acid (BCP). *J. Org. Chem.* **2021**, *86*, 14061-14068.

18. <https://www.combi-blocks.com/cgi-bin/find.cgi?QJ-0241>. Accessed 2/22/2023.

19. <https://www.combi-blocks.com/cgi-bin/find.cgi?JP-2803-1g>. Accessed 2/23/2023.

20. Dallaston, M. A.; Houston, S. D.; Williams, C. M., Cubane, Bicyclo[1.1.1]pentane and Bicyclo[2.2.2]octane: Impact and Thermal Sensitiveness of Carboxyl-, Hydroxymethyl- and Iodo-substituents. *Chem. Eur. J.* **2020**, *26*, 11966-11970.

21. Kuduva, S. S.; Craig, D. C.; Nangia, A.; Desiraju, G. R., Cubanecarboxylic Acids. Crystal Engineering Considerations and the Role of C-H...O Hydrogen Bonds in Determining O-H...O Networks. *J. Am. Chem. Soc.* **1999**, *121*, 1936-1944.

22. Kulbitski, K.; Nisnevich, G.; Gandelman, M., Metal-Free Efficient, General and Facile Iododecarboxylation Method with Biodegradable Co-Products. *Adv. Synth. Catal.* **2011**, *353*, 1438-1442.

23. 2a decomposed with an onset temperature of 102 °C and decomposition energy of -587 J/g followed by a second exotherm of -246 J/g starting at 221 °C (Table S3).

24. (A) Wiberg, K. B.; McMurdie, N., Formation and reactions of bicyclo[1.1.1]pentyl-1 cations. *J. Am. Chem. Soc.* **1994**, *116*, 11990-11998; (B) Wiberg, K. B.; Williams, V. Z., Solvolytic reactivity of 1-chlorobicyclo[1.1.1]pentane. *J. Am. Chem. Soc.* **1967**, *89*, 3373-3374.

25. Ramos-Villasenor, J. M.; Rodriguez-Cardenas, E.; Diaz, C. E. B.; Frontana-Urbe, B. A., Review-Use of 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP) Co-Solvent Mixtures in Organic Electrosynthesis. *J. Electrochem. Soc.* **2020**, *167*.

26. Tilstam, U., Sulfolane: A Versatile Dipolar Aprotic Solvent. *Org. Process Res. Dev.* **2012**, *16*, 1273-1278.

27. Kraus, G. A.; Hon, Y. S.; Thomas, P. J.; Laramay, S.; Liras, S.; Hanson, J., Organic-Synthesis Using Bridgehead Carbocations and Bridgehead Enones. *Chem. Rev.* **1989**, *89*, 1591-1598.

28. Gurudutt, K. N.; Ravindranath, B.; Srinivas, P., Solvolytic Displacement of Alkyl-Halides by Metal-Salts - Preparative Procedures for Allyl-Oxy, Benzyl-Oxy and Tertiary Alkyl-Oxy Derivatives Using the Zinc Salts. *Tetrahedron* **1982**, *38*, 1843-1846.

29. Della, E. W.; Grob, C. A.; Taylor, D. K., Bridgehead carbocations: A solvolytic study of 1-bromobicyclo[1.1.1]pentane and its bridgehead-substituted derivatives. *J. Am. Chem. Soc.* **1994**, *116*, 6159-6166.

30. Paiva, A. A.; Klakouski, C.; Li, S.; Johnson, B. M.; Shu, Y.-Z.; Josephs, J.; Zvyaga, T.; Zamora, I.; Shou, W. Z., Development, optimization and implementation of a centralized metabolic soft spot assay. *Bioanalysis* **2017**, *9*, 541-552.

