

# PyConSolv:

## A Python Package for Conformer Generation of (Metal-Containing) Systems in Explicit Solvent

*R. A. Talmazan<sup>[1]</sup> and M. Podewitz\*<sup>[1]</sup>*

[1] Institute of Materials Chemistry, TU Wien, Getreidemarkt 9, A-1060 Wien (Austria),  
[maren.podewitz@tuwien.ac.at](mailto:maren.podewitz@tuwien.ac.at)

Keywords: Conformer, Conformer generation, Python, Automation, MCPB, Transition-Metal  
Complex

### **Abstract**

We introduce PyConSolv, a freely available python package that automates the generation of conformers of metal and non-metal containing complexes in explicit solvent, through classical molecular dynamics simulations. Using a streamlined workflow and interfacing with widely used computational chemistry software, PyConSolv is an all-in-one tool for the generation of conformers in any solvent. Input requirements are minimal, only the geometry of the structure and the desired solvent in xyz (XMOL) format are needed. The package can also account for charged systems, by including arbitrary counterions in the simulation. A bonded model parametrization is

performed automatically, utilizing the AmberTools, ORCA, and Multiwfn software packages. PyConSolv provides a selection of pre-parametrized solvents and counterions for use in classical molecular dynamics simulations. We show the applicability of our package on a number of (transition-metal-containing) systems. The software is provided open-source and free of charge.

## **Introduction**

As quantum mechanical (QM) methods and computer hardware evolve and become more efficient, bigger systems,<sup>1-3</sup> that were previously far out of reach, can be tackled using quantum chemical wave function methods.<sup>4,5</sup> As the system size increases, it becomes necessary to consider greater flexibility, which in turn presents new challenges that need to be properly addressed.<sup>6,7</sup> One common approach is to generate conformational ensembles, which can be created in various ways.<sup>8</sup>

While the importance of conformers is well known and a key feature in drug design pipelines,<sup>9,10</sup> it has only recently begun gaining traction in the field of homogenous catalysis.<sup>11-16</sup> A systematic exploration of the conformational space, by generating rotamers, can be employed for simple systems, with few degrees of freedom, but quickly becomes unfeasible for larger molecules. While several more advanced methods for the generation of conformers exist, they are generally geared towards biological compounds<sup>17-20</sup>. Alternatively, molecular dynamics (MD) methods can be employed for the generation of conformers, by running simulations of the system (in explicit solvent) and clustering the resulting trajectory.<sup>21</sup> For this purpose, ab initio MD (AIMD)<sup>22</sup> or classical MD (cMD), in various flavors, can be used, each with its own advantages and disadvantages, as outlined below.

For AIMD, calculating the electronic structure is generally considered to be the limiting factor for the overall sampling. To overcome this issue, the use of very efficient semi-empirical methods,

such as GFN2-xTB,<sup>23</sup> provides a good compromise. One tailor-made approach for conformer generation is realized in the Conformer Rotamer Ensemble Tool (CREST),<sup>24</sup> which is shown to perform very well for a large a number of systems, yet it is not without its downsides. Due to the inherent limitations of semi-empirical quantum chemical methods, GFN2-xTB and CREST do not perform well for typical transition-metal systems and neither structures nor energies are reliable without any further processing.<sup>25</sup> Another caveat for CREST is the lack of explicit solvation. While implicit solvation models perform admirably for many complexes,<sup>26,27</sup> they fail to provide reasonable structures when looking at systems containing a cavity,<sup>15</sup> as intramolecular bonds are heavily favored. This can be counteracted by using explicit solvation, either in microsolvation approaches,<sup>28-30</sup> that use a few solvent molecules, or in condensed phase calculations, but in any case, at the cost of massively increasing the computational requirements.

On the other hand, cMD simulations rely on a force-field approach, which leads to a great speedup in calculations. Consequently, it allows for explicit treatment of solvent molecules, of course, at some cost in accuracy. While many force-fields exist, developed with either specific<sup>31-35</sup> or general use,<sup>36-40</sup> running a cMD simulation for metal containing systems requires the generation of custom parameters of the system at hand, to obtain reasonable results. While there are tools created for the parametrization of organic molecules,<sup>41</sup> the parametrization of transition metal containing complexes is a more involved process, requiring a special approach.<sup>42-46</sup> As metals can form very complex structures, with widely varying coordination numbers,<sup>47</sup> there are no predefined atom parameters available in the widely used general force fields. A full parametrization of the metal center is therefore required. One tool available in the AmberTools<sup>48</sup> suite, developed for parametrization of metal-containing biological systems, is the Metal Center Parameter Builder (MCPB).<sup>49</sup> MCPB utilized a bonded model<sup>42</sup> to describe the metal ion and its

surrounding environment. This approach requires defining the bonded and nonbonded parameters of the force field, for the metal atom and its ligands. These parameters entail structural information, force constants and potentials for bonds, angles and dihedrals as well as partial charges and van der Waals parameters. They are generated based on either experimental or QM optimized structures, by deriving their values from force constants and partial charges.<sup>49</sup> We would like to stress here that the force field parameters are generated in such way that the QM optimized structure is retained. While using a bonded model as implemented in MCPB provides accurate results for many different (transition-)metal containing complexes,<sup>14,15,42,49-51</sup> the generation of parameters is an intricate process and error prone, due to a lot of user intervention required.


In this work, we present a user-friendly python package, which builds upon AmberTools,<sup>48</sup> to provide an automated process for generating conformers of arbitrary, metal-containing or metal-free complexes, in explicit solvent. The user only needs to provide an input structure, the desired QM method to be used to the geometry optimization and force constant calculations, total system charge and multiplicity, as well as the solvent to be used to the simulation (if applicable). We provide full support for a large number of pre-parametrized solvents, as well as the ability to parametrize any solvent of choice. Additionally, as many metal-containing complexes are associated with a counterion, we provide seven pre-parametrized ions, as well as the ability to parametrize an ion of choice. By interfacing to ORCA 5,<sup>52</sup> the user has access to state-of-the-art QM methods for structure calculations. For system charge assignment, an interface to Multiwfn 3.8,<sup>53</sup> offers the user a plethora of charge calculations schemes, in addition to the Merz-Kollman RESP scheme<sup>54,55</sup> recommended by MCPB. After the parametrization is complete, the system is solvated and a suggested equilibration procedure is offered to the user, using the Amber MD package.<sup>56,57</sup> Once a simulation is performed, the resulting trajectory file can be analyzed via a

script provided by PyConSolv. The clustering itself is based on the popular cpptraj package<sup>58</sup> and returns a number of conformers, ranked by their QM energy, based on single point calculations.

## **Methodology**

In this work, conformational sampling is performed for an array of structures in explicit solvent using PyConSolv to show the applicability as a proof-of-concept. The systems of choice are a copper containing calix[8]arene catalyst for C-N coupling, a molybdenum based catalyst for olefin metathesis, the metal-free hydrogenobyrinic acid, as well as the vitamin B<sub>12</sub> metabolite methylcobalamin.

**Parametrization.** The protocol implemented in the PyConSolv python package is shown in Figure 1. The input required is a simple XMOL xyz formatted molecular structure. Input files for ORCA 5 are generated automatically, and geometry optimization and subsequent frequency calculations are performed at the electronic structure theory level chosen by the user. The optimized geometry is taken as input for subsequent parametrization steps. As the metal center parametrization builds upon the MCPB.py package provided in AmberTools, the steps follow those recommended in the



**Figure 1.** PyConSolv workflow steps.

aforementioned package. The structure must be split into fragments which are to be parametrized separately, with each atom requiring a unique identifier in the pdb files. This step is automated by building a connectivity matrix based on atom radii and pair distances of the atoms. If two atoms are closer together than 60% of the sum of their atomic radii, they are considered bound. If one of the atoms is a metal, the connectivity is checked, but it is not added to the matrix, as the metal needs to be parametrized separately from the rest of the fragments. Using a depth first search algorithm<sup>59</sup> to traverse the connectivity matrix, we are able to identify each individual fragment. The user is then presented with an interactive window where a Lewis structure of each fragment is displayed. Here, we require the user to provide total fragment charges to perform the parametrization of each individual fragment using antechamber.<sup>41</sup> The ORCA 5 output files are analyzed and converted into inputs for Multiwfn 3.8 and MCPB, also accounting for the usage of effective core potentials<sup>60</sup> for heavy

atoms. The RESP charges for the system are calculated using Multiwfn, with the recommended settings, as in the Multiwfn user manual (see SI). The MCPB input file is created, taking into account the carbon-metal bonds present in the system, which are not natively recognized. Using the parameters generated by MCPB, a simulation box is then set up, with either one of the 18 preconfigured solvents or any other user-defined solvent. Subsequently, an equilibration script, which follows an extensive heating and cooling procedure as outlined by Wallnöfer et al.<sup>61</sup>, is provided, as well as an input file for a simulation of 100 ns in the NVT ensemble, with T=300K.

**Supported Solvents and Ions.** As part of the PyConSolv package, the solvents shown in Table S1 (SI) have been pre-parametrized and verified that the equilibrated solvent box density at 300K is close to experimentally determined densities. All parametrizations were performed on structures optimized with BP86/def2-SVP/D4<sup>62-65</sup>, with the exception of acetonitrile, for which no D4 corrections were used, as the resulting structure was incorrect. For parametrization, antechamber was used, with RESP charges.

Alongside solvents, the following popular counterions have been parametrized:  $\text{ScF}_6^{3-}$ ,  $\text{BF}_4^-$ ,  $\text{B}[\text{Ar}^{\text{F}}]^-$ ,  $\text{B}(\text{Ph})^-$ ,  $\text{PF}_6^-$ ,  $\text{OTf}^-$  and  $\text{ClO}_4^-$ . The structures have been optimized with BP86/def2-SVP/D4<sup>62-65</sup> and all charges computed with RESP. During the parametrization, for  $\text{BF}_4^-$  and  $\text{ClO}_4^-$ , the angle parameters were unable to be determined automatically with antechamber/MCPB.py. In these cases, a scan of the angle was performed with ORCA and the parameter assigned manually.

## Analysis

After a cMD production run has completed, the resulting trajectory can be analyzed using PyConSolv. This can be done in two ways, either using one of the provided shell scripts or using

the python implementation present within PyConSolv. To obtain relevant conformers, the trajectory must be clustered. As we use a root mean square deviation (RMSD) of the distance of all heavy atoms as the clustering criteria, it is vital that the trajectory is properly aligned. This is resolved by aligning the solute based on a list of atom indices, provided by the user. The clustering can then be performed using one of the four methods provided by cptraj: dbscan, hierarchical, k-means or dpeaks<sup>58</sup>. On the resulting clusters, single point calculations are performed, using the same electronic structure theory level that was used during the parametrization. Finally, a list of clusters, ranked by the single point energy, is presented to the user.


## Results

The PyConSolv workflow, as explained above, was applied to the following systems: Cu(I)-calix[8]arene<sup>15</sup>, a molybdenum based catalyst for olefin metathesis<sup>66,67</sup>, hydrogenobyric acid<sup>68-70</sup> and methylcobalamin<sup>71</sup>. A suitable functional was chosen for each system, along with the appropriate solvent, as to mimic the experimental conditions. All charges were calculated using RESP. After parametrization, the system was equilibrated and a 100 ns cMD production run was performed, using the simulation input file provided by PyConSolv. The clusters were generated using k-means, with the script provided by PyConSolv, using the RMSD of all non-hydrogen atoms as the distance metric. The simulations were aligned on the “rigid” parts of the structure, as described in each case below. A total of 10 clusters was chosen for each system and evaluated.

As a note about runtime, when comparing PyConSolv with CREST, it can be seen that with increase in system size, the PyConSolv workflow becomes noticeably faster, as the time-consuming step becomes the frequency calculation, rather than the cMD simulation (see SI Table S2 for timings).



## Case 1: Cu(I)-Calix[8]arene



**Figure 2.** Cu(I)-calix[8]arene clusters. Upper Panel: Top three clusters with lowest QM energy. Lower Panel: all 10 clusters superimposed (left). The bottom right image depicts a collapsed cage structure in implicit solvation.


Cu(I)-calix[8]arene is a catalyst that has proven to be very complex to model, with the macrocyclic cage being particularly mobile. This inherent flexibility needs to be taken into account, in order to obtain an accurate energy profile for the reaction<sup>15</sup> or to explain the difference in activity between various regioisomers.<sup>14</sup>

Using PyConSolv, with PBE0/def2-SVP/D3 + CPCM(Chloroform)<sup>26,65,72,73</sup> the system shown in Figure 2 was parametrized and solvated in chloroform. The trajectory was aligned on the copper and phenanthroline moieties. The clusters shown in Figure 2

represent those with the lowest energy, as computed via DFT single points, with the above specified density functional, basis set and dispersion correction. For comparison, structures generated with CREST using GFN2-xTB and implicit solvation (chloroform) have been generated as well. From the representative cluster depicted in Figure 2, it can be seen that the calix[8]arene cage is collapsed around the copper center. when compared to the structures generated with PyConSolv. This is an artefact caused by the lack of explicit solvation in the CREST conformers and the tendency of dispersion corrections to favor compact structures with many intramolecular

interactions. The collapsed structure is in stark contrast to those generated in explicit solvent via PyConSolv (see Figure 2, bottom left).

### Case 2: Molybdenum N-heterocyclic carbene (NHC) catalyst



**Figure 3.** Mo imido alkylidene NHC catalyst clusters. Upper panel: Top three ranked clusters according to QM energy. Lower panel: All 10 clusters superimposed (left). The bottom right highlights the high flexibility of the triflate groups.


Molybdenum imido alkylidene N-heterocyclic carbene (NHC) catalysts represent one avenue of performing olefin metathesis.<sup>66,67</sup> It was shown during the investigation of the reaction mechanism, that conformer generation is crucial for explaining the reactivity.<sup>74</sup>

In the case of the Mo-based catalyst, the structures generated with PyConSolv, using BP86/def2-SVP/D3BJ+CPCM(CH<sub>2</sub>Cl<sub>2</sub>)<sup>26,62,63,65,75</sup>, for force field parametrization. The trajectory was aligned on the molybdenum, the carbon of the aryl group and the NHC carbon bound to the molybdenum. It can be seen that the

simulation successfully captured the movement of the ligands, as shown in Figure 3. The triflate ligands show the most prominent flexibility. For comparison, structures were generated with CREST (see Figure S1, SI). Here, the solvent does not seem to have a great influence on the overall structure of the catalyst. The movement of the triflate groups is more pronounced in the CREST

conformers, likely due to the enhanced sampling method implemented within its code. While similar enhanced sampling methods can easily be utilized in conjunction with PyConSolv, this is beyond the scope of this work. The NHC moiety, along with the aryl ligand appear to be considerably more rigid. While the CREST conformers display a rotation in the NHC group, this is not present in the cMD simulation set up by PyConSolv. The underlying electronic structure method of CREST, GFN2-xTB, has shown to be less appropriate for this particular catalyst and as such, the resulting structures require further refinement using DFT. As the coordination geometry of the Mo center is restrained in the force field to the QM optimized one, an adequate structure is conserved in the simulations at the cost that changes in the Mo coordination geometry cannot be captured.

### Case 3: Hydrogenobyric Acid




**Figure 4.** Hydrogenobyric acid clusters. Upper panel: Top three clusters with lowest QM energy. Lower panel: All 10 clusters superimposed (left) and a view of the corrin ring of the clusters (right).

Hydrogenobyric acid represents the metal-free precursor of vitamin B<sub>12</sub> from which numerous native or artificial metal-cobalamins can be synthesized.<sup>68-70</sup> Its structural variability can prove to be interesting to drug design. The electronic structure calculation for parametrization was performed with BP86/def2-SVP/D4+CPCM(Water).<sup>26,62-65</sup> and the trajectory aligned on the nitrogen atoms of the corrin ring.

It can be seen from Figure 4, that the flexibility of the side chains is quite well captured, alongside the flexibility of the corrin ring. The testcase also illustrates that PyConSolv works equally well for metal-free systems.

#### Case 4: Methylcobalamin



**Figure 5.** Methylcobalamin clusters. Upper panel: Top three clusters according to QM energy. Lower panel: All 10 clusters superimposed (left) and an example of a structure optimized in implicit solvent, dominated by intramolecular hydrogen bonding (right).

Methylcobalamin is a vitamin B<sub>12</sub> metabolite.<sup>71</sup> For the PyConSolv procedure, as in case 3, for parametrization, BP86/def2-SVP/ D4+CPCM(Water)<sup>26,62–65</sup> was the method of choice. The alignment was performed, as before, on the nitrogen atoms of the corrin ring.

It can be seen from the clusters in Figure 5, that the methylcobalamin structure is fairly rigid. The stabilization of the corrin ring is apparent when compared to that of the hydrogenobyric acid. It can be seen that all amide groups show some flexibility, while always pointing towards the solvent.

For sake of comparison, investigation of the complex in implicit solvent only, yielded a structure with many intramolecular hydrogen bonds of the amides and a loop conformation that is very different to the ones in explicit solvent (see Figure 5 bottom right, Figure S2 in SI). These intramolecular hydrogen bonds may affect the reactivity of the complex and it is unlikely that they

are formed under experimental conditions. Consequently, PyConSolv provides more realistic structures here.

## **Conclusion**

In this work, we present a python package to automate conformer generation of (metal containing) complexes in explicit solvent. Our tool performs parametrization of metal-containing structures and sets up a workflow for simulation and analysis. The automatization makes conformer generation accessible to the non-expert user. It allows for a straightforward implementation of conformational sampling (in explicit solvent) in standard computational chemistry workflows, e.g., for the determination of reaction mechanisms. With increasing system size, our python package outperforms semi-empirical methods, such as CREST. A subsequent reoptimization of the obtained clusters with DFT ensures that true energy minima are found.

The case studies highlight that conformer generation in explicit solvent yields more realistic structures. Our tool is a stepping stone towards more realistic modelling of reaction mechanisms and therefore a requisite for calculations becoming predictive.

## **Availability**

The PyConSolv package is available open-source, free of charge, on GitHub: <https://github.com/PodewitzLab/PyConSolv>

Likewise, it can be installed from the popular PyPi software repository using:

```
pip install PyConSolv
```

## **Acknowledgement**

The authors would like to thank the FWF for financial support (P33528), project granted to M.P.

# TOC



## References

- (1) Bowler, D. R.; Miyazaki, T. Calculations for Millions of Atoms with Density Functional Theory: Linear Scaling Shows Its Potential. *J. Phys.: Condens. Matter* **2010**, *22* (7), 074207. <https://doi.org/10.1088/0953-8984/22/7/074207>.
- (2) Grimme, S.; Schreiner, P. R. Computational Chemistry: The Fate of Current Methods and Future Challenges. *Angew. Chem. Int. Ed.* **2018**, *57* (16), 4170–4176. <https://doi.org/10.1002/anie.201709943>.
- (3) Houk, K. N.; Liu, F. Holy Grails for Computational Organic Chemistry and Biochemistry. *Acc. Chem. Res.* **2017**, *50* (3), 539–543. <https://doi.org/10.1021/acs.accounts.6b00532>.
- (4) Guo, Y.; Riplinger, C.; Becker, U.; Liakos, D. G.; Minenkov, Y.; Cavallo, L.; Neese, F. Communication: An Improved Linear Scaling Perturbative Triples Correction for the Domain Based Local Pair-Natural Orbital Based Singles and Doubles Coupled Cluster Method [DLPNO-CCSD(T)]. *J. Chem. Phys.* **2018**, *148* (1), 011101. <https://doi.org/10.1063/1.5011798>.
- (5) Riplinger, C.; Sandhoefer, B.; Hansen, A.; Neese, F. Natural Triple Excitations in Local Coupled Cluster Calculations with Pair Natural Orbitals. *J. Chem. Phys.* **2013**, *139* (13), 134101. <https://doi.org/10.1063/1.4821834>.
- (6) Kuhn, B.; Guba, W.; Hert, J.; Banner, D.; Bissantz, C.; Ceccarelli, S.; Haap, W.; Körner, M.; Kuglstatler, A.; Lerner, C.; Mattei, P.; Neidhart, W.; Pinard, E.; Rudolph, M. G.; Schulz-Gasch, T.; Woltering, T.; Stahl, M. A Real-World Perspective on Molecular Design: Miniperspective. *J. Med. Chem.* **2016**, *59* (9), 4087–4102. <https://doi.org/10.1021/acs.jmedchem.5b01875>.
- (7) Al-Hamdani, Y. S.; Nagy, P. R.; Zen, A.; Barton, D.; Kállay, M.; Brandenburg, J. G.; Tkatchenko, A. Interactions between Large Molecules Pose a Puzzle for Reference Quantum Mechanical Methods. *Nat Commun* **2021**, *12* (1), 3927. <https://doi.org/10.1038/s41467-021-24119-3>.
- (8) Hawkins, P. C. D. Conformation Generation: The State of the Art. *J. Chem. Inf. Model.* **2017**, *57* (8), 1747–1756. <https://doi.org/10.1021/acs.jcim.7b00221>.
- (9) De Vivo, M.; Masetti, M.; Bottegoni, G.; Cavalli, A. Role of Molecular Dynamics and Related Methods in Drug Discovery. *J. Med. Chem.* **2016**, *59* (9), 4035–4061. <https://doi.org/10.1021/acs.jmedchem.5b01684>.
- (10) Shim, J.; MacKerell, Jr., A. D. Computational Ligand-Based Rational Design: Role of Conformational Sampling and Force Fields in Model Development. *Med. Chem. Commun.* **2011**, *2* (5), 356. <https://doi.org/10.1039/c1md00044f>.
- (11) Besora, M.; Braga, A. A. C.; Ujaque, G.; Maseras, F.; Lledós, A. The Importance of Conformational Search: A Test Case on the Catalytic Cycle of the Suzuki–Miyaura Cross-Coupling. *Theor Chem Acc* **2011**, *128* (4–6), 639–646. <https://doi.org/10.1007/s00214-010-0823-6>.
- (12) Harvey, J. N.; Himo, F.; Maseras, F.; Perrin, L. Scope and Challenge of Computational Methods for Studying Mechanism and Reactivity in Homogeneous Catalysis. *ACS Catal.* **2019**, *9* (8), 6803–6813. <https://doi.org/10.1021/acscatal.9b01537>.
- (13) Eisenstein, O.; Ujaque, G.; Lledós, A. What Makes a Good (Computed) Energy Profile? In *New Directions in the Modeling of Organometallic Reactions*; Lledós, A., Ujaque, G., Eds.; Topics in Organometallic Chemistry; Springer International Publishing: Cham, 2020; pp 1–38. [https://doi.org/10.1007/3418\\_2020\\_57](https://doi.org/10.1007/3418_2020_57).
- (14) Berlanga-Vázquez, A.; Talmazan, R. A.; Reyes-Mata, C. A.; Percástegui, E. G.; Flores-Alamo, M.; Podewitz, M.; Castillo, I. Conformational Effects of Regioisomeric Substitution on the Catalytic Activity of Copper/Calix[8]Arene C–S Coupling. *Eur J Inorg Chem* **2022**. <https://doi.org/10.1002/ejic.202200596>.

- (15) Talmazan, R. A.; Refugio Monroy, J.; del Río-Portilla, F.; Castillo, I.; Podewitz, M. Encapsulation Enhances the Catalytic Activity of C-N Coupling: Reaction Mechanism of a Cu(I)/Calix[8]Arene Supramolecular Catalyst. *ChemCatChem* **2022**, *14* (20). <https://doi.org/10.1002/cctc.202200662>.
- (16) Podewitz, M.; Sen, S.; Buchmeiser, M. R. On the Origin of E-Selectivity in the Ring-Opening Metathesis Polymerization with Molybdenum Imido Alkylidene N-Heterocyclic Carbene Complexes. *Organometallics* **2021**, *40* (15), 2478–2488. <https://doi.org/10.1021/acs.organomet.1c00229>.
- (17) Kolossváry, I.; Guida, W. C. Low Mode Search. An Efficient, Automated Computational Method for Conformational Analysis: Application to Cyclic and Acyclic Alkanes and Cyclic Peptides. *J. Am. Chem. Soc.* **1996**, *118* (21), 5011–5019. <https://doi.org/10.1021/ja952478m>.
- (18) Cole, J. C.; Korb, O.; McCabe, P.; Read, M. G.; Taylor, R. Knowledge-Based Conformer Generation Using the Cambridge Structural Database. *J. Chem. Inf. Model.* **2018**, *58* (3), 615–629. <https://doi.org/10.1021/acs.jcim.7b00697>.
- (19) Friedrich, N.-O.; Flachsenberg, F.; Meyder, A.; Sommer, K.; Kirchmair, J.; Rarey, M. Conformer: A Novel Method for the Generation of Conformer Ensembles. *J. Chem. Inf. Model.* **2019**, *59* (2), 731–742. <https://doi.org/10.1021/acs.jcim.8b00704>.
- (20) Watts, K. S.; Dalal, P.; Murphy, R. B.; Sherman, W.; Friesner, R. A.; Shelley, J. C. ConfGen: A Conformational Search Method for Efficient Generation of Bioactive Conformers. *J. Chem. Inf. Model.* **2010**, *50* (4), 534–546. <https://doi.org/10.1021/ci100015j>.
- (21) Gaalswyk, K.; Rowley, C. N. An Explicit-Solvent Conformation Search Method Using Open Software. *PeerJ* **2016**, *4*, e2088. <https://doi.org/10.7717/peerj.2088>.
- (22) Marx, D.; Hutter, J. *Ab Initio Molecular Dynamics: Basic Theory and Advanced Methods*, 1st ed.; Cambridge University Press, 2009. <https://doi.org/10.1017/CBO9780511609633>.
- (23) Bannwarth, C.; Ehlert, S.; Grimme, S. GFN2-XTB—An Accurate and Broadly Parametrized Self-Consistent Tight-Binding Quantum Chemical Method with Multipole Electrostatics and Density-Dependent Dispersion Contributions. *J. Chem. Theory Comput.* **2019**, *15* (3), 1652–1671. <https://doi.org/10.1021/acs.jctc.8b01176>.
- (24) Pracht, P.; Bohle, F.; Grimme, S. Automated Exploration of the Low-Energy Chemical Space with Fast Quantum Chemical Methods. *Phys. Chem. Chem. Phys.* **2020**, *22* (14), 7169–7192. <https://doi.org/10.1039/C9CP06869D>.
- (25) Podewitz, M. Towards Predictive Computational Catalysis – a Case Study of Olefin Metathesis with Mo Imido Alkylidene N-Heterocyclic Carbene Catalysts. In *Chemical Modelling: Volume 17*; Bahmann, H., Tremblay, J. C., Eds.; The Royal Society of Chemistry, 2022; Vol. 17, p 0. <https://doi.org/10.1039/9781839169342-00001>.
- (26) Barone, V.; Cossi, M. Quantum Calculation of Molecular Energies and Energy Gradients in Solution by a Conductor Solvent Model. *J. Phys. Chem. A* **1998**, *102* (11), 1995–2001. <https://doi.org/10.1021/jp9716997>.
- (27) Ehlert, S.; Stahn, M.; Spicher, S.; Grimme, S. Robust and Efficient Implicit Solvation Model for Fast Semiempirical Methods. *J. Chem. Theory Comput.* **2021**, *17* (7), 4250–4261. <https://doi.org/10.1021/acs.jctc.1c00471>.
- (28) Simm, G. N.; Türtscher, P. L.; Reiher, M. Systematic Microsolvation Approach with a Cluster-Continuum Scheme and Conformational Sampling. *Journal of Computational Chemistry* **2020**, *41* (12), 1144–1155. <https://doi.org/10.1002/jcc.26161>.
- (29) Steiner, M.; Holzknacht, T.; Schauerperl, M.; Podewitz, M. Quantum Chemical Microsolvation by Automated Water Placement. *Molecules* **2021**, *26* (6), 1793. <https://doi.org/10.3390/molecules26061793>.



- (30) Spicher, S.; Plett, C.; Pracht, P.; Hansen, A.; Grimme, S. Automated Molecular Cluster Growing for Explicit Solvation by Efficient Force Field and Tight Binding Methods. *J. Chem. Theory Comput.* **2022**, *18* (5), 3174–3189. <https://doi.org/10.1021/acs.jctc.2c00239>.
- (31) Tian, C.; Kasavajhala, K.; Belfon, K. A. A.; Raguette, L.; Huang, H.; Migués, A. N.; Bickel, J.; Wang, Y.; Pincay, J.; Wu, Q.; Simmerling, C. Ff19SB: Amino-Acid-Specific Protein Backbone Parameters Trained against Quantum Mechanics Energy Surfaces in Solution. *J. Chem. Theory Comput.* **2020**, *16* (1), 528–552. <https://doi.org/10.1021/acs.jctc.9b00591>.
- (32) Dickson, C. J.; Walker, R. C.; Gould, I. R. Lipid21: Complex Lipid Membrane Simulations with AMBER. *J. Chem. Theory Comput.* **2022**, *18* (3), 1726–1736. <https://doi.org/10.1021/acs.jctc.1c01217>.
- (33) Galindo-Murillo, R.; Robertson, J. C.; Zgarbová, M.; Šponer, J.; Otyepka, M.; Jurečka, P.; Cheatham, T. E. Assessing the Current State of Amber Force Field Modifications for DNA. *J. Chem. Theory Comput.* **2016**, *12* (8), 4114–4127. <https://doi.org/10.1021/acs.jctc.6b00186>.
- (34) Madarász, Á.; Berta, D.; Paton, R. S. Development of a True Transition State Force Field from Quantum Mechanical Calculations. *J. Chem. Theory Comput.* **2016**, *12* (4), 1833–1844. <https://doi.org/10.1021/acs.jctc.5b01237>.
- (35) Rosales, A. R.; Quinn, T. R.; Wahlers, J.; Tomberg, A.; Zhang, X.; Helquist, P.; Wiest, O.; Norrby, P.-O. Application of Q2MM to Predictions in Stereoselective Synthesis. *Chem. Commun.* **2018**, *54* (60), 8294–8311. <https://doi.org/10.1039/C8CC03695K>.
- (36) Wang, J.; Wolf, R. M.; Caldwell, J. W.; Kollman, P. A.; Case, D. A. Development and Testing of a General Amber Force Field. *J. Comput. Chem.* **2004**, *25* (9), 1157–1174. <https://doi.org/10.1002/jcc.20035>.
- (37) MacKerell, A. D.; Bashford, D.; Bellott, M.; Dunbrack, R. L.; Evanseck, J. D.; Field, M. J.; Fischer, S.; Gao, J.; Guo, H.; Ha, S.; Joseph-McCarthy, D.; Kuchnir, L.; Kuczera, K.; Lau, F. T. K.; Mattos, C.; Michnick, S.; Ngo, T.; Nguyen, D. T.; Prodhom, B.; Reiher, W. E.; Roux, B.; Schlenkrich, M.; Smith, J. C.; Stote, R.; Straub, J.; Watanabe, M.; Wiórkiewicz-Kuczera, J.; Yin, D.; Karplus, M. All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins. *J. Phys. Chem. B* **1998**, *102* (18), 3586–3616. <https://doi.org/10.1021/jp973084f>.
- (38) van Gunsteren, W. F.; Daura, X.; Mark, A. E. GROMOS Force Field. In *Encyclopedia of Computational Chemistry*; von Ragué-Schleyer, P., Allinger, N. L., Clark, T., Gasteiger, J., Kollman, P. A., Schaefer, H. F., Schreiner, P. R., Eds.; John Wiley & Sons, Ltd: Chichester, UK, 2002; p cga011. <https://doi.org/10.1002/0470845015.cga011>.
- (39) Jorgensen, W. L.; Maxwell, D. S.; Tirado-Rives, J. Development and Testing of the OPLS All-Atom Force Field on Conformational Energetics and Properties of Organic Liquids. *J. Am. Chem. Soc.* **1996**, *118* (45), 11225–11236. <https://doi.org/10.1021/ja9621760>.
- (40) *Development and Benchmarking of Open Force Field v1.0.0—the Parsley Small-Molecule Force Field | Journal of Chemical Theory and Computation.* <https://pubs.acs.org/doi/10.1021/acs.jctc.1c00571> (accessed 2023-05-16).
- (41) Wang, J.; Wang, W.; Kollman, P. A.; Case, D. A. Automatic Atom Type and Bond Type Perception in Molecular Mechanical Calculations. *Journal of Molecular Graphics and Modelling* **2006**, *25* (2), 247–260. <https://doi.org/10.1016/j.jmglm.2005.12.005>.
- (42) Peters, M. B.; Yang, Y.; Wang, B.; Füsti-Molnár, L.; Weaver, M. N.; Merz, K. M. Structural Survey of Zinc-Containing Proteins and Development of the Zinc AMBER Force Field (ZAFF). *J. Chem. Theory Comput.* **2010**, *6* (9), 2935–2947. <https://doi.org/10.1021/ct1002626>.
- (43) Li, P.; Roberts, B. P.; Chakravorty, D. K.; Merz, K. M. Rational Design of Particle Mesh Ewald Compatible Lennard-Jones Parameters for +2 Metal Cations in Explicit Solvent. *J. Chem. Theory Comput.* **2013**, *9* (6), 2733–2748. <https://doi.org/10.1021/ct400146w>.

- (44) Li, P.; Merz, K. M. Taking into Account the Ion-Induced Dipole Interaction in the Nonbonded Model of Ions. *J. Chem. Theory Comput.* **2014**, *10* (1), 289–297. <https://doi.org/10.1021/ct400751u>.
- (45) Aqvist, J.; Warshel, A. Free Energy Relationships in Metalloenzyme-Catalyzed Reactions. Calculations of the Effects of Metal Ion Substitutions in Staphylococcal Nuclease. *J. Am. Chem. Soc.* **1990**, *112* (8), 2860–2868. <https://doi.org/10.1021/ja00164a003>.
- (46) Grossfield, A.; Ren, P.; Ponder, J. W. Ion Solvation Thermodynamics from Simulation with a Polarizable Force Field. *J. Am. Chem. Soc.* **2003**, *125* (50), 15671–15682. <https://doi.org/10.1021/ja037005r>.
- (47) Frenking, G.; Fröhlich, N. The Nature of the Bonding in Transition-Metal Compounds. *Chem. Rev.* **2000**, *100* (2), 717–774. <https://doi.org/10.1021/cr980401l>.
- (48) Case, D. A.; Ben-Shalom, I. Y.; Brozell, S. R.; Cerutti, D. S.; Cheatham III, T. E.; Cruzeiro, V. W. D.; Darden, T. A.; Duke, R. E.; Ghoreishi, D.; Gilson, M. K. AMBER20. *University of California, San Francisco* **2020**.
- (49) Li, P.; Merz, K. M. MCPB.Py: A Python Based Metal Center Parameter Builder. *J. Chem. Inf. Model.* **2016**, *56* (4), 599–604. <https://doi.org/10.1021/acs.jcim.5b00674>.
- (50) Kastner, D. W.; Nandy, A.; Mehmood, R.; Kulik, H. J. Mechanistic Insights into Substrate Positioning That Distinguish Non-Heme Fe(II)/ $\alpha$ -Ketoglutarate-Dependent Halogenases and Hydroxylases. *ACS Catal.* **2023**, *13* (4), 2489–2501. <https://doi.org/10.1021/acscatal.2c06241>.
- (51) Piskorz, T. K.; Martí-Centelles, V.; Young, T. A.; Lusby, P. J.; Duarte, F. Computational Modeling of Supramolecular Metallo-Organic Cages—Challenges and Opportunities. *ACS Catal.* **2022**, 5806–5826. <https://doi.org/10.1021/acscatal.2c00837>.
- (52) Neese, F.; Wennmohs, F.; Becker, U.; Riplinger, C. The ORCA Quantum Chemistry Program Package. *J. Chem. Phys.* **2020**, *152* (22), 224108. <https://doi.org/10.1063/5.0004608>.
- (53) Lu, T.; Chen, F. Multiwfn: A Multifunctional Wavefunction Analyzer. *J. Comput. Chem.* **2012**, *33* (5), 580–592. <https://doi.org/10.1002/jcc.22885>.
- (54) Singh, U. C.; Kollman, P. A. An Approach to Computing Electrostatic Charges for Molecules. *J. Comput. Chem.* **1984**, *5* (2), 129–145. <https://doi.org/10.1002/jcc.540050204>.
- (55) Bayly, C. I.; Cieplak, P.; Cornell, W.; Kollman, P. A. A Well-Behaved Electrostatic Potential Based Method Using Charge Restraints for Deriving Atomic Charges: The RESP Model. *J. Phys. Chem.* **1993**, *97* (40), 10269–10280. <https://doi.org/10.1021/j100142a004>.
- (56) Götz, A. W.; Williamson, M. J.; Xu, D.; Poole, D.; Le Grand, S.; Walker, R. C. Routine Microsecond Molecular Dynamics Simulations with AMBER on GPUs. 1. Generalized Born. *J. Chem. Theory Comput.* **2012**, *8* (5), 1542–1555. <https://doi.org/10.1021/ct200909j>.
- (57) Salomon-Ferrer, R.; Götz, A. W.; Poole, D.; Le Grand, S.; Walker, R. C. Routine Microsecond Molecular Dynamics Simulations with AMBER on GPUs. 2. Explicit Solvent Particle Mesh Ewald. *J. Chem. Theory Comput.* **2013**, *9* (9), 3878–3888. <https://doi.org/10.1021/ct400314y>.
- (58) Roe, D. R.; Cheatham, T. E. PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular Dynamics Trajectory Data. *J. Chem. Theory Comput.* **2013**, *9* (7), 3084–3095. <https://doi.org/10.1021/ct400341p>.
- (59) Tarjan, R. Depth-First Search and Linear Graph Algorithms. In *12th Annual Symposium on Switching and Automata Theory (swat 1971)*; IEEE: East Lansing, MI, USA, 1971; pp 114–121. <https://doi.org/10.1109/SWAT.1971.10>.
- (60) Kahn, L. R.; Goddard, W. A., III. Ab Initio Effective Potentials for Use in Molecular Calculations. *The Journal of Chemical Physics* **1972**, *56* (6), 2685–2701. <https://doi.org/10.1063/1.1677597>.
- (61) Wallnoefer, H. G.; Liedl, K. R.; Fox, T. A Challenging System: Free Energy Prediction for Factor Xa. *J. Comput. Chem.* **2011**, *32* (8), 1743–1752. <https://doi.org/10.1002/jcc.21758>.
- (62) Becke, A. D. Density-Functional Exchange-Energy Approximation with Correct Asymptotic Behavior. *Phys. Rev. A* **1988**, *38* (6), 3098–3100. <https://doi.org/10.1103/PhysRevA.38.3098>.

- (63) Perdew, J. P. Density-Functional Approximation for the Correlation Energy of the Inhomogeneous Electron Gas. *Phys. Rev. B* **1986**, *33* (12), 8822–8824. <https://doi.org/10.1103/PhysRevB.33.8822>.
- (64) Caldeweyher, E.; Ehlert, S.; Hansen, A.; Neugebauer, H.; Spicher, S.; Bannwarth, C.; Grimme, S. A Generally Applicable Atomic-Charge Dependent London Dispersion Correction. *The Journal of Chemical Physics* **2019**, *150* (15), 154122. <https://doi.org/10.1063/1.5090222>.
- (65) Weigend, F.; Ahlrichs, R. Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn: Design and Assessment of Accuracy. *Phys. Chem. Chem. Phys.* **2005**, *7* (18), 3297–3305. <https://doi.org/10.1039/B508541A>.
- (66) Herz, K.; Podewitz, M.; Stöhr, L.; Wang, D.; Frey, W.; Liedl, K. R.; Sen, S.; Buchmeiser, M. R. Mechanism of Olefin Metathesis with Neutral and Cationic Molybdenum Imido Alkylidene N-Heterocyclic Carbene Complexes. *J. Am. Chem. Soc.* **2019**, *141* (20), 8264–8276. <https://doi.org/10.1021/jacs.9b02092>.
- (67) Buchmeiser, M. R.; Sen, S.; Unold, J.; Frey, W. N-Heterocyclic Carbene, High Oxidation State Molybdenum Alkylidene Complexes: Functional-Group-Tolerant Cationic Metathesis Catalysts. *Angewandte Chemie International Edition* **2014**, *53* (35), 9384–9388. <https://doi.org/10.1002/anie.201404655>.
- (68) Kieninger, C.; Wurst, K.; Podewitz, M.; Stanley, M.; Deery, E.; Lawrence, A. D.; Liedl, K. R.; Warren, M. J.; Kräutler, B. Replacement of the Cobalt Center of Vitamin B12 by Nickel: Nibalamin and Nibyrinic Acid Prepared from Metal-Free B12 Ligands Hydrogenobalamin and Hydrogenobyric Acid. *Angewandte Chemie International Edition* **2020**, *59* (45), 20129–20136. <https://doi.org/10.1002/anie.202008407>.
- (69) Kieninger, C.; Baker, J. A.; Podewitz, M.; Wurst, K.; Jockusch, S.; Lawrence, A. D.; Deery, E.; Gruber, K.; Liedl, K. R.; Warren, M. J.; Kräutler, B. Zinc Substitution of Cobalt in Vitamin B12: Zincobyric Acid and Zincobalamin as Luminescent Structural B12-Mimics. *Angewandte Chemie International Edition* **2019**, *58* (41), 14568–14572. <https://doi.org/10.1002/anie.201908428>.
- (70) Kieninger, C.; Deery, E.; Lawrence, A. D.; Podewitz, M.; Wurst, K.; Nemoto-Smith, E.; Widner, F. J.; Baker, J. A.; Jockusch, S.; Kreutz, C. R.; Liedl, K. R.; Gruber, K.; Warren, M. J.; Kräutler, B. The Hydrogenobyric Acid Structure Reveals the Corrin Ligand as an Entatic State Module Empowering B12 Cofactors for Catalysis. *Angewandte Chemie International Edition* **2019**, *58* (31), 10756–10760. <https://doi.org/10.1002/anie.201904713>.
- (71) Kräutler, B. 2.11 - Cobalt Enzymes. In *Comprehensive Inorganic Chemistry III (Third Edition)*; Reedijk, J., Poepelmeier, K. R., Eds.; Elsevier: Oxford, 2023; pp 268–301. <https://doi.org/10.1016/B978-0-12-823144-9.00146-1>.
- (72) Adamo, C.; Barone, V. Toward Reliable Density Functional Methods without Adjustable Parameters: The PBE0 Model. *The Journal of Chemical Physics* **1999**, *110* (13), 6158–6170. <https://doi.org/10.1063/1.478522>.
- (73) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. A Consistent and Accurate Ab Initio Parametrization of Density Functional Dispersion Correction (DFT-D) for the 94 Elements H-Pu. *J Chem Phys* **2010**, *132* (15), 154104. <https://doi.org/10.1063/1.3382344>.
- (74) *On the Origin of E-Selectivity in the Ring-Opening Metathesis Polymerization with Molybdenum Imido Alkylidene N-Heterocyclic Carbene Complexes | Organometallics.* <https://pubs.acs.org/doi/10.1021/acs.organomet.1c00229> (accessed 2023-04-25).
- (75) Grimme, S.; Ehrlich, S.; Goerigk, L. Effect of the Damping Function in Dispersion Corrected Density Functional Theory. *Journal of Computational Chemistry* **2011**, *32* (7), 1456–1465. <https://doi.org/10.1002/jcc.21759>.