

Synthesis of Nickel(I)–Bromide Complexes via Oxidation and Ligand Displacement: Evaluation of Ligand Effects on Speciation and Reactivity

Samuel H. Newman-Stonebraker,[‡] T. Judah Raab,[‡] Hootan Roshandel, Abigail G. Doyle^{*}

ABSTRACT: Nickel's +1 oxidation state has received much interest due to its varied and often enigmatic behavior in increasingly popular catalytic methods. In part, the lack of understanding about Ni^I results from common synthetic strategies limiting the breadth of complexes that are accessible for mechanistic study and catalyst design. We report an oxidative approach using tribromide salts that allows for the generation of a well-defined precursor, [Ni^I(COD)Br]₂, as well as several new Ni^I complexes. Included among them are complexes bearing bulky monophosphines, for which structure–speciation relationships are established and catalytic reactivity in a Suzuki–Miyaura coupling (SMC) is investigated. Notably, these routes also allow for the synthesis of well-defined monomeric *t*-Bu^{bpy}-bound Ni^I complexes, which has not previously been achieved. These complexes, which react with aryl halides, can enable previously challenging mechanistic investigations and present new opportunities for catalysis and synthesis.

INTRODUCTION

Nickel-catalyzed cross-couplings have emerged as broadly useful methodologies in synthetic organic chemistry—in large part due to the complementary reactivity of Ni to precious metals like Pd.^{1,2} In addition to Ni's ability to facilitate challenging oxidative addition reactions,^{3,4} its relative propensity⁵ to access open-shell electron configurations (i.e., Ni^I & Ni^{III}) underlies many modern cross-coupling reactions (e.g., Ni/photoredox,⁶ Ni/electrocatalysis,⁷ cross-electrophile coupling⁸). Conversely, this same property can attenuate turnover in traditional Ni⁰/Ni^{II} cycles, where Ni^I species have been identified as off-cycle.⁹ It is therefore of great interest to understand the nuanced factors that control the formation, reactivity, and speciation of Ni^I intermediates.^{10,11}

Stoichiometric synthesis of well-defined Ni^I complexes presents an opportunity to investigate these nuances and characterize species that are relevant in catalysis. Examples of such investigations have examined several ligand classes on Ni^I centers: Matsubara¹² and Schoenebeck¹³ have studied *N*-heterocyclic carbene (NHC) ligands, Hazari^{9,14–16} and Schoenebeck¹⁷ bisphosphine ligands, and Hazari^{18,19} and Martin^{19,20} phenanthroline ligands. For each ligand class, tractable conclusions about the role(s) of Ni^I species in the respective catalytic mechanisms have been put forth. However, existing strategies for synthesizing Ni^I species (Figure 1A) limit the breadth of complexes that can be accessed with catalytically relevant ligands. We postulate two contributing factors therein: (1) common preparations for Ni^I species necessitate the formation of L_nNi^{II}X₂ complexes that are intractable for bulky ligands²¹ and (2) components of [Ni⁰] and [Ni^{II}] precursors used for Ni^I synthesis can induce undesired redox equilibria and speciation effects.^{20,22} These limitations are borne out in a literature survey of Ni^I

complexes: certain structural motifs are well-explored while others remain elusive (Figure 1B).

Bulky monophosphines are among the ligand classes that have scarcely been characterized on Ni^I centers. Notably, small monophosphine ligands have been known to form monomeric (PR₃)_nNi^IX (n = 2 or 3) species via comproportionation and oxidative addition,^{23,24} but dimeric [(PR₃)Ni^IX]₂ complexes bearing larger phosphine congeners were unknown prior to a 2022 report from our group.^{25,26} This is in stark contrast to precedent for Pd^I, for which complexes of the type [(PR₃)Pd^IX]₂ are long-known²⁷ and have been thoroughly studied by Schoenebeck and coworkers.^{28,29} As precedent for bisligated (PR₃)₂Ni^I species has shown that such species are capable of catalytic cycle (re)entry,^{9,14–16} the structure and speciation of monoligated (PR₃)Ni^I complexes warrants further investigation.

Moreover, while 4,4'-di-*tert*-butylbipyridine (*t*-Bu^{bpy}) is the ligand of choice for many Ni-catalyzed cross-couplings that invoke open-shell intermediates,^{30–33} only two examples of formal (*t*-Bu^{bpy})Ni^I species have been structurally characterized: Hazari's [(*t*-Bu^{bpy})Ni^ICl]₂ and Nocera's [(*t*-Bu^{bpy})Ni^{1.5+}(quinuclidine)Cl]₂Cl.^{18,34} Importantly, these complexes do not replicate the behavior of (*t*-Bu^{bpy})Ni^I species in catalysis due to irreversible dimerization of reactive (*t*-Bu^{bpy})Ni^IX. While *in situ* generation of monomeric (*t*-Bu^{bpy})Ni^I species has clarified the reactivity of important catalytic intermediates,^{35,36} these approaches are more challenging to apply to future catalyst design and mechanistic studies than stoichiometric synthesis.

Strategies to enable the synthesis of previously unknown Ni^I complexes create downstream opportunities for more effective Ni catalysis. Contributions from Morandi and coworkers, who recently reported a phenoxide-bound Ni^I precursor,³⁷ have already led to mechanistic insight and development in Ni-catalyzed methodologies.³⁸ Further goals

for the synthetic organometallic community in this area include: the complexation of catalytically relevant ligands to structurally novel Ni^I centers, the incorporation of easily derivatized X-type ligands such as halides, and application of synthetic discoveries to catalytic systems.

Herein, we present the synthesis of over 20 previously unidentified Ni^I complexes bearing olefins and catalytically relevant monophosphines and bipyridine ligands. We access these complexes through a mild one-pot oxidation or via ligand displacement of well-defined Ni^I precursors. For the bulky monophosphine class, we conduct a structure-speciation analysis and catalytic studies to elucidate these ligands' behavior on Ni^I centers. For the bipyridine class, we identify the first examples of monomeric (^t-Bu₂bpy)Ni^I complexes and investigate their electronic structure and stoichiometric reactivity with aryl halides.





Figure 1. (A) Ni^I complex synthetic strategies and their prevalence. (B) Examples of structurally characterized Ni^I complexes for (I) phosphine ligands, (II) bipyridine/phenanthroline ligands, and (III) olefin ligands.

RESULTS AND DISCUSSION


In our recent study of dialkylbiaryl phosphine ligands in Ni catalysis, we serendipitously identified the structure of a unique monophosphine-bound Ni^I halide dimer.²⁵ This species, [(Cy)JohnPhos]Ni^ICl₂ (**1-Cl**), formed in trace amounts following oxidative addition of 2-chlorotoluene to

(Cy)JohnPhos)₂Ni⁰. To the best of our knowledge, **1-Cl** was the first structurally characterized example of a [(PR₃)Ni^IX]₂ dimer. To further investigate the role of **1** in the catalytic system of interest, we sought to synthesize this complex independently by comproportionation. However, comproportionation attempts with several Ni⁰ and Ni^{II} precursors failed, perhaps due to challenges forming phosphine-bound Ni^{II}Cl₂ species. Taking inspiration from studies from Fout³⁹ and Uyeda⁴⁰, we were able to isolate [(Cy)JohnPhos]Ni^IBr₂ (**1-Br**) by oxidizing (Cy)JohnPhos)₂Ni⁰ with trimethylphenylammonium tribromide (TMPhATB), a commercially available, easily manipulated solid (Figure 2B). **1-Br** assumes an analogous solid-state structure to **1-Cl**, with κ¹-P phosphine binding, μ₂-Br ligand bridging, and a Ni-Ni bonding interaction (2.5955(4) Å).


A. Ni^I with bulky phosphines inaccessible via comproportionation



B. Synthesis of [(Cy)JohnPhos]Ni^IBr₂ using tribromide oxidant



C. Synthesis of (XPhos)Ni^IBr from Ni(COD)₂



D. SCXRD and EPR characterization of **1-Br** and **2**




Figure 2. (A) Attempted syntheses of **1-Cl** and **1-Br** via comproportionation (X = Cl or Br). (B) Initial discovery of tribromide oxidation to generate **1-Br**. (C) One-pot oxidation from Ni(COD)₂ to give **2**. (D) Solid state structures of **1-Br** and **2**, with thermal ellipsoids displayed at 50% probability and hydrogen atoms omitted for clarity. Selected solid-state bond distances are tabulated. EPR simulation parameters: $g_1 = 2.306$, $g_2 = 2.138$, $g_3 = 2.039$. For simulated hyperfine splitting and strain, see SI.

Seeking to extend this oxidative approach to bulkier Buchwald phosphines, we treated Ni(COD)₂ (COD = 1,5-cyclooctadiene) with TPhATB in the presence of XPhos, generating (XPhos)Ni^IBr (**2**). In the solid state, **2** is monomeric, with XPhos adopting a κ¹-P, η²-C_{arene} binding mode. This speciation contrasts with both **1-Cl** and **1-Br**, in which CyJohnPhos binds κ¹-P only in a dimeric complex. EPR measurements in glassy toluene at 77 K suggest that the solid-state speciation is conserved in the solution phase: no signal is observed for **1-Br**, while spectra of **2** feature a rhombic signal with large hyperfine splitting from the ³¹P nucleus of XPhos (Figure 2D). The low solubility of its solution magnetic moment. However, **2** was observed to have a solution-phase magnetic moment of 1.73 μ_B, consistent with a monomeric d⁹ species.⁴¹

Having identified tribromide oxidation as a tenable route to Ni^I species, we sought to evaluate the scope of ligands that could be complexed to Ni^I centers. We were able to generate Ni^I halide complexes for a variety of catalytically important ligand classes including mono- and bisphosphine, *N*-heterocyclic carbene (NHC), and polypyridyl (Table 1). In most cases, yields are comparable to other reported syntheses,^{4,13,42–44} but the unique generality of Br₃[−] oxidation is highly enabling.

Table 1. Ligand generality of tribromide oxidation

Ligand	<i>n</i>	Ni ^I complex	Br ₃ [−] oxidation yield (isolated)
PPh ₃	3	L ₃ Ni ^I Br	75%
PCy ₃	2	L ₂ Ni ^I Br	67%
dppf	1	LNi ^I Br/[LNi ^I Br] ₂	77%
<i>t</i> -BuXantPhos	1	LNi ^I Br	48%
IPr	1	[LNi ^I Br] ₂	69%
tpy	1	LNi ^I Br	42%
dtbbpy	1	[LNi ^I Br] ₂	66% ^a

^aNMR yield using displaced COD as an internal standard

However, challenges arose in the isolation of Ni^I complexes bearing two ligand classes: bipyridines and bulky trialkylphosphines. The former case will be discussed in detail (*vide infra*). Our challenges with the latter case are best summarized with the product mixture resulting from the oxidation of a Ni(COD)₂/P(*t*-Bu)₃ mixture. The desired product, a dimeric [(P(*t*-Bu)₃)Ni^IBr]₂ complex, was not detected. Instead, this reaction generated two unanticipated P(*t*-Bu)₃-bound species: an over-oxidized Ni^{1.5+} dimer, [(P(*t*-Bu)₃)₂Ni^{1.5+}₂Br₃] (**3**), and a nickelate complex with an outer-sphere ammonium cation, [(P(*t*-Bu)₃)Ni^IBr₂]Me₃PhN (**4**) (Figure 3). **3** likely forms as a product along with Ni⁰ black in a redox equilibrium with the expected Ni^I dimer, while **4** is presumably the product of ammonium bromide coordination to a (P(*t*-Bu)₃)Ni^IBr species.

While complexes **3** and **4** are interesting from a structural and electronic perspective, we sought an alternative route to the desired [(P(*t*-Bu)₃)Ni^IBr]₂ complex that did not

employ ammonium salts. To this end, we identified that the formation of complexes **2**, **3**, and **4** implies a reactive intermediate from which ligation of these bulky phosphines is relatively facile. Both XPhos and P(*t*-Bu)₃ are too sterically encumbered to displace COD from Ni(COD)₂ directly,²⁵ but both ligate to Ni following tribromide oxidation. We therefore aimed to isolate this putative intermediate, which could serve as an ammonium-free Ni^I precursor.

Synthesis and solid-state structures of P(*t*-Bu)₃/Ni(COD)₂ oxidation products




Figure 3. Synthesis and solid state structures of **3** and **4**, with thermal ellipsoids displayed at 50% probability and hydrogen atoms omitted for clarity. Selected bond distances (Å) for **3**: Ni1–Ni2: 2.378(1); Ni1–P1: 2.306(1); Ni1–Br1: 2.4637(1). Selected bond distances (Å) for **4**: Ni3–P3: 2.209(1); Ni3–Br4: 2.4096(7).

Following tribromide oxidation of a Ni(COD)₂ solution in THF-*d*₈, we observed the formation of a single paramagnetic species by ¹H NMR. Use of a more soluble tribromide oxidant, tetrabutylammonium tribromide (TBATB),⁴⁵ cooling, and addition of solvent-quantity free COD to the oxidation mixture stabilized the resulting complex. With these modifications, we were able to isolate and characterize [Ni^I(COD)Br]₂ (**5**), a golden yellow solid that is stable at –35 °C under inert atmosphere (Figure 4).^{46,47} **5** is a rare example of a Ni^I–olefin complex and a tractable precursor to a variety of Ni^I–Br complexes (*vide infra*).

Synthesis and solid-state structure of [Ni^I(COD)Br]₂




Figure 4. Synthesis, selected interatomic distances, and solid state structure of **5**. Thermal ellipsoids are displayed at 50% probability and hydrogen atoms are omitted for clarity. Selected solid-state bond distances are tabulated.

Structure–Speciation Relationships of Monodentate Phosphines with [Ni^I(COD)Br]₂. With access to [Ni^I(COD)Br]₂, we studied its ability to serve as a precursor

to generate L_nNi^I complexes with bulky monophosphines. Unlike with $Ni(COD)_2$,²⁵ Buchwald-type ligands of all steric profiles were capable of displacing the COD ligands of $[Ni^I(COD)Br]_2$ to generate the desired Ni^I complex. In addition to **1-Br** and **2** with CyJohnPhos and XPhos, respectively, Ni^I complexes with SPhos, DavePhos, JohnPhos, and *t*-BuBrettPhos formed readily from **5**; each of these complexes was characterized by SCXRD (see SI). For Buchwald-type ligands, three distinct classes of L_1Ni^I species were observed in the solid-state: (i) dimers with Ni–Ni bonds and no η^2 -C_{arene} interaction, (ii) dimers with no Ni–Ni bond (distances >3.0 Å) and a strong η^2 -C_{arene} interaction between each Ni and the ligand B ring, and (iii) monomers with a strong η^2 -C_{arene} interaction to the B ring (Figure 5). The ligand structural feature that most clearly leads to the observed speciation outcome is the presence of a 4-*i*-Pr group on the B ring, which prevents dimer formation altogether. The structural feature(s) that distinguishes the two classes of dimers is less clear, though only the smaller Buchwald-type ligands formed species with Ni–Ni bonds.




Figure 5. (A) Buchwald ligand-bound Ni^I complex binding modes in the solid state. (B) Solid-state structure of $[(JohnPhos)Ni^I]_2$ (**6**) with thermal ellipsoids displayed at 50% probability and hydrogen atoms omitted for clarity.

Given the success of bulky Buchwald-type ligands at displacing COD from **5**, we reexamined the synthesis of bulky trialkylphosphine-bound Ni^I complexes using the NR_4Br salt-free precursor (Figure 6). We were pleased to find that reaction of **5** with 2 equiv of $P(t-Bu)_3$ (1:1 L:Ni) led to the formation of $[(P(t-Bu)_3)Ni^I]_2$ (**7**). The X-ray crystal structure of **7** confirmed its identity as a Ni^I dimer, with a Ni–Ni bond distance (2.6005(6) Å) similar to that observed in **1-Br**. We found that $P(t-Bu)_3$:Ni ratios >1:1 did not affect the speciation of the resulting complexes, with only the L_1Ni^I dimer complex observed.

Recent work by our lab and the Sigman lab has found that minimum percent buried volume ($\%V_{bur}(min)$)—a steric quantification of the smallest energetically accessible conformation of a ligand within 3.5 Å of the metal center—enables the discovery of structure–speciation relationships of phosphine ligand/metal complexes in cross-coupling.⁴⁸ With the demonstrated success of $\%V_{bur}(min)$ in rationalizing phosphine ligand effects at Ni^0/Ni^{II} , we were curious to investigate similar effects at Ni^I .

For ligands with $\%V_{bur}(min)$ values⁴⁹ slightly lower than $P(t-Bu)_3$ (36.3%), such as $CyP(t-Bu)_2$ (34.3%) and $Cy_2P(t-Bu)$ (32.0%), both the L_1Ni^I dimer⁵⁰ (**8** (L = $CyP(t-Bu)_2$) and **9** (L = $Cy_2P(t-Bu)$)) and L_2Ni^I monomer species (**10** (L = $CyP(t-Bu)_2$) and **11** (L = $Cy_2P(t-Bu)$)) could be generated and were structurally characterized, with the L:Ni stoichiometric ratio controlling the outcome.⁵¹ While these ligands normally do not form L_2Ni complexes at Ni^0 or Ni^{II} , the relatively small size of the single halide ligand leaves the majority of Ni's coordination sphere unencumbered, allowing the coordination of large phosphines. The crystal structure of $[(CyP(t-Bu)_2)_2Ni^I]Br$ revealed a nearly T-shaped complex, as the high amount of steric pressure between the bulky phosphines distorted the complex from the ideal trigonal geometry. For ligands smaller than $P(t-Bu)_3$ where more than one Ni^I species can form, L:Ni stoichiometry controlled the outcome for isolated material in the solid-state. These trends were generally conserved in solution-state NMR characterization, though a small amount of phosphine dissociation and dimerization was observed for L_2Ni^I monomer **10** with $CyP(t-Bu)_2$ (see SI).

We also found that PCy_3 ($\%V_{bur}(min) = 30.2\%$) could form a L_1Ni^I dimer (**12** (L = PCy_3)) from **5** with a 1:1 ratio of phosphine:Ni. In the presence of excess COD, the PCy_3 complex is more prone to disproportionation—generating $Ni(COD)_2$ and $(PCy_3)_2Ni^{II}Br_2$ —than ligands with higher $\%V_{bur}(min)$ values. This is unsurprising given that stable L_2Ni^0 and L_2Ni^{II} complexes can readily form with PCy_3 . Nonetheless, the ability to synthesize $[(PCy_3)Ni^I]_2$ differs from the ligand structure–speciation trends observed in generation of the analogous Pd^I dimers.²⁹ Like $CyP(t-Bu)_2$ and $Cy_2P(t-Bu)$, the L_2Ni^I Br monomer (**13** (L = PCy_3)) was readily obtained with PCy_3 .

For even smaller PPh_3 ($\%V_{bur}(min) = 28.2\%$), Ni^I species were obtainable for 3:1, 2:1, and 1:1 ratios of phosphine:Ni. Unlike the examined phosphines with greater $\%V_{bur}(min)$ values, a L_3Ni^I (**14** (L = PPh_3)) monomer is sterically accessible for PPh_3 . The L_2Ni^I (**15** (L = PPh_3)) monomer is also isolable. Uniquely, treatment of **5** with only one equivalent of PPh_3 relative to Ni resulted in the formation of a rare monomeric COD-bound Ni^I complex, $(PPh_3)Ni^I(COD)Br$ (**16**) (Figure 6B).⁵² Similar to COD-bound **5**, complex **16** rapidly decomposes in solution in the absence of added COD or ligand.




Figure 6. (A) Monophosphine structure–speciation relationships. (B) Synthesis and characterization of $(\text{PPh}_3)\text{Ni}^{\text{I}}(\text{COD})\text{Br}$. EPR simulation parameters: $g_1 = 2.593$, $g_2 = 2.075$, $g_3 = 2.027$. For simulated hyperfine splitting and strain, see SI. (C) Solid-state structures of selected $\text{L}_1\text{Ni}^{\text{I}}$ dimers, with thermal ellipsoids displayed at 50% probability and hydrogen atoms omitted for clarity. (D) Solid-state structures of selected $\text{L}_2\text{Ni}^{\text{I}}$ monomers, with thermal ellipsoids displayed at 50% probability and hydrogen atoms omitted for clarity.

Overall, we have found that $[\text{Ni}^{\text{I}}(\text{COD})\text{Br}]_2$ is a versatile precursor for synthesizing monophosphine-bound Ni^{I} complexes. The high lability of COD in **5** enables formation of otherwise challenging-to-access complexes with bulky phosphines such as Buchwald-type ligands and $\text{P}(t\text{-Bu})_3$. Furthermore, the excellent stoichiometric control on product outcome when using **5** as a precursor allows for a detailed understanding of phosphine structural effects on Ni^{I} speciation (Figure 6).

Implications of Bulky Monophosphine Speciation at Ni^{I} in Cross-Coupling. Recently, in a collaborative study with the Sigman group, we identified phosphine ligand reactivity thresholds in Ni - and Pd -catalyzed cross-coupling datasets;⁴⁸ these reactivity thresholds were linked to the % V_{bur} (min) steric descriptor.⁴⁹ For Ni , only monodentate phosphines with % V_{bur} (min) values less than 32% were effective at catalyzing the studied Suzuki–Miyaura coupling (SMC) reactions. This value corresponded to the region of chemical space where two phosphines could bind and stabilize Ni^0 and Ni^{II} complexes; these even oxidation states of Ni are believed to be the catalytically relevant species in SMC reactions.⁹ Given Ni 's propensity to engage in unproductive side pathways, we hypothesized that attaining bisligated $\text{Ni}^0/\text{Ni}^{\text{II}}$ complexes was necessary to stabilize these on-cycle species.⁵³ The inability of bulky monodentate phosphines (i.e., with % V_{bur} (min) values greater than 32%) to adequately stabilize these species would leave Ni more prone to falling into off-cycle thermodynamic sinks. However, this hypothesis has eluded testing due to previous synthetic challenges in accessing Ni complexes of any oxidation

state with bulky monodentate phosphines. Indeed, ligands like $\text{P}(t\text{-Bu})_3$ do not displace olefin ligands from Ni^0 precursors like $\text{Ni}(\text{COD})_2$,²⁵ nor do they ligate $\text{Ni}^{\text{II}}\text{X}_2$ salts.²¹

Given the demonstrated ability of well-defined, bidentate phosphine-bound Ni^{I} complexes to re-enter $\text{Ni}^0/\text{Ni}^{\text{II}}$ catalytic cycles,¹⁵ we were interested to determine if monodentate phosphine-bound Ni^{I} complexes could also reenter the cycle and if the % V_{bur} (min) threshold behavior was retained. In order to evaluate the behavior of these monophosphine Ni^{I} species in a catalytic system, $[\text{Ni}^{\text{I}}(\text{COD})\text{Br}]_2$ was employed as a precatalyst in two SMC reactions (Figure 7A). For the three phosphines tested with % V_{bur} (min) values less than 32%, the observed yields are lower when $[\text{Ni}^{\text{I}}(\text{COD})\text{Br}]_2$ is used as a precursor relative to $\text{Ni}(\text{COD})_2$, in line with previous precedent with $\text{dppf}/\text{Ni}^{\text{I}}$ species.

However, moderate product formation did occur for these three ligands, indicating reentry to the $\text{Ni}^0/\text{Ni}^{\text{II}}$ cycle. Notably, a precipitous decline in yield occurs for monophosphines possessing % V_{bur} (min) values >32%, analogous to the reactivity threshold we recently reported with $\text{Ni}(\text{COD})_2$ as a precursor. Even in the absence of COD, **7** was an ineffective precatalyst, giving trace yields in both reactions.

These data suggest that bulky monophosphine-bound Ni^{I} complexes are thermodynamic sinks that are recalcitrant towards $\text{Ni}^0/\text{Ni}^{\text{II}}$ catalytic cycle (re)entry. Specifically, the ability to coordinate two or more phosphine ligands to substrate bound Ni^0 or Ni^{II} complexes appears to be necessary for stability of these even oxidation states in catalysis. The unique geometric and electronic structure of Ni^{I} monomer

and dimer species appears to be ideal for supporting bulky ligands, whereas typical coordination spheres of Ni⁰ and/or Ni^{II} complexes are not amenable to coordination of multiple bulky phosphines (Figure 7B).




Figure 7. (A) Catalytic reactivity studies comparing **5** and Ni(COD)₂ as precatalysts in an SMC reaction. (B) Rationalization of catalytic reactivity by phosphine steric environment across oxidation states.

Synthesis of Ni^I Complexes with Bipyridine Ligands.

Motivated by the catalytic importance of *t*-Bu₃bpy, we sought to access (*t*-Bu₃bpy)Ni^I complexes using the synthetic methods discussed thus far. Tribromide oxidation of a Ni(COD)₂/*t*-Bu₃bpy mixture did not afford isolable Ni^I species; bidentate, π-accepting COD induced the formation of COD-bound Ni⁰ and L_nNiBr₂ species upon concentration (Table 1).²² Likewise, reaction of *t*-Bu₃bpy with the well-defined precursor **5** did not afford isolable [(*t*-Bu₃bpy)Ni^IBr]₂. With these results in hand, we envisioned that reaction of *t*-Bu₃bpy with precursors that contain supporting ligands other than COD would enable access to Ni^I complexes of catalytic interest.

First, we investigated bulky monophosphine-bound Ni^I dimers as synthetic precursors of (*t*-Bu₃bpy)Ni^I complexes. Matsubara and coworkers have previously demonstrated that unsubstituted 2,2'-bipyridine (bpy) could displace the bridging μ-Cl/Br interactions of NHC-bound Ni^I halide dimers to give monomeric (NHC)(bpy)Ni^IX complexes.¹² We evaluated an analogous route from **7** and were able to isolate (*t*-Bu₃bpy)(P(*t*-Bu)₃)Ni^IBr (**17**) by facile ligand displacement. By varying bipyridine ligand identity, it was also

possible to identify (^{CO₂Et}bpy)(P(*t*-Bu)₃)Ni^IBr (see SI, ^{CO₂Et}bpy = diethyl 2,2'-bipyridine-4,4'-dicarboxylate). Indeed, the affinity of large phosphines for the steric environment of Ni^I species (*vide supra*) seems to favor the formation of monomeric species in heteroleptic complexes with bipyridine ligands. It is exciting to recognize the synergy of this relationship: monomeric bipyridine-ligated Ni^I systems are coveted. Furthermore, this class of compounds may suggest previously unexplored mechanistic possibilities for methodologies in which both (poly)pyridyl and bulky monophosphine ligands are employed in one pot.^{54,55}

However, we foresaw potential limitations of a (*t*-Bu₃bpy)Ni^I complex bearing a strongly σ-donating phosphine ligand. Such a mixed-ligand system may lead to ambiguity about which ancillary ligand is responsible for reactivity, among other shortcomings. A more general monomeric (*t*-Bu₃bpy)(L)Ni^IX precursor would fill its coordination sphere with a highly labile ligand, such as an olefin. To this end, we attempted a tribromide oxidation from the recently reported Ni⁰ precursor, Ni⁰(stb)₃ (stb = (*E*)-stilbene).⁵⁶ Upon treatment of a Ni⁰(stb)₃/*t*-Bu₃bpy mixture with 0.5 equiv TBATB, we observed the formation of a previously undetected paramagnetic species by ¹H NMR. Layering the resulting THF solution with pentane and cooling to -35 °C afforded red-black crystals suitable for SCXRD, which identified the complex (*t*-Bu₃bpy)Ni^I(stb)Br (**18**).

Solid-state structural analysis of **17** and **18** reveals subtle differences in the coordination sphere of the metal center (Figure 8D). The *t*-Bu₃bpy ligand of **17** is positioned closer to Ni than in **18**, while **18** features a more tightly bound bromo ligand. The Ni–P(*t*-Bu)₃ bond of **17** is longer than for bpy-free congeners **4** and **10**, consistent with greater steric hindrance around the metal center. Bond metrics on the *t*-Bu₃bpy ligand are within error for the two complexes, suggesting a similar extent of donation from the Ni^I center. The C–C bonds in the backbones of the *t*-Bu₃bpy ligands for **17** and **18** are significantly longer than in (*t*-Bu₃bpy)₂Ni⁰ (C–C_{bpy} = 1.439(6) Å, see SI). This observation is consistent with a relatively π-basic metal center endowing greater *t*-Bu₃bpy-character for Ni⁰ species. Also in accordance with π-basicity, **18** is observed to activate the olefin of stilbene to a lesser extent than its Ni⁰ congener, (*t*-Bu₃bpy)Ni⁰(stb) (**19**).

The electronic structure of these complexes was studied with EPR spectroscopy (Figure 8E). Continuous-wave spectra of **17** and **18** in glassy frozen solutions afforded rhombic signals. The spectrum of a frozen toluene solution containing **17** features no resolved hyperfine splitting and an observed value of *g*_{avg} = 2.261, which is similar to previous reports of well-defined monomeric (bpy/phen)Ni^I(halide) complexes (*g*_{avg} = 2.19–2.24).^{19,22,57,58} For **18**, a color change from dark green to red was observed upon freezing in both 2-MeTHF and toluene, and spectra exhibited abnormal line-shape (see SI). Thorough investigation of a chemical process that may occur upon freezing has not been conducted, as spin relaxation for **18** is sufficiently long to observe a room-temperature spectrum. The observed *g*_{iso} value is 2.211, which is also consistent with prior reports.^{19,22,57,58} Overall, both EPR spectra are consistent with monomeric d⁹ Ni^I complexes as observed by SCXRD.




Figure 8. (A) Precedented dimeric (*t*-Bu₃bpy)Ni^I halide complex and reactivity. (B) Monomeric (*t*-Bu₃bpy)(L)Ni^IBr complexes accessed in this work. (C) Synthesis of complex **18** via complex **19**. Conditions (a): 1.35 equiv (*E, E, E*)-1,5,9-cyclododecatriene, 2.3 equiv Al(OEt)₂Et₂, Et₂O, -35 °C to rt, 16 h. Conditions (b): 1 equiv stb, 1 equiv *t*-Bu₃bpy, Et₂O. (D) Solid-state structures of **17** and **18**, with thermal ellipsoids displayed at 30% probability and hydrogen atoms omitted for clarity. Selected solid-state bond distances (Å) for **17**: Ni1–P1: 2.249(1); Ni1–Br1: 2.4438(7); Ni1–N1: 1.983(3); N1–C1: 1.365(3); C1–C2: 1.472(3). Selected solid-state bond distances (Å) for **18**: Ni2–Br2: 2.4062(6); Ni2–N3: 1.997(3); N3–C3: 1.351(4); C3–C4: 1.471(4); Ni2–C5: 2.057(5); C5–C6: 1.406(5). (E) X-band EPR spectra of **17** (toluene glass, 77 K) and **18** (THF, 298 K). Simulation parameters (**17**): $g_1 = 2.557, g_1 = 2.148, g_3 = 2.077$. Simulation parameters (**18**): $g_{iso} = 2.211$. Synthesis of **17** from **7** not shown. See SI for more information.

Complexes **17** and **18** are the first structurally characterized monomeric *t*-Bu₃bpy-bound Ni^I complexes. The nuclearity of similar complexes is observed to be crucial to their reactivity: dimeric [(*t*-Bu₃bpy)Ni^IX]₂ species (X = Cl, Br) cannot kinetically disaggregate and therefore are inert towards aryl halides (Figure 8A),¹⁸ whereas *in situ* generated monomeric (*t*-Bu₃bpy)Ni^IX species have been shown to activate even aryl chlorides.³⁶ Indeed, **17** and **18** are capable of activating aryl halide bonds (Figure 8B). Furthermore, complex **18** is an especially useful Ni^I model system for mechanistic and stoichiometric studies: it has a direct Ni⁰ analog, **19**, against which its reactivity can be evaluated. A three-step, two-pot synthesis from Ni(acac)₂—which proceeds via **19**—ultimately provided **18** cleanly (Figure 8C). Efforts to prepare **8** through an analogous pathway were unsuccessful, likely due to the instability of a putatively trigonal planar (*t*-Bu₃bpy)(P(*t*-Bu)₃)Ni⁰ complex.

Interested in evaluating the reactivity of these monomeric (*t*-Bu₃bpy)(L)Ni^IBr complexes, we undertook stoichiometric oxidative addition experiments. Both **17** and **18** are observed to react with 1,4-bromofluorobenzene, affording a 1:1 mixture of (*t*-Bu₃bpy)Ni^{III}Br₂ and (*t*-Bu₃bpy)Ni^{III}(4-fluorophenyl)Br (Figure 9). An analogous product mixture has been observed in other stoichiometric oxidative studies with (bpy/phen)Ni^I complexes; it is proposed to arise from rapid comproportionation of a putative Ni^{III} oxidative adduct with remaining Ni^I.^{22,55}

Complexes **17** and **18** are also observed to activate the C(sp²)-Cl bond of 1,4-chlorofluorobenzene (Figure 9). While the oxidative addition of aryl chlorides to (bpy/phen)Ni^I

species has been invoked in methodologies,^{59,60} it has not previously been demonstrated for a well-defined, isolable (bpy/phen)Ni^I complex.^{22,34,55} Complex **18** is observed to convert the anticipated amount of aryl chloride within minutes, while **17** requires several hours of reaction time for the expected 50% conversion. This is perhaps due to the increased lability of olefinic stilbene relative to P(*t*-Bu)₃. Rigorous kinetic and mechanistic investigations into the reactivity profile of these complexes is beyond the scope of this study, but work is underway in our laboratory to further interrogate the behavior of this unique class of compounds.

Stoichiometric oxidative addition studies with 17 and 18




Figure 9. Stoichiometric reactivity with aryl halides for **17** and **18**. ^aConversion measured after 4 h.

CONCLUSIONS

In summary, we have identified a mild oxidative approach and precursor to a variety of Ni^I complexes bearing catalytically relevant ligands. These strategies enabled access to previously elusive complexes, including L₁Ni monophosphine dimers and the first examples of monomeric *t*-Bu₃bpy-bound Ni^I species. For the monophosphine ligand

class, we have elucidated structure–speciation relationships at Ni^I and conducted catalytic studies to relate established steric relationships to observed trends for Ni^I complexes. For the bipyridine ligand class, we have synthesized and characterized well-defined monomeric complexes that are capable of activating aryl bromides and chlorides. We anticipate that our findings will enable future mechanistic studies and catalyst design for Ni-catalyzed cross-coupling reactions.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures, experimental data, and characterization and spectral data for new compounds (PDF)

Accession Codes

CCDC 2204445–2204452, 2204454–2204455, 2204457, 2204459, 2245704–2245705, 2245707–2245709, 2245759, and 2262986–2262993 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Abigail G. Doyle – Department of Chemistry and Biochemistry, University of California Los Angeles, Los Angeles, California 90095, United States; orcid.org/0000-0002-6641-0833
Email: agdoyle@chem.ucla.edu

Authors

Samuel H. Newman-Stonebraker[†] – Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States; Department of Chemistry and Biochemistry, University of California Los Angeles, Los Angeles, California 90095, United States; orcid.org/0000-0001-6611-8480

T. Judah Raab – Department of Chemistry and Biochemistry, University of California Los Angeles, Los Angeles, California 90095, United States; orcid.org/0000-0001-9449-5574

Hootan Roshandel – Department of Chemistry and Biochemistry, University of California Los Angeles, Los Angeles, California 90095, United States; orcid.org/0000-0002-7744-0958

Present Addresses

[†]Department of Chemistry, Yale University, New Haven, CT 06511

Author Contributions

[‡]S. H. N.-S. and T. J. R. contributed equally and are listed in alphabetical order.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We thank Dr. Robert Taylor (UCLA) and Dr. Paul Oyala (Caltech) for assistance with EPR measurements and Dr. Saeed Khan (UCLA) for assistance with X-ray diffraction data collection. We thank Prof. Franziska Schoenebeck for helpful discussions. Financial support for this work was provided by the NIGMS (R35 GM126986). These studies were supported by shared instrumentation grants from the National Science Foundation under equipment grants CHE-1048804 and 2117480, along with the NIH Office of Research Infrastructure Programs under grant S100D028644.

REFERENCES

- (1) Tasker, S. Z.; Standley, E. A.; Jamison, T. F. Recent Advances in Homogeneous Nickel Catalysis. *Nature* **2014**, *509*, 299–309.
- (2) Ananikov, V. P. Nickel: The “Spirited Horse” of Transition Metal Catalysis. *ACS Catal.* **2015**, *5*, 1964–1971.
- (3) Su, B.; Cao, Z.-C.; Shi, Z.-J. Exploration of Earth-Abundant Transition Metals (Fe, Co, and Ni) as Catalysts in Unreactive Chemical Bond Activations. *Acc. Chem. Res.* **2015**, *48*, 886–896.
- (4) Bajo, S.; Laidlaw, G.; Kennedy, A. R.; Sproules, S.; Nelson, D. J. Oxidative Addition of Aryl Electrophiles to a Prototypical Nickel(0) Complex: Mechanism and Structure/Reactivity Relationships. *Organometallics* **2017**, *36*, 1662–1672.
- (5) Diccianni, J. B.; Diao, T. Mechanisms of Nickel-Catalyzed Cross-Coupling Reactions. *Trends Chem.* **2019**, *1*, 830–844.
- (6) Chan, A. Y.; Perry, I. B.; Bissonnette, N. B.; Buksh, B. F.; Edwards, G. A.; Frye, L. I.; Garry, O. L.; Lavagnino, M. N.; Li, B. X.; Liang, Y.; Mao, E.; Millet, A.; Oakley, J. V.; Reed, N. L.; Sakai, H. A.; Seath, C. P.; MacMillan, D. W. C. Metallaphotoredox: The Merger of Photoredox and Transition Metal Catalysis. *Chem. Rev.* **2022**, *122*, 1485–1542.
- (7) Malapit, C. A.; Prater, M. B.; Cabrera-Pardo, J. R.; Li, M.; Pham, T. D.; McFadden, T. P.; Blank, S.; Minter, S. D. Advances on the Merger of Electrochemistry and Transition Metal Catalysis for Organic Synthesis. *Chem. Rev.* **2022**, *122*, 3180–3218.
- (8) Richmond, E.; Moran, J. Recent Advances in Nickel Catalysis Enabled by Stoichiometric Metallic Reducing Agents. *Synthesis* **2017**, *50*, 499–513.
- (9) Beromi, M. M.; Nova, A.; Balcells, D.; Brasacchio, A. M.; Brudvig, G. W.; Guard, L. M.; Hazari, N.; Vinyard, D. J. Mechanistic Study of an Improved Ni Precatalyst for Suzuki–Miyaura Reactions of Aryl Sulfamates: Understanding the Role of Ni(I) Species. *J. Am. Chem. Soc.* **2017**, *139*, 922–936.
- (10) Lin, C.-Y.; Power, P. P. Complexes of Ni(I): A “Rare” Oxidation State of Growing Importance. *Chem. Soc. Rev.* **2017**, *46*, 5347–5399.
- (11) Bismuto, A.; Finkelstein, P.; Müller, P.; Morandi, B. The Journey of Ni(I) Chemistry. *Helv. Chim. Acta* **2021**, *104*, e202100177.
- (12) Matsubara, K.; Fukahori, Y.; Inatomi, T.; Tazaki, S.; Yamada, Y.; Koga, Y.; Kanegawa, S.; Nakamura, T. Monomeric Three-Coordinate N-Heterocyclic Carbene Nickel(I) Complexes: Synthesis, Structures, and Catalytic Applications in Cross-Coupling Reactions. *Organometallics* **2016**, *35*, 3281–3287.
- (13) Guven, S.; Kundu, G.; Weßels, A.; Ward, J. S.; Rissanen, K.; Schoenebeck, F. Selective Synthesis of Z-Silyl Enol Ethers via Ni-Catalyzed Remote Functionalization of Ketones. *J. Am. Chem. Soc.* **2021**, *143*, 8375–8380.
- (14) Beromi, M. M.; Banerjee, G.; Brudvig, G. W.; Charboneau, D. J.; Hazari, N.; Lant, H. M. C.; Mercado, B. Q. Modifications to the Aryl Group of Dppf-Ligated Ni σ-Aryl Precatalysts: Impact on Speciation and Catalytic Activity in Suzuki–Miyaura Coupling Reactions. *Organometallics* **2018**, *37*, 3943–3955.
- (15) Guard, L. M.; Beromi, M. M.; Brudvig, G. W.; Hazari, N.; Vinyard, D. J. Comparison of Dppf-Supported Nickel Precatalysts for the Suzuki–Miyaura Reaction: The Observation and Activity of Nickel(I). *Angew. Chem. Int. Ed.* **2015**, *54*, 13352–13356.

- (16) Beromi, M. M.; Banerjee, G.; Brudvig, G. W.; Hazari, N.; Mercado, B. Q. Nickel(I) Aryl Species: Synthesis, Properties, and Catalytic Activity. *ACS Catal.* **2018**, *8*, 2526–2533.
- (17) Kalvet, I.; Guo, Q.; Tizzard, G. J.; Schoenebeck, F. When Weaker Can Be Tougher: The Role of Oxidation State (I) in P- vs N-Ligand-Derived Ni-Catalyzed Trifluoromethylthiolation of Aryl Halides. *ACS Catal.* **2017**, *7*, 2126–2132.
- (18) Beromi, M. M.; Brudvig, G. W.; Hazari, N.; Lant, H. M. C.; Mercado, B. Q. Synthesis and Reactivity of Paramagnetic Nickel Polypyridyl Complexes Relevant to C(sp²)-C(sp³) Coupling Reactions. *Angew. Chem. Int. Ed.* **2019**, *58*, 6094–6098.
- (19) Somerville, R. J.; Odena, C.; Obst, M. F.; Hazari, N.; Hopmann, K. H.; Martin, R. Ni(I)-Alkyl Complexes Bearing Phenanthroline Ligands: Experimental Evidence for CO₂ Insertion at Ni(I) Centers. *J. Am. Chem. Soc.* **2020**, *142*, 10936–10941.
- (20) Day, C. S.; Rentería-Gómez, Á.; Ton, S. J.; Gogoi, A. R.; Gutierrez, O.; Martin, R. Elucidating Electron-Transfer Events in Polypyridine Nickel Complexes for Reductive Coupling Reactions. *Nat. Catal.* **2023**, *6*, 244–253.
- (21) Standley, E. A.; Smith, S. J.; Müller, P.; Jamison, T. F. A Broadly Applicable Strategy for Entry into Homogeneous Nickel(0) Catalysts from Air-Stable Nickel(II) Complexes. *Organometallics* **2014**, *33*, 2012–2018.
- (22) Ting, S. I.; Williams, W. L.; Doyle, A. G. Oxidative Addition of Aryl Halides to a Ni(I)-Bipyridine Complex. *J. Am. Chem. Soc.* **2022**, *144*, 5575–5582.
- (23) Tsou, T. T.; Kochi, J. K. Mechanism of Oxidative Addition. Reaction of Nickel(0) Complexes with Aromatic Halides. *J. Am. Chem. Soc.* **1979**, *101*, 6319–6322.
- (24) Beck, R.; Shoshani, M.; Krasinkiewicz, J.; Hatnean, J. A.; Johnson, S. A. Synthesis and Chemistry of Bis(Triisopropylphosphine) Nickel(I) and Nickel(0) Precursors. *Dalton Trans.* **2012**, *42*, 1461–1475.
- (25) Newman-Stonebraker, S. H.; Wang, J. Y.; Jeffrey, P. D.; Doyle, A. G. Structure-Reactivity Relationships of Buchwald-Type Phosphines in Nickel-Catalyzed Cross-Couplings. *J. Am. Chem. Soc.* **2022**, *144*, 19635–19648.
- (26) We are aware of a concurrent study under review from the Schoenebeck group that also studies the formation of such complexes.
- (27) Vilar, R.; Mingos, D. M. P.; Cardin, C. J. Synthesis and Structural Characterisation of [Pd₂(μ-Br)₂(PBUt₃)₂], an Example of a Palladium(I)-Palladium(I) Dimer. *J. Chem. Soc., Dalton Trans.* **1996**, 4313–4314.
- (28) Fricke, C.; Sperger, T.; Mendel, M.; Schoenebeck, F. Catalysis with Palladium(I) Dimers. *Angew. Chem. Int. Ed.* **2021**, *60*, 3355–3366.
- (29) Hueffel, J. A.; Sperger, T.; Funes-Ardoiz, I.; Ward, J. S.; Risananen, K.; Schoenebeck, F. Accelerated Dinuclear Palladium Catalyst Identification through Unsupervised Machine Learning. *Science* **2021**, *374*, 1134–1140.
- (30) Zuo, Z.; Ahneman, D. T.; Chu, L.; Terrett, J. A.; Doyle, A. G.; MacMillan, D. W. C. Merging Photoredox with Nickel Catalysis: Coupling of α-Carboxyl Sp³-Carbons with Aryl Halides. *Science* **2014**, *345*, 437–440.
- (31) Huihui, K. M. M.; Caputo, J. A.; Melchor, Z.; Olivares, A. M.; Spiewak, A. M.; Johnson, K. A.; DiBenedetto, T. A.; Kim, S.; Ackerman, L. K. G.; Weix, D. J. Decarboxylative Cross-Electrophile Coupling of N-Hydroxyphthalimide Esters with Aryl Iodides. *J. Am. Chem. Soc.* **2016**, *138*, 5016–5019.
- (32) Zultanski, S. L.; Fu, G. C. Nickel-Catalyzed Carbon-Carbon Bond-Forming Reactions of Unactivated Tertiary Alkyl Halides: Suzuki Arylations. *J. Am. Chem. Soc.* **2013**, *135*, 624–627.
- (33) Kawamata, Y.; Vantourout, J. C.; Hickey, D. P.; Bai, P.; Chen, L.; Hou, Q.; Qiao, W.; Barman, K.; Edwards, M. A.; Garrido-Castro, A. F.; deGruyter, J. N.; Nakamura, H.; Knouse, K.; Qin, C.; Clay, K. J.; Bao, D.; Li, C.; Starr, J. T.; Garcia-Irizarry, C.; Sach, N.; White, H. S.; Neurock, M.; Minter, S. D.; Baran, P. S. Electrochemically Driven, Ni-Catalyzed Aryl Amination: Scope, Mechanism, and Applications. *J. Am. Chem. Soc.* **2019**, *141*, 6392–6402.
- (34) Sun, R.; Qin, Y.; Ruccolo, S.; Schnedermann, C.; Costentin, C.; Nocera, D. G. Elucidation of a Redox-Mediated Reaction Cycle for Nickel-Catalyzed Cross Coupling. *J. Am. Chem. Soc.* **2019**, *141*, 89–93.
- (35) Till, N. A.; Oh, S.; MacMillan, D. W. C.; Bird, M. J. The Application of Pulse Radiolysis to the Study of Ni(I) Intermediates in Ni-Catalyzed Cross-Coupling Reactions. *J. Am. Chem. Soc.* **2021**, *143*, 9332–9337.
- (36) Cagan, D. A.; Bím, D.; McNicholas, B. J.; Kazmierczak, N. P.; Oyala, P. H.; Hadt, R. G. Photogenerated Ni(I)-Bipyridine Halide Complexes: Structure-Function Relationships for Competitive C(sp²)-Cl Oxidative Addition and Dimerization Reactivity Pathways. *Inorg. Chem.* **2023**, ASAP. DOI: 10.1021/acs.inorgchem.3c00917
- (37) Bismuto, A.; Müller, P.; Finkelstein, P.; Trapp, N.; Jeschke, G.; Morandi, B. One to Find Them All: A General Route to Ni(I)-Phenolate Species. *J. Am. Chem. Soc.* **2021**, *143*, 10642–10648.
- (38) Ni, S.; Yan, J.; Tewari, S.; Reijerse, E. J.; Ritter, T.; Cornella, J. Nickel Meets Aryl Thianthrenium Salts: Ni(I)-Catalyzed Halogenation of Arenes. *J. Am. Chem. Soc.* **2023**, *145*, 9988–9993.
- (39) Martinez, G. E.; Ocampo, C.; Park, Y. J.; Fout, A. R. Accessing Pincer Bis(Carbene) Ni(IV) Complexes from Ni(II) via Halogen and Halogen Surrogates. *J. Am. Chem. Soc.* **2016**, *138*, 4290–4293.
- (40) Zhou, Y.-Y.; Hartline, D. R.; Steiman, T. J.; Fanwick, P. E.; Uyeda, C. Dinuclear Nickel Complexes in Five States of Oxidation Using a Redox-Active Ligand. *Inorg. Chem.* **2014**, *53*, 11770–11777.
- (41) Evans, D. F. 400. The Determination of the Paramagnetic Susceptibility of Substances in Solution by Nuclear Magnetic Resonance. *J. Chem. Soc.* **1959**, 2003–2005.
- (42) Dicciani, J. B.; Katigbak, J.; Hu, C.; Diao, T. Mechanistic Characterization of (Xantphos)Ni(I)-Mediated Alkyl Bromide Activation: Oxidative Addition, Electron Transfer, or Halogen-Atom Abstraction. *J. Am. Chem. Soc.* **2019**, *141*, 1788–1796.
- (43) Ciszewski, J. T.; Mikhaylov, D. Y.; Holin, K. V.; Kadirov, M. K.; Budnikova, Y. H.; Sinyashin, O.; Vivic, D. A. Redox Trends in Terpyridine Nickel Complexes. *Inorg. Chem.* **2011**, *50*, 8630–8635.
- (44) Dible, B. R.; Sigman, M. S.; Arif, A. M. Oxygen-Induced Ligand Dehydrogenation of a Planar Bis-μ-Chloronickel(I) Dimer Featuring an NHC Ligand. *Inorg. Chem.* **2005**, *44*, 3774–3776.
- (45) [Ni(COD)Br]₂ could also be synthesized using TPhATB, but we found that use of TBATB as the oxidant led to less decomposition and higher yields.
- (46) Porri, L.; Vitulli, G.; Gallazzi, M. C. 1,5-Cyclooctadienenickel Bromide and Iodide. *Angew. Chem. Int. Ed.* **1967**, *6*, 452.
- (47) Lohrey, T. D.; Cusumano, A. Q.; Goddard, W. A.; Stoltz, B. M. Identifying the Imperative Role of Metal-Olefin Interactions in Catalytic C-O Reductive Elimination from Nickel(II). *ACS Catal.* **2021**, *11*, 10208–10222.
- (48) Newman-Stonebraker, S. H.; Smith, S. R.; Borowski, J. E.; Peters, E.; Gensch, T.; Johnson, H. C.; Sigman, M. S.; Doyle, A. G. Univariate Classification of Phosphine Ligation State and Reactivity in Cross-Coupling Catalysis. *Science* **2021**, *374*, 301–308.
- (49) Gensch, T.; Gomes, G. dos P.; Friederich, P.; Peters, E.; Gaudin, T.; Pollice, R.; Jorner, K.; Nigam, A.; Lindner-D'Addario, M.; Sigman, M. S.; Aspuru-Guzik, A. A Comprehensive Discovery Platform for Organophosphorus Ligands for Catalysis. *J. Am. Chem. Soc.* **2022**, *144*, 1205–1217.
- (50) Within minutes at ambient temperature in the solution phase, Ni⁰ black and a color change from green to dark blue was observed for L₁Ni^I dimers with bulky phosphines (L = P(*t*-Bu)₃, CyP(*t*-Bu)₂). Per SCXRD analysis, the dark blue solution likely contains Ni^{1.5+} dimer, as in 3.
- (51) For ligands smaller than P(*t*-Bu)₃ where more than one Ni^I species can form, L:Ni stoichiometry controlled the outcome for isolated material in the solid state. Speciation was generally conserved in solution-state NMR characterization (C₆D₆), though a

small amount of phosphine dissociation and dimerization was observed for L_2Ni^I monomer **10** with $CyP(t-Bu)_2$.

(52) Hoberg, H.; Radine, K.; Krüger, C.; Romão, M. J. Synthesis of New Phosphine Nickel(I) Complexes and Crystal Structure of μ_3 -Iodo-Tris- μ -Iodo-Cyclotris(Triphenylphosphine Nickel), $(TPP)_3Ni_3I_4$. *Z. Naturforsch. B* **1985**, *40*, 607–614.

(53) Borowski, J. E.; Newman-Stonebraker, S. H.; Doyle, A. G. Comparison of Monophosphine and Bisphosphine Precatalysts for Ni-Catalyzed Suzuki–Miyaura Cross-Coupling: Understanding the Role of the Ligation State in Catalysis. *ACS Catal.* **2023**, *13*, 7966–7977.

(54) Sheng, J.; Ni, H.-Q.; Liu, G.; Li, Y.; Wang, X.-S. Combinatorial Nickel-Catalyzed Monofluoroalkylation of Aryl Boronic Acids with Unactivated Fluoroalkyl Iodides. *Org. Lett.* **2017**, *19*, 4480–4483.

(55) Hamby, T. B.; LaLama, M. J.; Sevov, C. S. Controlling Ni Redox States by Dynamic Ligand Exchange for Electroreductive Csp³–Csp² Coupling. *Science* **2022**, *376*, 410–416.

(56) Nattmann, L.; Saeb, R.; Nöthling, N.; Cornella, J. An Air-Stable Binary Ni(0)–Olefin Catalyst. *Nat. Catal.* **2020**, *3*, 6–13.

(57) Lin, Q.; Diao, T. Mechanism of Ni-Catalyzed Reductive 1,2-Dicarbonylfunctionalization of Alkenes. *J. Am. Chem. Soc.* **2019**, *141*, 17937–17948.

(58) Humphrey, E. L. B. J.; Kennedy, A. R.; Sproules, S.; Nelson, D. J. Evaluating a Dispersion of Sodium in Sodium Chloride for the Synthesis of Low-Valent Nickel Complexes. *Eur. J. Inorg. Chem.* **2022**, e202101006.

(59) Mirabi, B.; Marchese, A. D.; Lautens, M. Nickel-Catalyzed Reductive Cross-Coupling of Heteroaryl Chlorides and Aryl Chlorides. *ACS Catal.* **2021**, *11*, 12785–12793.

(60) Yue, H.; Zhu, C.; Rueping, M. Cross-Coupling of Sodium Sulfinates with Aryl, Heteroaryl, and Vinyl Halides by Nickel/Photoredox Dual Catalysis. *Angew. Chem. Int. Ed.* **2018**, *57*, 1371–1375.

