

Strain induced reactivity of cyclic iminoboranes: The (2 + 2) cycloaddition of a 1-*H*-1,3,2-diazaborepine with ethene

Divanshu Gupta, Ralf Einholz, Holger F. Bettinger*

Institut für Organische Chemie, Universität Tübingen, Auf der Morgenstelle 18, 72076 Tübingen, Germany

E-mail: holger.bettinger@uni-tuebingen.de

Abstract


Iminoboranes have gathered immense attention due to their reactivity and potential applications as isoelectronic and isosteric alkynes. While cyclic alkynes are well investigated and useful reagents, cyclic iminoboranes are underexplored and their existence was inferred only via trapping experiments. We report the first direct spectroscopic evidence of a cyclic seven-membered iminoborane, 1-(*tert*-butyldimethylsilyl)-1-*H*-1,3,2-diazaborepine **2**, under cryogenic matrix isolation conditions. The amino-iminoborane **2** was photochemically generated in solid argon at 4 K from 1-(*tert*-butyldimethylsilyl)-2-azido-1,2-dihydro-1,2-azaborine (**3**) and was

characterized using FT-IR, UV-Vis spectroscopy, and computational chemistry. The characteristic BN stretching vibration (1751 cm^{-1}) is shifted by about 240 cm^{-1} compared to linear amino-iminoboranes indicating a significant weakening of the bond. The Lewis acidity value determined computationally ($LA_B = 9.1 \pm 2.6$) is similar to that of boron trichloride, and twelve orders of magnitude lower than that of 1,2-azaborinine (BN-aryne, $LA_B = 21.5 \pm 2.6$), a six-membered cyclic iminoborane. In contrast to the latter, the reduced ring strain of **2** precludes nitrogen fixation, but it unexpectedly allows facile (2 + 2) cycloaddition reaction with C_2H_4 under matrix isolation conditions at 30 K. The reaction of **2** with ethene provides the first example of a (2 + 2) cycloaddition reaction of an iminoborane.

Introduction

Iminoboranes are an important class of BN containing compounds.¹⁻³ The BN/CC isosterism relates iminoboranes and alkynes (Scheme 1a).⁴ In contrast to the latter, iminoboranes are kinetically unstable towards cyclooligomerization or polymerization and thus special conditions or steric protection are required for their synthesis and isolation.^{1-2, 5-12} The first iminoborane stable at room temperature, $F_5C_6\text{-BN-tBu}$, was reported by Paetzold et al.,¹¹ and followed by numerous examples.^{2, 9, 13-15} Since then, iminoboranes have attracted tremendous attention.¹⁶⁻²³ Various iminoboranes have been investigated over the past decades to understand their reactivity²⁴⁻²⁷ including the formation of BN-doped polycyclic aromatic hydrocarbons (PAHs),²⁸⁻³² formation of N-heterocyclic carbene coordinated iminoboranes,^{25, 33-36} synthesis of BN containing heterocycles,³⁷⁻⁴⁴ (2 + 2) and (2 + 3) cycloaddition reactions with a number of polar double bonds (e. g., $RR'C=O$) and


dipolar reagents,⁴⁵⁻⁵⁰ formation of iminoboryl carboranes,^{19, 51} and use of frustrated Lewis pairs to stabilize iminoboranes.^{19, 52-57} A series of ab initio computational studies performed by Gilbert compared iminoboranes and alkynes with respect to the electronic and geometric structure, as well as the reactivity in (2 + 2) and (2 + 4) cycloadditions towards alkenes and alkynes.⁵⁸⁻⁶⁰



Scheme 1. a) Iminoboranes and their analogy with alkynes. b) Resonance forms of 1,2-azaborinine (**1**) and its analogy with *ortho*-benzyne. c) Representative examples of cyclic iminoboranes previously inferred by Paetzold et al.⁶¹ and Braunschweig et al.⁴²

Studies on cyclic iminoboranes are quite rare.^{42, 61} Due to ring strain, they are more reactive than linear iminoboranes,¹ but coordination by N-heterocyclic carbenes (NHC) provides a way of stabilization.^{36, 62-63} A special case of cyclic iminoboranes is the aromatic BN-aryne, 1,2-azaborinine **1**, the BN analogue of *ortho*-benzyne (Scheme 1b).⁶⁴⁻⁶⁶ This was detected by matrix isolation methods by our group and shows remarkably high reactivity towards inert molecules.⁶⁴⁻⁶⁶ The polarity of the strained BN link of **1** results in a bonding situation that differs from that of the strained triple bond in arynes and from

that of the BN unit in linear iminoboranes.^{64, 66-67} The dibenzo derivative of **1**, dibenzo[*c,e*][1,2]azaborinine, was inferred as reactive intermediate in solution,⁶⁸⁻⁷⁰ and can even activate the strong Si–F bond for subsequent insertion reaction.⁷⁰



Scheme 2. Photogeneration of 1-(*tert*-butyldimethylsilyl)-1,3,2-diazaborepine **2** under matrix isolation conditions and its reaction with CO and C₂H₄.

As iminoborane units in larger than six-membered rings have never been observed directly, but only inferred from trapping experiments (Scheme 1c),^{42, 61} we studied the seven membered 1-(*tert*-butyldimethylsilyl)-1-*H*-1,3,2-diazaborepine **2** to elucidate the impact of ring size on the reactivity of cyclic iminoboranes. The matrix isolation technique is ideally suited to study such highly reactive intermediates directly, and furthermore probe their reactivity towards interesting target compounds. We generated target

compound **2** in solid argon by the photolysis of 2-azido-1-(*tert*-butyldimethylsilyl)-1,2-dihydro-1,2-azaborinine (**3**) under cryogenic matrix conditions (see Scheme 2). On one hand, ring strain will be reduced in this seven-ring BN iminoborane compared to **1** and this is expected to reduce its reactivity, while on the other hand, the eight π -electron count may induce some degree of antiaromaticity that will potentially destabilize **2**. Seven-membered cyclic iminoboranes were never directly observed, but recently inferred as reactive intermediate in the formation of diazadiboretidines by Braunschweig.⁴² We provide here for the first time direct spectroscopic evidence for a cyclic seven-ring iminoborane and reveal its unexpectedly facile (2 + 2) cycloaddition reaction (Scheme 1c).

Results and Discussion

Generation and characterization of cyclic iminoborane 2. Matrix-isolation infrared spectroscopic studies of 2-azido-1-(*tert*-butyldimethylsilyl)-1,2-dihydro-1,2-azaborinine, **3**, were performed in solid Ar at 4 K. The most distinguished peak at 2141 cm^{-1} corresponds to the $\nu(\text{N}_3)$ stretching vibration (Figure 1 and Figure S10, Table S12 in Supporting Information). Irradiation of matrix isolated precursor **3** with $\lambda = 254 \text{ nm}$ until it was completely bleached, as shown in the difference spectrum of Figure 1d, resulted in a new set of IR bands. This set is assigned to **2** based on comparison with computational data obtained at the B3LYP-D3(BJ)/6-311+G(d,p) level of theory. The formation of the structural isomer 1-*H*-1,2,3-diazaborepine or the triplet boryl nitrene, the expected product of N_2 extrusion from **3**, can be excluded based on the computations (see Figure S16). Among the new bands formed during photolysis of **3**, the most intense peak, 1751 cm^{-1}

corresponds to the $\nu(\text{BN})$ stretching vibration of **2** (Table S14) by comparison with the computations (Figure 1e). The band at 1809 cm^{-1} corresponds to the $\nu(\text{BN})$ stretching vibration of the ^{10}B isotopologue, and the isotopic shift of 58 cm^{-1} is in good agreement with computations (60 cm^{-1}). The formation of the 1-*H*-1,3,2-diazaborepine isomer is in agreement with the known behavior of azidoboranes: thermolysis or photolysis of diorganyl azidoboranes results in iminoboranes without trappable borylnitrenes,¹ while only in the case of donor substitution ($\text{X} = \text{NR}_2, \text{OR}$) the borylnitrenes X_2BN could be trapped or directly observed.⁷¹⁻⁷⁷ The photolysis of azide **3** is thus expected to result in ring enlargement by shift of the carbon rather than the nitrogen center to give the cyclic iminoborane **2**.




Figure 1. a) IR spectrum obtained after deposition of **3** in Ar matrix at 28 K. b) IR spectrum obtained after irradiation of Ar matrix with $\lambda = 254\text{ nm}$ (following the deposition of **3**). c) Spectrum for ^{11}B and ^{10}B isotopologues (81:19) of **3** calculated at the B3LYP-

D3(BJ)/6–311+G(d,p) level of theory. d) Difference spectrum obtained after irradiation of Ar matrix with $\lambda = 254$ nm (following the deposition of **3**). e) Spectrum for ^{11}B and ^{10}B isotopologues (81:19) of **2** calculated at the B3LYP-D3(BJ)/6–311+G(d,p) level of theory. (● corresponds to the overtones and combination bands according to computed anharmonic vibrational frequency analysis).

For linear amino-iminoboranes, the BN stretching mode was reported around 1990 cm^{-1} and 2030 cm^{-1} for the ^{11}B and ^{10}B isotopologues, respectively.⁷⁸⁻⁷⁹ The large shift of the BN stretching frequency of **2** from linear diorganyl substituted amino-iminoboranes of around 240 cm^{-1} is due to the cyclic nature of **2** which results in considerable weakening of the BN bond. Note that the BN stretching for **1**⁶⁴ at 1637/1634 cm^{-1} indicates an even weaker BN bond in the six-membered ring.

UV/vis spectroscopy (Figure S8) of the azide **3** in Ar at 8 K shows a strong feature around 290 nm with fine structure and maxima at 296 nm, 288 nm, and 244 nm that is typical for 1,2-azaborines.⁸⁰⁻⁸¹ Irradiation with $\lambda = 254$ nm results in quick decrease of the azide absorption band with an isosbestic point at 313 nm. The photoproduct **2** only has a relatively weaker and broad absorption maximum centered at 305 nm. The measured spectra of **3** and **2** are in good agreement with TD-CAM-B3LYP/6-311+G(d,p) computations (Figure S9).

The most remarkable feature of the computed geometry (M06-2X/6-311+G(d,p)) of **2** is the distortion of the heptagon with a small angle of 104.9° at nitrogen and a large angle of 161.4° at boron in the singlet ground state (Figure 2a). Similarly to the case of

1,2-azaborinine, this feature can be rationalized by the difference in electronegativity: the more electronegative N atom prefers to have a σ -type lone pair (HOMO-1), while the more electropositive B atom prefers an empty σ -type orbital (LUMO) which has a strong p character (Figure 2b). The natural bond orbital (NBO) analysis at the M06-2X/6-311+G(d,p) level of theory arrives at occupancies of 1.83 e⁻ and 0.33 e⁻ for the lone pair and empty orbitals, respectively, slightly increased compared to the NBO occupancies of **1** (1.82 e⁻ and 0.21 e⁻ for the lone pair and empty orbitals). There is a pronounced n(N) → n*(B) interaction [$E(2)$ =34.5 kcal/mol] according to the second-order perturbation estimate of the donor–acceptor interaction in the NBO basis that is smaller than that in **1** (39.7 kcal/mol). The natural charges obtained from the NBO analysis on N and B are large (-0.91 on N and +1.16 on B) while the Wiberg bond index between B and N is only 1.55. Compared to 1,2-azaborinine **1** the Wiberg bond index is larger (1.43 in **1**) and the polarity of the BN bond (natural charges in **1**: -0.81 on N and +1.00 on B; NBO occupancies of 1.82 e⁻ and 0.21 e⁻ for the lone pair and empty orbitals) is increased in **2**.





Figure 2. a) Geometrical parameters of **2**. b) Natural bond orbitals (NBOs) and their occupation numbers of **2**. Important bond lengths [Å] and bond angles [°] are given. All computations at the M06-2X/6-311+G(d,p) level of theory. c) Plot of NICS_{zz} value calculated along a line perpendicular to the molecular plane, starting at the ring center, at B3LYP/6-311+G(d,p)//M06-2X/6-311+G(d,p) level of theory.

We compare the aromatic character of **2** with that of 1,2-azaborinine **1**, benzene, and benzyne using nuclear independent chemical shift (NICS) calculations,⁸²⁻⁸⁴ placing


the dummy atom 1 Å above the center of the ring to get the NICS(1)_{zz} value. 1,2-Azaborinine **1** has a NICS(1)_{zz} value of -25.2 which is lower than that of benzene (-29.3) and benzyne (-33.5), but still suggests aromatic character that is slightly larger than that of 1,2-dihydro-1,2-azaborine (NICS(1)_{zz} = -21.1).⁸⁵ The NICS(1)_{zz} value of -1.1 and -3.1 computed for **2** is significantly lower suggesting that the antiaromaticity expected on the basis of the formal electron count is not relevant (see Figure 2c). The two slightly differing values for **2** are due to the non-planar ring.



Scheme 3. Homodesmotic reaction for the calculation of strain energy.

We compared the strain of the two cyclic iminoboranes (**1** and **2**) by considering homodesmotic reactions (Scheme 3). As expected, the strain energy of the seven-ring iminoborane **2** is lower than that of 1,2-azaborinine by approximately 6.3 kcal/mol at the

M06-2X/6-311+G(d,p) level of theory. We also computed the Lewis acidities (LA_B) of **1** and **2** using the method of Ofial et al. from known Lewis basicities LB_B and Lewis acidities LA_B , computed equilibrium constants (Scheme 4, $\Delta G_{iso}^\circ = -RT \ln K_B$), and the equation $\log K_B = LA_B + LB_B$.⁸⁶ Ofial et al.⁸⁶ obtained the LA_B parameters from ΔG_{iso} to give $LA_B = 8.4 \pm 2.0$ for BF_3 , $LA_B = 9.3 \pm 1.8$ for BCl_3 , and $LA_B = 10.1 \pm 1.3$ for BBr_3 . The qualitative ordering obtained by Ofial et al.⁸⁶ of Lewis acidities LA_B with $BF_3 < BCl_3 < BBr_3$ is in accordance with Lewis acidity rankings based on spectroscopic data.⁸⁷⁻⁸⁹ The individual LA_B parameters obtained from ΔG_{iso} for three different reference bases were averaged to give $LA_B = 21.5 \pm 2.6$ for **1** and $LA_B = 9.1 \pm 2.6$ for **2** (see the Supporting Information for further details). While the Lewis acidity of **2** is thus similar to that of BCl_3 , that of **1** is 12 orders of magnitudes larger.



Scheme 4. Combining the isodesmic reaction [Eq. I] with an experimental reference reaction [Eq. II] allows one to determine the Lewis acidities of X (X= **1** or **2**) from ΔG_{iso} [Eq. III]. The Lewis bases pyridine, acetonitrile, and benzaldehyde and the Lewis acids BAr_3 (Ar = C_6H_5 , 4- ClC_6H_4 , 3,4,5- $F_3C_6H_2$) were chosen as references. The borane Lewis acidity scale is defined so that $LA_B(\text{triphenylborane}) = 0$.

Interaction of 2 with N₂. A hallmark of the superelectrophilic 1,2-azaborinine **1** is its ability to bind nitrogen to give adduct **1•N₂** under cryogenic conditions.⁶⁴ In order to study the Lewis acidity of the photogenerated 1-H-1,3,2-diazaborepine **2** experimentally, the matrix was annealed up to 35 K. This did not lead to any further spectral changes, indicating that **2** cannot bind the photochemically extruded N₂ that is lying in its vicinity. The potential energy paths computed with the B-N₂ distance as parameter (B3LYP-D3(BJ)/6-311+G(d,p)) reveal that formation of the dative complex **2•N₂** is energetically unfavorable and involves a barrier (Figure 3a). In contrast, formation of **1•N₂** is without barrier and exothermic (Figure 3b).




Figure 3. Plot for the relative energy in kcal/mol as a function of the boron nitrogen distance in Å for a) 1-(*tert*-butyldimethylsilyl)-1,3,2-diazaborepine **2** and b) 1,2-azaborinine **1**, respectively, computed at B3LYP-D3(BJ)/6-311+G(d,p) level of theory.

Interaction of 2 with CO. The stronger Lewis base CO can undergo formation of adduct **4** as revealed in experiments using mixtures of Ar and CO (1 – 2 % in Ar) (Scheme 2, Figure 4). In separate experiments, the isotopologues ¹³CO and C¹⁸O (Figures S14-S16) were

employed in order to obtain isotopic shifts of vibrational bands for comparison with computations.




Figure 4. Infrared spectra obtained after irradiation of **3** in CO (2–3 %) doped argon matrix. a) After 60 min irradiation with $\lambda=254$ nm at $T=4$ K. b) Difference spectrum after annealing for 30 min at 30 K (following the irradiation with $\lambda=254$ nm). c) Difference spectrum after irradiation with $435 \text{ nm} > \lambda > 620 \text{ nm}$ for 30 min (following the annealing at 30 K).

Irradiation of matrix-isolated precursor **3** with $\lambda = 254$ nm forms diazaborepine **2** in the presence of CO. Annealing the matrix to 30 K allows the formation of distinct bands which decrease upon subsequent irradiation with $435 \text{ nm} > \lambda > 620 \text{ nm}$, while the bands corresponding to **2** collectively form again (see Figure 4). We assign the bands formed

after annealing of the matrix to 30 K to the Lewis acid-base adduct **4** based on the comparison with the vibrational spectrum computed at B3LYP-D3(BJ)/6-311+G(d,p) level of theory (see Figure 5).




Figure 5. a) Spectrum for ¹¹B and ¹⁰B isotopologues (81:19) of **4** calculated at the B3LYP-D3(BJ)/6-311+G(d,p) level of theory. b) Difference IR spectrum after irradiation with 435 nm > λ > 620 nm for 30 min (following the annealing step). c) Spectrum for ¹¹B and ¹⁰B isotopologues (81:19) of **2** calculated at the B3LYP-D3(BJ)/6-311+G(d,p) level of theory.

The most intense peak at 2112 cm⁻¹ is due to the ν(CO) stretching vibration of the Lewis acid – base adduct **4** (Table S15). It is observable at 2065 cm⁻¹ and 2064 cm⁻¹ for the C¹⁸O and ¹³CO isotopologues, respectively (Table S16, S17). These isotopic shifts of

47 cm⁻¹ (¹²C¹⁸O vs. ¹²C¹⁶O) and 48 cm⁻¹ (¹³C¹⁶O vs. ¹²C¹⁶O) are in very good agreement with those computed for **4** (47 cm⁻¹ and 50 cm⁻¹, respectively, see Table S18).

The trapping of **2** with CO demonstrates that Lewis acid–base interaction is significantly preferred over (2 + 1) or (2 + 2) cycloaddition reactions, similar to the reaction of 1,2-azaborinine with CO.⁶⁶ The formation of **4** has a barrier of 1.3 kcal/mol from the complex **4_comp'** at the DLPNO-CCSD(T)/cc-pVTZ//M06-2X/6-311+G(d,p) level of theory (Figure 6). The formation of the (2+1) product **4_(2 + 1)** from **4** does not proceed thermally as its barrier of 6.4 kcal/mol (Figure 6) is too high for this reaction to be observable at 30 K under matrix isolation conditions. The formation of the (2 + 2) cycloaddition product via complex **4_comp** involves a barrier of 23.1 kcal/mol at the DLPNO-CCSD(T)/cc-pVTZ//M06-2X/6-311+G(d,p) level of theory. This barrier is also too high for the reaction to proceed under our experimental conditions. The trapping of **2** with CO demonstrates that Lewis acid–base interaction is significantly preferred over (2 + 1) or (2 + 2) cycloaddition reactions.




Figure 6. Reaction pathways for the reaction of **2** with CO at (Bold: M06-2X/6-311+G(d,p); Italics: DLPNO-CCSD(T)/cc-pVTZ//M06-2X/6-311+G(d,p)). Calculated ZPVE corrected energies in kcal/mol are shown.

Reaction of 2 with ethene. After the successful generation of **2** and its trapping with CO, we used ethene as the simplest olefin for studying of the reactivity of **2** in (2 + 2) cycloaddition reactions. To examine the trapping of **2** with ethene, matrix-isolated precursor **3** was irradiated in the presence of 5 % C₂H₄ in Ar with $\lambda = 254$ nm, which again resulted in 1,3,2-diazaborepine **2**. But apart from **2**, new signals were also observed (Figure 7) that we assigned with the help of the computed spectrum to **5**, the (2 + 2) cycloaddition product between **2** and C₂H₄ (Scheme 2, Figure 8). Subsequently, the matrix was annealed to 30 K and it was observed that the set of signals corresponding to

2 were diminished (Figure 7c), while the set of signals assigned to **5** increased in intensity (Figure 7c). This shows that the formation of **5** from **2** and ethene can proceed thermally with a very low activation barrier.




Figure 7. Infrared spectra obtained after a) deposition of **3** in C₂H₄ (5 %) doped argon matrix. b) Difference spectrum after 60 min irradiation with $\lambda = 254$ nm at T=4 K. c) Difference spectrum after annealing for 30 min at 30 K (following the irradiation with $\lambda = 254$ nm).

The prominent and characteristic signals of **5**, 1637 cm⁻¹ and 1615 cm⁻¹, are attributed to the $\nu(\text{C}=\text{C})$ stretching vibration of the diazaborepine ring (Table S19, S20). The same $\nu(\text{C}=\text{C})$ stretching vibrations at 1637 cm⁻¹ and 1615 cm⁻¹ were also observed

for the C_2D_4 isotope as the stretching mode does not involve any vibration of the ethene unit (Figure S15). The most intense peak was observed at 828 cm^{-1} , which we assigned to a vibration including ring stretching and $\nu(\text{CH}\text{ wag})$ of C_2H_4 , while a peak at 823 cm^{-1} was observed in case of reaction with C_2D_4 . The experimentally observed isotopic shift of 5 cm^{-1} is in good agreement with the isotopic shift of 4 cm^{-1} computed at the B3LYP-D3(BJ)/6-311+G(d,p) level of theory (see Table S21 in SI).




Figure 8. a) Spectrum for ^{11}B and ^{10}B isotopologues (81:19) of **2** calculated at the B3LYP-D3(BJ)/6-311+G(d,p) level of theory. b) Difference IR spectrum after annealing to 30 K for 60 min (following the irradiation step). c) Spectrum for ^{11}B and ^{10}B

isotopologues (81:19) of **5** calculated at the B3LYP-D3(BJ)/6–311+G(d,p) level of theory.
(▲ corresponds to the peaks of C₂H₄)

According to computations at the DLPNO-CCSD(T)/cc-pVTZ//M06-2X/6-311+G(d,p) level of theory, **2** can interact with ethene to give two complexes, **5_comp'** and **5_comp**, that are lower in energy than the separated reactants by 4.6 and 6.7 kcal/mol, respectively (see Figure 9). The interconversion between complexes **5_comp'** and **5_comp** is associated with a barrier of 1.2 kcal/mol. The formation of the (2 + 2) cycloaddition product **5** from complex **5_comp** can proceed thermally as its barrier of 1.0 kcal/mol (Figure 9) is small enough for this reaction to be observable at 30 K under matrix isolation conditions. Formation of the C-H insertion product **6** via a concerted pathway from complex **5_comp'** involves a barrier of 23.0 kcal/mol. This is significantly higher than the barrier of (2 + 2) addition product formation (Figure 9) and explains the preferred formation of **5** in the experiment.




Figure 9. Pathways for the reaction of **2** with C₂H₄ at M06-2X/6-311+G(d,p) (**bold**) and DLPNO-CCSD(T)/cc-pVTZ//M06-2X/6-311+G(d,p) (*italics*). Calculated ZPVE corrected energies in kcal/mol are shown. Geometrical parameters of complex **5_comp** computed at the M06-2X/6-311+G(d,p) level of theory. Important bond lengths [Å] and bond angles [°] are given.

Complex **5_comp** has carbon-boron distances (1.771 Å and 1.887 Å) that are shorter than the sum of van-der-Waals radii (3.62 Å) (see Figure 9) indicative of a Lewis acid – base interaction.⁹⁰⁻⁹¹ The H-C-H and H-C-C angles have changed so that the sum of bond angles around carbon is 357.1 and 359.6°, showing a slight deviation from planarity around carbon atoms in C₂H₄. The complex **5_comp** is formed due to the

interaction between the empty boron orbital with high p-character and the C=C π -bond. Further insight into the nature of bonding in **5_comp** is provided by natural bond orbital analysis.⁹²⁻⁹⁵ The second-order perturbation theory analysis in the NBO basis gives an E(2) value of 314.8 kcal/mol, implying strong stabilization due to $\pi(\text{C}_2\text{H}_4) \rightarrow n^*(\text{B})$ (empty orbital on boron) delocalization. The corresponding natural localized molecular orbitals (NLMO) have major contributions from the $\pi(\text{C}=\text{C})$ bond orbitals and “delocalization tails” of 20.6 % from a slightly hybridized vacant orbital at boron (Table S10 and S11).

The comparison of the reaction mechanism for the reaction of ethene with 1,2-azaborinine **1** and 1-(*tert*-butyldimethylsilyl)-1,3,2-diazaborepine **2** is instructive. The former interacts with ethene to give a Lewis acid-base complex whose formation is thermodynamically highly exothermic (-23.7 kcal/mol at CCSD(T)/cc-pVTZ//M06-2X/6-311+G(d,p)).⁶⁷ This complex is so low in energy that the formation of the (2 + 2) cycloaddition product has a barrier of 14.6 kcal/mol at CCSD(T)/cc-pVTZ//M06-2X/6-311+G(d,p).⁶⁷ This is significantly higher than the barrier for the (2 + 2) cycloaddition product **5** (1.0 kcal/mol from **5_comp**).⁶⁷ It is highly interesting and unexpected that the much less Lewis acidic **2** is undergoing unprecedented (2 + 2) cycloaddition with ethene while the highly Lewis acidic **1** is expected to be trapped in the Lewis acid-base complex with ethene as this reaction should not be observable at 30 K under matrix isolation conditions. Preliminary experiments of **1** + C₂H₄ indeed indicate that thermally initiated (2 + 2) cycloaddition is not occurring under similar matrix isolation conditions.

Conclusions

We here for the first time reveal the generation, spectroscopic detection by matrix isolation, and reactivity studies of a seven membered cyclic iminoborane of the 1-*H*-1,3,2-diazaborepine type. The precursor 2-azido-1-(*tert*-butyldimethylsilyl)-1,2-dihydro-1,2-azaborinine (**3**) undergoes a photoinduced N₂ extrusion and rearrangement to the 1-(*tert*-butyldimethylsilyl)-1-*H*-1,3,2-diazaborepine **2** without detectable intermediates. Detailed combined experimental and computational investigations reveal that **2** shows high reactivity towards carbon monoxide and ethene even at the very low temperature. With CO, **2** reacts to form Lewis acid-base adduct **4** while the alternative (2 + 1) or (2 + 2) cycloaddition reactions were not observed due to high energy barriers. With C₂H₄, **2** undergoes (2 + 2) cycloaddition reaction to form product **5**. The (2 + 2) cycloaddition with ethene discovered here shows that the **2** has the potential to provide novel modes of reactivity for the construction of BN-containing heterocycles.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This work was supported by the German Research Foundation (DFG) through grant BE 3183/6-2 and in part by the Dr. Eberle foundation. The computations were performed on BwForCluster JUSTUS2 cluster. The authors acknowledge support by the state of Baden-Württemberg through bwHPC and the German Research Foundation (DFG) through grant no INST 40/575-1 FUGG (JUSTUS 2 cluster) for computation facilities.

References

1. Paetzold, P., Iminoboranes. In *Advances in Inorganic Chemistry*, Elsevier: 1987; Vol. 31, pp 123-170.
2. Nöth, H., The Chemistry of Amino Imino Boranes. *Angew. Chem. Int. Ed. Engl.* **1988**, *27* (12), 1603-1623.
3. Fan, Y.; Cui, J.; Kong, L., Recent Advances in the Chemistry of Iminoborane Derivatives. *Eur. J. Org. Chem.* **2022**, *2022* (43), e202201086.
4. Liu, Z.; Marder, T. B., B=N versus C≡C: How Similar Are They? *Angew. Chem. Int. Ed.* **2008**, *47* (2), 242-244.
5. Lory, E. R.; Porter, R. F., Infrared studies of matrix isolated species in the hydrogen-boron-nitrogen system. *J. Am. Chem. Soc.* **1973**, *95* (6), 1766-1770.
6. Thompson, C. A.; Andrews, L., Reactions of B atoms with NH₃ to produce HBNH, BNBH, and B₂N. *J. Am. Chem. Soc.* **1995**, *117* (40), 10125-10126.
7. Zhang, F.; Maksyutenko, P.; Kaiser, R. I.; Mebel, A. M.; Gregušová, A.; Perera, S. A.; Bartlett, R. J., On the Directed Gas Phase Synthesis of the Imidoborane Molecule (HNBH) — An Isoelectronic Molecule of Acetylene (HCCH). *J. Phys. Chem. A* **2010**, *114* (46), 12148-12154.
8. Baird, N. C.; Datta, R. K., Ab initio molecular orbital study of iminoborane. *Inorg. Chem.* **1972**, *11* (1), 17-19.
9. Paetzold, P.; Plotho, C. V.; Schmid, G.; Boese, R.; Schrader, B.; Bougeard, D.; Pfeiffer, U.; Gleiter, R.; Schüfer, W., Darstellung, Reaktionen und Struktur von tert-Butyl(tert-butylimino)boran. *Chem. Ber.* **1984**, *117* (3), 1089-1102.
10. Rosas-Garcia, V. M., The electronic structure of substituted iminoboranes. *J. Theor. Comput. Chem.* **2011**, *967* (1), 160-164.
11. Paetzold, P.; Richter, A.; Thijssen, T.; Würtenberg, S., Bildung, Struktur und Reaktivität von (Pentafluorphenyl)bor-tert-butylimid und seinem Cyclodimeren. *Chem. Ber.* **1979**, *112* (12), 3811-3827.
12. Paetzold, P.; von Plotho, C., Über weitere monomere Borimide und ihre Reaktionen. *Chem. Ber.* **1982**, *115* (8), 2819-2825.
13. Paetzold, P., New perspectives in boron-nitrogen chemistry - I. *Pure Appl. Chem.* **1991**, *63* (3), 345-350.
14. Steuber, E. v.; Elter, G.; Noltemeyer, M.; Schmidt, H.-G.; Meller, A., First B-Organyloxy-Substituted Iminoboranes: Preparation, Stabilization, and Reactivity. *Organometallics* **2000**, *19* (24), 5083-5091.
15. Haase, M.; Klingebiel, U., Simple Synthesis of Stable Iminoboranes. *Angew. Chem. Int. Ed. Engl.* **1985**, *24* (4), 324-324.
16. Guo, R.; Li, T.; Wei, R.; Zhang, X.; Li, Q.; Liu, L. L.; Tung, C.-H.; Kong, L., Boraiminolithium: An Iminoborane-Transfer Reagent. *J. Am. Chem. Soc.* **2021**, *143* (34), 13483-13488.
17. Guo, R.; Zhang, X.; Li, T.; Li, Q.; Ruiz, D. A.; Liu, L. L.; Tung, C.-H.; Kong, L., Unraveling the reactivity of a cationic iminoborane: avenues to unusual boron cations. *Chem. Sci.* **2022**, *13* (8), 2303-2309.
18. Schleier, D.; Schaffner, D.; Gerlach, M.; Hemberger, P.; Fischer, I., Threshold photoelectron spectroscopy of iminoborane, HBNH. *Phys. Chem. Chem. Phys.* **2022**, *24* (1), 20-24.
19. Wang, J.; Jia, P.; Sun, W.; Wei, Y.; Lin, Z.; Ye, Q., Synthesis of Iminoboryl o-Carboranes by Lewis Base Promoted Aminoborirane-to-Iminoborane Isomerization. *Inorg. Chem.* **2022**, *61* (23), 8879-8886.
20. Wrackmeyer, B., The B-N Bond in Some Aminoboranes and an Iminoborane, Studied by ¹¹B and ¹⁵N NMR Spectroscopy and DFT Methods. *Z. Anorg. Allg. Chem.* **2015**, *641* (14), 2525-2529.
21. Bao, M.; Dai, Y.; Liu, C.; Su, Y., Acid/Base-Free Acyclic Anionic Oxoborane and Iminoborane Bearing Diboryl Groups. *Inorg. Chem.* **2022**, *61* (29), 11137-11142.

22. Raiser, D.; Schubert, H.; Bettinger, H. F.; Wesemann, L., Germaborenes: Borylene Transfer Agents for the Synthesis of Iminoboranes. *Chem. Eur. J.* **2021**, *27* (6), 1981-1983.
23. Winner, L.; Ewing, W. C.; Geetharani, K.; Dellermann, T.; Jouppi, B.; Kupfer, T.; Schaefer, M.; Braunschweig, H., Spontaneous Metal-Free Transfer Hydrogenation of Iminoboranes with Ammonia Borane and Amine Boranes. *Angew. Chem. Int. Ed.* **2018**, *57* (38), 12275-12279.
24. Braunschweig, H.; Damme, A.; Jimenez-Halla, J. O. C.; Pfaffinger, B.; Radacki, K.; Wolf, J., Metal-mediated synthesis of 1,4-di-tert-butyl-1,4-azaborine. *Angew. Chem. Int. Ed.* **2012**, *51* (40), 10034.
25. Braunschweig, H.; Ewing, W. C.; Geetharani, K.; Schaefer, M., The Reactivities of Iminoboranes with Carbenes: BN Isosteres of Carbene-Alkyne Adducts. *Angew. Chem. Int. Ed.* **2015**, *54* (5), 1662-1665.
26. Schaefer, M.; Beattie, N. A.; Geetharani, K.; Schaefer, J.; Ewing, W. C.; Krahfuss, M.; Hoerl, C.; Dewhurst, R. D.; Macgregor, S. A.; Lambert, C.; Braunschweig, H., Synthesis of functionalized 1,4-azaborinines by the cyclization of di-tert-butyliminoborane and alkynes. *J. Am. Chem. Soc.* **2016**, *138* (26), 8212-8220.
27. Schaefer, M.; Schaefer, J.; Dewhurst, R. D.; Ewing, W. C.; Krahfuss, M.; Kuntze-Fechner, M. W.; Wehner, M.; Lambert, C.; Braunschweig, H., Regioselective Catalytic and Stepwise Routes to Bulky, Functional-Group-Appended, and Luminescent 1,2-Azaborinines. *Chem. Eur. J.* **2016**, *22* (25), 8603-8609.
28. Ayhan, O.; Eckert, T.; Plamper, F. A.; Helten, H., Poly(iminoborane)s: An Elusive Class of Main-Group Polymers? *Angew. Chem. Int. Ed.* **2016**, *55* (42), 13321-13325.
29. Lorenz, T.; Crumbach, M.; Eckert, T.; Lik, A.; Helten, H., Poly(p-phenylene iminoborane): A Boron-Nitrogen Analogue of Poly(p-phenylene vinylene). *Angew. Chem. Int. Ed.* **2017**, *56* (10), 2780-2784.
30. Ayhan, O.; Riensch, N. A.; Glasmacher, C.; Helten, H., Cycloliner Oligo- and Poly(iminoborane)s: The Missing Link in Inorganic Main-Group Macromolecular Chemistry. *Chem. Eur. J.* **2018**, *24* (22), 5883-5894.
31. Beweries, T.; Helten, H. In *Poly(aminoborane)s and Poly(iminoborane)s*, John Wiley & Sons Ltd.: 2020; p eibc2717.
32. Winner, L.; Hermann, A.; Belanger-Chabot, G.; Gonzalez-Belman, O. F.; Jimenez-Halla, J. O. C.; Kelch, H.; Braunschweig, H., Cleavage of BN triple bonds by main group reagents. *Chem. Commun. (Cambridge, U. K.)* **2018**, *54* (59), 8210-8213.
33. Franz, D.; Inoue, S., Systematic Investigation of the Ring-Expansion Reaction of N-Heterocyclic Carbenes with an Iminoborane Dihydride. *Chem Asian J* **2014**, *9* (8), 2083-2087.
34. Braunschweig, H.; Dewhurst, R. D.; Ewing, W. C.; Kuntze-Fechner, M. W.; Schaefer, M., Isolation of a Reactive Cyclopropane Intermediate via a Unique Catalyst-Free Spontaneous Cyclopropanation Step at 0 °C. *Chem. Eur. J.* **2017**, *23* (25), 5953-5956.
35. Cui, P.; Guo, R.; Kong, L.; Cui, C., New Approaches to N-Heterocyclic-Carbene-Coordinated Iminoborane and Borenum Species. *Inorg. Chem.* **2020**, *59* (8), 5261-5265.
36. Koner, A.; Sergeieva, T.; Morgenstern, B.; Andrada, D. M., A Cyclic Iminoborane-NHC Adduct: Synthesis, Reactivity, and Bonding Analysis. *Inorg. Chem.* **2021**, *60* (18), 14202-14211.
37. Prieschl, D.; Belanger-Chabot, G.; Guo, X.; Dietz, M.; Mueller, M.; Krummenacher, I.; Lin, Z.; Braunschweig, H., Synthesis of Complex Boron-Nitrogen Heterocycles Comprising Borylated Triazenes and Tetrazenes Under Mild Conditions. *J. Am. Chem. Soc.* **2020**, *142* (2), 1065-1076.
38. Noguchi, M.; Suzuki, K.; Kobayashi, J.; Yurino, T.; Tsurugi, H.; Mashima, K.; Yamashita, M., Planar and Bent BN-Embedded p-Quinodimethanes Synthesized by Transmetalation of Bis(trimethylsilyl)-1,4-dihydropyrazines with Chloroborane. *Organometallics* **2018**, *37* (12), 1833-1836.
39. Lui, M. W.; Paisley, N. R.; McDonald, R.; Ferguson, M. J.; Rivard, E., Metal-Free Dehydrogenation of Amine Boranes by Tunable N-Heterocyclic Iminoboranes. *Chem. Eur. J.* **2016**, *22* (6), 2134-2145.
40. Xie, L.; Zhang, J.; Cui, C., 2-Chloro-Azaborolyl Anion: A Source of 1,2-Azaborole Isosteric to Cyclopentadienyldiene. *Chem. Eur. J.* **2014**, *20* (31), 9500-9503.

41. Xie, L.; Zhang, J.; Hu, H.; Cui, C., Synthesis and Reactions of π -Conjugated Iminoboranes Stabilized by Intramolecular Imine Groups. *Organometallics* **2013**, *32* (23), 6875-6878.
42. Thiess, T.; Belanger-Chabot, G.; Fantuzzi, F.; Michel, M.; Ernst, M.; Engels, B.; Braunschweig, H., Diborane(4) Azides: Surprisingly Stable Sources of Transient Iminoboranes. *Angew. Chem. Int. Ed.* **2020**, *59* (36), 15480-15486.
43. Matler, A.; Arrowsmith, M.; Schorr, F.; Hermann, A.; Hofmann, A.; Lenczyk, C.; Braunschweig, H., Reactivity of Terminal Iron Borylenes and Bis(borylenes) with Carbodiimides: Cycloaddition, Metathesis, Insertion and C-H Activation Pathways. *Eur. J. Inorg. Chem.* **2021**, *2021* (45), 4619-4631.
44. Liu, T.-T.; Chen, J.; Guan, B.-T.; Lin, Z.; Shi, Z.-J., Distance-Triggered Distinct Aryl Migrations on Azidodiboranes. *Chem. Eur. J.* **2023**, *29* (13), e202203676.
45. Paetzold, P.; von Plotho, C.; Niecke, E.; Ruger, R., [2 + 2]-Cycloaddition von Iminoboran und Iminophosphan. *Chem. Ber.* **1983**, *116* (4), 1678-1681.
46. Paetzold, P.; Delpy, K.; Hughes, R. P.; Herrmann, W. A., [2 + 2]-Cycloaddition eines Iminoborans an einen Methylen-Titan-Komplex. *Chem. Ber.* **1985**, *118* (4), 1724-1725.
47. Braunschweig, H.; Paetzold, P.; Boese, R., Weitere Reaktionen der Iminoborane $RB\equiv NR'$ mit Neopentylidentantal-Komplexen. *Chem. Ber.* **1993**, *126* (7), 1571-1577.
48. Krockert, B.; van Bonn, K.-H.; Paetzold, P., The Iminoborane $tBuB\equiv NtBu$ as a Dipolarphile in (2+3) Cycloadditions. *Z. Anorg. Allg. Chem.* **2005**, *631* (5), 866-868.
49. Nutz, M.; Borthakur, B.; Dewhurst, R. D.; Deissenberger, A.; Dellermann, T.; Schaefer, M.; Krummenacher, I.; Phukan, A. K.; Braunschweig, H., Synthesis and Trapping of Iminoboranes by M:B/C:N Bond Metathesis. *Angew. Chem. Int. Ed.* **2017**, *56* (27), 7975-7979.
50. Winner, L.; Belanger-Chabot, G.; Celik, M. A.; Schaefer, M.; Braunschweig, H., Intriguing migrations in transient iminoborane adducts: two new pathways to aminoboranes. *Chem. Commun. (Cambridge, U. K.)* **2018**, *54* (67), 9349-9351.
51. Zhang, H.; Wang, J.; Yang, W.; Xiang, L.; Sun, W.; Ming, W.; Li, Y.; Lin, Z.; Ye, Q., Solution-Phase Synthesis of a Base-Free Benzoborirene and a Three-Dimensional Inorganic Analogue. *J. Am. Chem. Soc.* **2020**, *142* (41), 17243-17249.
52. Frenette, B. L.; Omana, A. A.; Ferguson, M. J.; Zhou, Y.; Rivard, E., Access to adducts of parent iminoborane isomers, HBNH and NBH₂, using frustrated Lewis pair chelation. *Chem. Commun. (Cambridge, U. K.)* **2021**, *57* (83), 10895-10898.
53. Wong, A.; Chu, J.; Wu, G.; Telsner, J.; Dobrovetsky, R.; Menard, G., Redox-Controlled Reactivity at Boron: Parallels to Frustrated Lewis/Radical Pair Chemistry. *Inorg. Chem.* **2020**, *59* (14), 10343-10352.
54. Mehta, M.; Goicoechea, J. M., Carboboration of isocyanates with tris(pentafluorophenyl)borane and evidence for dissociative FLP chemistry of an acid-base pair. *Chem. Commun. (Cambridge, U. K.)* **2019**, *55* (48), 6918-6921.
55. Omana, A. A.; Watt, R.; Zhou, Y.; Ferguson, M. J.; Rivard, E., Frustrated Lewis Pair Chelation and Reactivity of Complexed Parent Iminoborane and Aminoborane. *Inorg. Chem.* **2022**, *61* (41), 16430-16440.
56. Swarnakar, A. K.; Hering-Junghans, C.; Nagata, K.; Ferguson, M. J.; McDonald, R.; Tokitoh, N.; Rivard, E., Encapsulating Inorganic Acetylene, HBNH, Using Flanking Coordinative Interactions. *Angew. Chem. Int. Ed.* **2015**, *54* (36), 10666-10669.
57. Swarnakar, A. K.; Hering-Junghans, C.; Ferguson, M. J.; McDonald, R.; Rivard, E., Reactivity of a coordinated inorganic acetylene unit, HBNH, and the azidoborane cation $[HB(N_3)]^+$. *Chem. Sci.* **2017**, *8* (3), 2337-2343.
58. Gilbert, T. M., Ab Initio Computational Studies of Heterocycloalkynes: Structures, Natural Bond Orders, Ring Strain Energies, and Isomerizations of Cyclic Iminoboranes and Iminoalanes. *Organometallics* **2000**, *19* (6), 1160-1165.
59. Gilbert, T. M., Computational Studies of Pericyclic Reactions of Iminoboranes. [4+2] Diels-Alder vs [2+2] Dimerization of $RB:NR'$ Compounds. *Organometallics* **2003**, *22* (11), 2298-2304.

60. Gilbert, T. M., Computational Prediction of Regiospecificity in the [4+2] Diels–Alder Cyclizations between the Iminoborane (F₃C)₃C–B:N–(t-Bu) and Substituted cis-Butadienes. *Organometallics* **2005**, *24* (26), 6445-6449.
61. Münster, J.; Paetzold, P.; Schröder, E.; Schwan, H.; von Bennigsen-Mackiewicz, T., Cyclic Iminoboranes: Analogues of Cycloalkynes. *Z. Anorg. Allg. Chem.* **2004**, *630* (15), 2641-2651.
62. Giustra, Z. X.; Liu, S.-Y., The State of the Art in Azaborine Chemistry: New Synthetic Methods and Applications. *J. Am. Chem. Soc.* **2018**, *140* (4), 1184-1194.
63. Campbell, P. G.; Marwitz, A. J. V.; Liu, S.-Y., Recent Advances in Azaborine Chemistry. *Angew. Chem. Int. Ed.* **2012**, *51* (25), 6074-6092.
64. Edel, K.; Brough, S. A.; Lamm, A. N.; Liu, S.-Y.; Bettinger, H. F., 1,2-Azaborine: The Boron-Nitrogen Derivative of ortho-Benzyne. *Angew. Chem. Int. Ed.* **2015**, *54* (27), 7819-7822.
65. Edel, K.; Fink, R. F.; Bettinger, H. F., Isomerization and fragmentation pathways of 1,2-azaborine. *J. Comput. Chem.* **2016**, *37* (1), 110-116.
66. Edel, K.; Ishibashi, J. S. A.; Liu, S.-Y.; Bettinger, H. F., Superelectrophilicity of 1,2-Azaborine: Formation of Xenon and Carbon Monoxide Adducts. *Angew. Chem. Int. Ed.* **2019**, *58* (12), 4061-4064.
67. Gupta, D.; Bettinger, H. F., Reactions of 1,2-Azaborinine, a BN-Benzyne, with Organic π Systems. *The Journal of Organic Chemistry* **2023**, *88* (13), 8369-8378.
68. Müller, M.; Maichle-Mössmer, C.; Bettinger, H. F., BN-Phenanthryne: Cyclotetramerization of an 1,2-Azaborine Derivative. *Angew. Chem. Int. Ed.* **2014**, *53* (35), 9380-9383.
69. Bettinger, H. F.; Müller, M., Pathways for the cyclotetramerization of dibenz[c,e][1,2]azaborine, a BN-phenanthryne. *J. Phys. Org. Chem.* **2015**, *28* (2), 97-103.
70. Keck, C.; Hahn, J.; Gupta, D.; Bettinger, H. F., Solution Phase Reactivity of Dibenzo[c,e][1,2]azaborinine: Activation and Insertion into Si-E Single Bonds (E=H, OSi(CH₃)₃, F, Cl) by a BN-Aryne. *Chem. Eur. J.* **2022**, *28* (5), e202103614.
71. Pieper, W.; Schmitz, D.; Paetzold, P., Borimide und Borylnitrene beim Zerfall von Diaminoazidoboranen. *Chem. Ber.* **1981**, *114* (12), 3801-3812.
72. Bettinger, H. F.; Bornemann, H., Donor stabilized borylnitrene: A highly reactive BN analogue of vinylidene. *J. Am. Chem. Soc.* **2006**, *128* (34), 11128-11134.
73. Bettinger, H. F.; Filthaus, M.; Bornemann, H.; Oppel, I. M., Metal-Free Conversion of Methane and Cycloalkanes to Amines and Amides by Employing a Borylnitrene. *Angew. Chem. Int. Ed.* **2008**, *47* (25), 4744-4747.
74. Bettinger, H. F.; Filthaus, M., Borylnitrenes: electrophilic reactive intermediates with high reactivity towards C–H bonds. *Organic & Biomolecular Chemistry* **2010**, *8* (24), 5477-5482.
75. Filthaus, M.; Schwertmann, L.; Neuhaus, P.; Seidel, R. d. W.; Oppel, I. M.; Bettinger, H. F., C–H Bond Amination by Photochemically Generated Transient Borylnitrenes at Room Temperature: A Combined Experimental and Theoretical Investigation of the Insertion Mechanism and Influence of Substituents. *Organometallics* **2012**, *31* (10), 3894-3903.
76. Müller, M.; Maichle-Mössmer, C.; Bettinger, H. F., C–H functionalization of tetramethylsilane employing a borylnitrene. *Chem. Comm.* **2013**, *49* (100), 11773-11775.
77. Bettinger, H. F.; Filthaus, M.; Neuhaus, P., Insertion into dihydrogen employing the nitrogen centre of a borylnitrene. *Chem. Comm.* **2009**, (16), 2186-2188.
78. Bonn, K.-H. v.; Bennigsen-Mackiewicz, T. v.; Kiesgen, J.; Plotho, C. v.; Paetzold, P., Neue Iminoborane und ihre Reaktionen / Novel Iminoboranes and their Reactions. *Zeitschrift für Naturforschung B* **1988**, *43* (1), 61-68.
79. NOETH, H.; Weber, S., CONTRIBUTION TO THE CHEMISTRY OF BORON. 137. A CONVENIENT SYNTHESIS OF AN AMINOIMINOBORANE AND ITS ADDUCTS WITH BORON TRICHLORIDE AND BORON TRIBROMIDE. *Chemischer Informationsdienst* **1984**, *15* (10), no-no.

80. Chrostowska, A.; Xu, S.; Lamm, A. N.; Mazière, A.; Weber, C. D.; Dargelos, A.; Baylère, P.; Graciaa, A.; Liu, S.-Y., UV-Photoelectron Spectroscopy of 1,2- and 1,3-Azaborines: A Combined Experimental and Computational Electronic Structure Analysis. *J. Am. Chem. Soc.* **2012**, *134* (24), 10279-10285.
81. Edel, K.; Yang, X.; Ishibashi, J. S. A.; Lamm, A. N.; Maichle-Mössmer, C.; Giustra, Z. X.; Liu, S.-Y.; Bettinger, H. F., The Dewar Isomer of 1,2-Dihydro-1,2-azaborinines: Isolation, Fragmentation, and Energy Storage. *Angew. Chem. Int. Ed.* **2018**, *57* (19), 5296-5300.
82. Chen, Z.; Wannere, C. S.; Corminboeuf, C.; Puchta, R.; Schleyer, P. v. R., Nucleus-Independent Chemical Shifts (NICS) as an Aromaticity Criterion. *Chem. Rev.* **2005**, *105* (10), 3842-3888.
83. Schleyer, P. v. R.; Maerker, C.; Dransfeld, A.; Jiao, H.; van Eikema Hommes, N. J. R., Nucleus-Independent Chemical Shifts: A Simple and Efficient Aromaticity Probe. *J. Am. Chem. Soc.* **1996**, *118* (26), 6317-6318.
84. Gershoni-Poranne, R.; Stanger, A., Magnetic criteria of aromaticity. *Chem. Soc. Rev.* **2015**, *44* (18), 6597-6615.
85. Papadopoulos, A. G.; Charistos, N. D.; Kyriakidou, K.; Sigalas, M. P., Study of Electron Delocalization in 1,2-, 1,3-, and 1,4-Azaborines Based on the Canonical Molecular Orbital Contributions to the Induced Magnetic Field and Polyelectron Population Analysis. *J. Phys. Chem. A* **2015**, *119* (39), 10091-10100.
86. Mayer, R. J.; Hampel, N.; Ofial, A. R., Lewis Acidic Boranes, Lewis Bases, and Equilibrium Constants: A Reliable Scaffold for a Quantitative Lewis Acidity/Basicity Scale. *Chem. Eur. J.* **2021**, *27* (12), 4070-4080.
87. Sivaev, I. B.; Bregadze, V. I., Lewis acidity of boron compounds. *Coordination Chemistry Reviews* **2014**, *270-271*, 75-88.
88. Döring, S.; Erker, G.; Fröhlich, R.; Meyer, O.; Bergander, K., Reaction of the Lewis acid tris (pentafluorophenyl) borane with a phosphorus ylide: competition between adduct formation and electrophilic and nucleophilic aromatic substitution pathways. *Organometallics* **1998**, *17* (11), 2183-2187.
89. Massey, A.; Park, A., Perfluorophenyl derivatives of the elements: VII. further studies on tris (pentafluorophenyl) boron. *Journal of Organometallic Chemistry* **1966**, *5* (3), 218-225.
90. Bondi, A. v., van der Waals volumes and radii. *J. Phys. Chem.* **1964**, *68* (3), 441-451.
91. Mantina, M.; Chamberlin, A. C.; Valero, R.; Cramer, C. J.; Truhlar, D. G., Consistent van der Waals radii for the whole main group. *J. Phys. Chem. A* **2009**, *113* (19), 5806-5812.
92. Reed, A. E.; Curtiss, L. A.; Weinhold, F., Intermolecular interactions from a natural bond orbital, donor-acceptor viewpoint. *Chem. Rev.* **1988**, *88* (6), 899-926.
93. Glendening, E.; Landis, C.; Weinhold, F., Wiley Interdiscip. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **2012**, *2* (1), 1-42.
94. Weinhold, F., Natural bond orbital analysis: A critical overview of relationships to alternative bonding perspectives. *J. Comput. Chem.* **2012**, *33* (30), 2363-2379.
95. Glendening, E. D.; Landis, C. R.; Weinhold, F., NBO 6.0: Natural bond orbital analysis program. *J. Comput. Chem.* **2013**, *34* (16), 1429-1437.