

Shining light on inverted singlet-triplet emitters

Matteo Bedogni, Davide Giavazzi, Francesco Di Maiolo,* and Anna Painelli

*Department of Chemistry, Life Science and Environmental Sustainability, Università di
Parma, 43124 Parma, Italy.*

E-mail: francesco.dimaiolo@unipr.it

Abstract

The inversion of the lowest singlet and triplet excited states, observed in several triangle-shaped organic molecules containing conjugated carbon and nitrogen atoms, is an astonishing result that implies the breakdown of Hund's rule. The phenomenon attracted interest for its potential towards triplet harvesting in organic LEDs. On a more fundamental vein, the singlet-triplet (ST) inversion sheds new light on the role of electron correlations in the excited state landscape of π -conjugated molecules. Relying on the celebrated Pariser-Parr-Pople model, the simplest model for correlated electrons in π -conjugated systems, we demonstrate that the ST inversion does not require triangle-shaped molecules nor any specific molecular symmetry. Indeed, the ST inversion does not require strictly non-overlapping HOMO and LUMO orbitals, but rather a small gap and a small exchange integral between the frontier orbitals.

1 Introduction

In closed shell molecular systems, the Hund's multiplicity rule would locate the first excited singlet state S_1 at higher energy than the corresponding triplet state T_1 , thus leading to a positive singlet-triplet (ST) gap ΔE_{ST} . However, recent experimental¹ and theoretical²⁻¹⁰

studies have shown that several triangle-shaped π -conjugated molecules (also called triangles) decorated with nitrogen atoms display a negative ΔE_{ST} , thus violating Hund's rule. Indeed, in these systems the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO) are marginally overlapping and the lowest energy transition has a dominant multi resonant charge-transfer (MR-CT) character, with the electronic charge being transferred from HOMO to LUMO. In these conditions, the exchange integral, responsible for the ST splitting, is tiny, and the spin polarization correction, typically negligible, may be large enough to revert the ST gap.¹¹⁻¹⁵

The family of MR-CT dyes is interesting in several respects. Even systems with the normal order of singlet and triplet states have small ΔE_{ST} , making it possible to thermally populate the S_1 state from the T_1 state.¹⁶⁻¹⁹ Fluorescent MR-CT dyes are then intensively investigated as emitters featuring thermally activated delayed fluorescence (TADF). They are of special interest for OLED applications, since the reduced polarity of the MR-CT state and the molecular rigidity guarantee for very narrow and marginally solvatochromic emission spectra, properties in demand for emitters for (blue) OLED with high color purity.^{1,2,20-31} Here we will not address the family of MR-CT dyes at large, but will focus on the specific issue of ST inversion. The theoretical interest in molecules breaking Hund's rule is backed up by their applicative potential. The reverted ST order, in fact, allows for easy and efficient triplet harvesting in organic light-emitting diodes (OLEDs), increasing their maximum internal efficiency from a disappointing 25% to a reassuring 100%.²⁰

Spin-polarization, the mechanism responsible for ST-inversion, is due to electron correlation so that conventional TD-DFT approaches to excited states are inadequate to address ST inversion, as it was early pointed out.^{2,32,33} High-quality, computationally expensive methods have been exploited so far to deal with ST-inversion, including CIS(D),^{3,6} SCS-CC2,^{2,3,8,9} EOM-CCSD,^{2,4,5,34,35} SCS-ADC(2),^{2-4,7,8} CASPT2,² SA-CASSCF,^{3,8} SC-NEVPT2^{3,4,8} and Bethe-Salpeter equation in the GW approximation.¹⁰

In an effort to better understand the physics underlying the ST inversion, we undertook

an extensive study of several molecular structures described by the celebrated Pariser-Parr-Pople (PPP) model Hamiltonian, the simplest model for correlated electrons in π -conjugated molecules. The model, proposed in the 1950s,^{36–38} successfully described the behavior of several families of small organic molecules.^{37,39–47} In the field of π -conjugated polymers, PPP was pivotal in understanding the anomalous behavior of polyacetylene,⁴⁸ where correlated electrons are responsible for the appearance of a low-lying dark state, thus solving the puzzle of its non-fluorescent behavior. More recently, density matrix renormalization group (DMRG) approaches allowed to extend the PPP model to deal with graphene-based structures,⁴⁹ to address singlet-fission in polyenes⁵⁰ and to build models for dark states in polyene and carotenoid systems.⁵¹

Here, we exploit the comparative simplicity of the PPP approach to understand ST inversion in triangulenes and related molecules. Comparing with exact diagonalization results, we critically discuss the validity of the double configuration interactions (CI) and related approaches, typically adopted to deal with *all electrons* models of triangulenes. The adopted PPP model cannot provide quantitatively accurate results on specific molecules. However, it is a solid and elegant approach that, applied here to a variety of molecular structures, will disclose qualitative trends about the effects of electron correlations in π -conjugated molecules. The novel insights gained along these lines will contribute to establish general and reliable guidelines towards the rational design of molecules featuring ST inversion. As an additional bonus, we show that information easily obtained at the Hartree-Fock (HF) level (including the HOMO-LUMO gap and exchange energy) offers safe guidance to recognize molecular structures with negative ΔE_{ST} . An easy-to-use computational tool is made available for a quick screening of interesting structures.




Figure 1: Sketches of the molecular structures of two prototypical triangulenes: 2T-N, also called cyclazine, and 2T-7N, also called heptazine.

2 Theoretical and computational background

Much as Hückel model, PPP model describes the physics of π -electrons in planar conjugated molecules in terms of a minimal basis set that only accounts, on each atomic site, for just the p orbital orthogonal to the molecular plane. At variance with the Hückel model, PPP also accounts for electron-electron (e-e) interaction in the zero differential overlap (ZDO) approximation. The PPP Hamiltonian reads:

$$\begin{aligned} \hat{H}_{PPP} = & \sum_{\mu} \varepsilon_{\mu} \hat{n}_{\mu} - \sum_{\mu\nu, \nu > \mu} \sum_{\sigma} t_{\mu\nu} (\hat{a}_{\mu\sigma}^{\dagger} \hat{a}_{\nu\sigma} + \hat{a}_{\nu\sigma}^{\dagger} \hat{a}_{\mu\sigma}) \\ & + \sum_{\mu} U_{\mu} \hat{n}_{\mu\uparrow} \hat{n}_{\mu\downarrow} + \sum_{\mu\nu, \nu > \mu} V_{\mu\nu} (Z_{\mu} - \hat{n}_{\mu}) (Z_{\nu} - \hat{n}_{\nu}) \end{aligned} \quad (1)$$

where μ and ν run on atomic sites, $\hat{a}_{\mu,\sigma}^{(\dagger)}$ is the fermionic annihilation (creation) operator for an electron with spin σ in the atomic orbital (AO) on site μ , and $\hat{n}_{\mu} = \sum_{\sigma} \hat{a}_{\mu\sigma}^{\dagger} \hat{a}_{\mu\sigma}$ counts the total number of electrons on site μ . The first two terms of the above Hamiltonian are single electron terms, with ε_{μ} measuring the energy of the on-site orbital and $t_{\mu\nu}$ the hopping integral between sites μ and ν . As in the Hückel model, PPP only accounts for hopping integrals between sites connected by a σ -bond. The remaining terms in the above Hamiltonian describe e-e interaction: U_{μ} measures the repulsion between two electrons on the same site, $V_{\mu\nu}$ is the repulsion between electrons on different sites and Z_{μ} measures the charge on atom μ upon removal of the π -electrons ($Z_{\mu} = 1$ for carbon and aza nitrogen

atoms, $Z_\mu = 2$ for pyrrole nitrogens). We adopt the Ohno expression for $V_{\mu\nu}$:^{42,45,52}

$$V_{\mu\nu} = \frac{e^2}{4\pi\epsilon_0} \left[r_{\mu\nu}^2 + \left(\frac{\epsilon_r e^2}{4\pi\epsilon_0(U_\mu + U_\nu)} \right)^2 \right]^{-1/2} \quad (2)$$

where $\epsilon_r = 2$ is the relative dielectric constant, as relevant to organic media.⁴⁵

The PPP Hamiltonian can be written on the so-called real space (RS) basis, comprising all states obtained by distributing the n π -electrons on the N atoms. In the valence bond (VB) approach, the basis states are combined, following the Pauling rules, to be eigenstates of the total spin operator, \hat{S}^2 , so that the Hamiltonian factorizes in subspaces having the same S value.^{53,54} The VB approach is extremely efficient as it allows to work with comparatively small matrices, but the VB basis is non-orthogonal and hence difficult to manage. On the other hand, in the RS representation, the basis states are selected as eigenstates of the z -component spin operator, \hat{S}_z . The RS subspaces are therefore much larger than the VB subspaces, but the basis set is orthogonal. For closed shell systems, as discussed here, diagonalizing the PPP Hamiltonian in the $S_z = 0$ and 1 subspaces, allows to single out triplet (and higher spin) states as those states that are present in both subspaces. The dimension of the RS basis grows fast with N , leading to very large Hamiltonian matrices, that however are very sparse. Exploiting specific algorithms for the storage and diagonalization of sparse matrices,⁵⁵ the PPP Hamiltonian for the 2T-N and 2T-7N molecules in Fig. 1 can be diagonalized numerically, in spite of a basis that in the $S_z = 0$ subspace contains 2,944,656 states.

The RS diagonalization leads to exact *full-CI* results. However, in order to address larger systems and for a direct comparison with current quantum chemical approaches to triangulene-like systems, we adopt a molecular orbital (MO) approach to the PPP Hamiltonian. Specifically, in the Hartree Fock (HF) approximation the PPP Hamiltonian in Eq. 1

reduces to the one-electron Fock operator:

$$\hat{\mathcal{F}}_{PPP} = \sum_{\mu} (\varepsilon_{\mu} + J_{\mu\mu} - K_{\mu\mu}) \hat{n}_{\mu} + \sum_{\mu\nu, \mu \neq \nu} (-t_{\mu\nu} - K_{\mu\nu}) \sum_{\sigma} (\hat{a}_{\mu\sigma}^{\dagger} \hat{a}_{\nu\sigma} + \hat{a}_{\nu\sigma}^{\dagger} \hat{a}_{\mu\sigma}) \quad (3)$$

where, in line with the PPP model, we exploited the ZDO approximation, so that the Coulomb operator has only diagonal terms:

$$J_{\mu\mu} = \sum_{\lambda=1}^N (P_{\lambda\lambda} - Z_{\lambda}) V_{\lambda\mu} \quad (4)$$

while the exchange operator has elements:

$$K_{\mu\nu} = (P_{\mu\nu}/2 - Z_{\nu} \delta_{\mu\nu}) V_{\mu\nu} \quad (5)$$

In Eqs. 4 and 5 the matrix elements of the density operator are defined as :

$$P_{\mu\nu} = 2 \sum_k c_{k\mu} c_{k\nu} \quad (6)$$

where k runs on the occupied MOs in the ground state configuration and $c_{k\mu}$ are the expansion coefficients of the k -th MO on the AOs (see ESI, Eqs. S1 and S2). The HF Hamiltonian is solved self-consistently. Typically, 40-50 HF iterations are needed to reach convergence on the density matrix elements.

Once the HF MOs are obtained, as relevant to the ground state configuration, $|g\rangle$, single, double, triple, etc. excited configurations are defined transferring one, two, three, etc. electrons from the occupied to the virtual MOs. The PPP Hamiltonian is then written on the basis defined by the different configurations and diagonalized in the CI approach (see ESI, Sec. S1). The PPP Hamiltonian on the CI basis is far less sparse than on the RS basis, so that full-CI is only feasible for very small molecules (up to ~ 12 π -electrons on 12 atoms). However, as it will be shown below, the results converge quickly with the number

of excitations, making it possible to obtain reliable results on fairly large systems.

The diagonalization of the CI Hamiltonian leads to the molecular excited states $|f\rangle$, as needed to calculate optical spectra. The electric dipole moment operator reads:

$$\hat{\vec{\mu}} = \hat{\mu}_x \vec{i} + \hat{\mu}_y \vec{j} = \sum_{\mu} (Z_{\mu} - \hat{n}_{\mu})(x_{\mu} \vec{i} + y_{\mu} \vec{j}) \quad (7)$$

where x_{μ}, y_{μ} are the Cartesian coordinates of site μ in the molecular plane. Absorption spectra are calculated assigning a Gaussian bandshape to each transition:

$$A(\omega) \propto \omega \sum_f |\mu_{fg}|^2 \exp \left[-\frac{(\omega - \omega_{fg})^2}{2\sigma^2} \right] \quad (8)$$

where the sum runs over the excited eigenstates and the transition energies and dipole moments are $\omega_{fg} = E_f - E_g$ and $\mu_{fg} = \langle f | \hat{\vec{\mu}} | g \rangle$, respectively (in the following $\sigma = 0.1$ eV).

3 Results and discussion

3.1 2T-N and 2T-7N as reference systems

To start with, we target the two triangulene structures in Fig. 1. In recent years, these molecules have been extensively investigated theoretically as prototypical examples of ST inversion.^{2,7,32,56} Experimental absorption spectra of both dyes, collected in the 80's,^{57,58} will be exploited to parametrize and validate the PPP model. PPP parameters for C atoms, derived in the 1950s,³⁷ are well established and transferable among different molecules.^{42,45,53,59} Specifically, setting the on-site energy of C atoms to zero, we adopt the standard value for the on-site e-e interaction, $U_C = 11.26$ eV, and for the nearest-neighbor C-C hopping, $t = 2.4$ eV. As for nitrogen atoms, a unique set of PPP parameters is not available.^{41-45,53,60-66} We set the hopping integral for C-N bonds to the same value as for C-C bonds.⁵³ Typically, the on-site e-e repulsion is slightly larger for N atoms than that for C atoms.⁴⁵ Best agreement

with experimental data is obtained here setting $U_N \sim 15$ eV and 15.5 eV for pyrrole and aza nitrogens, respectively. These choices marginally affect the results (see ESI, Section S2). As for the molecular geometry, for the sake of simplicity, we set all angles to 120° and all bond lengths to 1.4 \AA (results for different geometries are marginally different, see ESI Fig. 2S). The most delicate issue is the on-site energy of the N atoms. We fix the site energy of the aza nitrogen $\varepsilon_N^{aza} = -5$ eV and adjust the pyrrole nitrogen site energy ε_N^{py} as to best simulate experimental spectra (Fig. 3S shows results obtained for different ε_N^{aza} values).

Experimental absorption spectra of 2T-N and 2T-7N from Refs.^{57,58} are shown as red lines in Figs. 2a and 2d, respectively. For both molecules, the lowest energy absorption band is very weak and is ascribed to a forbidden transition that gains intensity through vibronic coupling. This dark transition is located at ~ 1 - 1.5 eV and ~ 2.6 - 3 eV for 2T-N and 2T-7N, respectively. An intense and sharp transition is seen in 2T-N at ~ 2.7 eV followed by a broad absorption for energies higher than ~ 3.9 eV. The 2T-7N spectrum is similar but blue-shifted, with the first intense absorption observed at 4 - 4.3 eV and a second broad absorption at energies higher than ~ 5.3 eV. In the same figure, superimposed to experimental spectra, we show spectra calculated accounting for up to triple excitations (i.e., CISDT), as to ensure converged results (see below). Results are shown for different values of the on-site energy of the triangulene central nitrogen, ε_N^{py} . For both molecules, the lowest energy state, S_1 , is a dark state (A'_2 symmetry in the D_{3h} point group) and its location is marked by a cross in the figure.

For 2T-N (Fig. 2a), upon increasing the electron-withdrawing nature of the central N atom (i.e., for more negative ε_N^{py}) the dark S_1 state moves to the red. Similarly, the highest energy excitation redshifts, while the band at intermediate energy, ascribed to S_2 , a doubly degenerate E' state, is marginally affected by ε_N^{py} . For 2T-N, acceptable agreement with experiment is observed in the region $-13 \text{ eV} < \varepsilon_N^{py} < -11 \text{ eV}$. Quite interestingly, Fig. 2b shows that in this region (highlighted in orange) the ST gap is consistently negative.

Moving to 2T-7N (Fig. 2d), the dark state again moves to the red when ε_N^{py} becomes more




Figure 2: Simulating the photophysics of 2T-N (panels a-c) and 2T-7N (panels d-f). (a) Absorption spectra of 2T-N: red lines are experimental data from Ref.,⁵⁷ colored lines show results for several ϵ_N^{py} values (same color code as in panel b), the crosses in the low energy region mark the position of the dark state. Calculated spectra are normalized to the maximum absorbance of the most intense spectrum. (b) 2T-N singlet-triplet energy gaps calculated for different ϵ_N^{py} values, the shaded area highlights ϵ_N^{py} values giving acceptable agreement between calculated and experimental absorption spectra. (c) 2T-N frontier HF-MO for two values of ϵ_N^{py} , the corresponding exchange integral K_{H-L} , and energy gaps, ΔE_{H-L} . (d) Absorption spectra of 2T-7N: red lines are experimental data from Ref.,⁵⁸ colored lines show results for several ϵ_N^{py} values (same color code as in panel e), the crosses in the low energy region mark the position of the dark state. Calculated spectra are normalized to the maximum absorbance of the most intense spectrum. (e) The same as (b) for 2T-7N. (f) The same as (c) for 2T-7N.

negative, but, at variance with 2T-N, the intermediate $S_2 \leftarrow S_0$ transition is also affected by ε_N^{py} , with the corresponding peak moving to the blue when ε_N^{py} becomes more negative. An acceptable agreement with experiment is obtained for ε_N^{py} values ranging from -14 eV to -12 eV (the region highlighted in orange in panel e). So, we will stick to $\varepsilon_N^{py} = -13$ eV as a good compromise to describe both 2T-N and 2T-7N. In the following, when not otherwise stated, results will be shown for this value of ε_N^{py} . Quite interestingly, the sign of the ST gap of 2T-7N in the relevant region goes from negative values, $\Delta E_{ST} = -0.22$ eV at $\varepsilon_N^{py} = -14$ eV, to positive values, $\Delta E_{ST} = +0.26$ eV at $\varepsilon_N^{py} = -12$ eV. Accordingly, the ST inversion, safely assessed in 2T-N, is marginal in 2T-7N.

The HF MOs of 2T-N and 2T-7N in Fig. 2c and f, respectively, give interesting clues. When the ST gap is negative (points ① in Fig. 2b and e), the HOMO and LUMO are localized on two complementary sublattices and the HF HOMO-LUMO exchange energy K_{H-L} is as low as -0.04 eV (2T-N) and -0.02 eV (2T-7N). In either case, the LUMO+1 orbital is doubly degenerate and is clearly not complementary to HOMO. When ΔE_{ST} becomes positive (points ② in Fig. 2b and e), the nature of LUMO and LUMO+1 is reversed, so that the HOMO-LUMO complementarity is lost, and sizable K_{H-L} values are calculated (-0.5 eV for 2T-N and -0.23 eV for 2T-7N). We stress that this notable reversal in the nature of LUMO and LUMO+1 is not shown by the Hückel MOs, demonstrating once again that electron correlations are the key to ST inversion.

The different behavior observed for 2T-N and 2T-7N in terms of ΔE_{ST} reflects the diverse nature of the triangulene rim in the two cases. In Fig. 3, we show the dependence of the ST gap on the parameters that define the triangulene rim, namely ε_N^{aza} and U_N^{aza} . The red dot on the map corresponds to the 2T-N parameters, where all rim atoms are C atoms. If, starting from this position, we decorate the odd sites on the rim with electron-withdrawing atoms, whose on-site energy becomes more and more negative, the ST inversion is lost when ε_N^{aza} becomes more negative than ~ -2.1 eV. However, increasing the on site electron repulsion from the value relevant to C atoms to that relevant to N atoms, the ST inversion is favored.




Figure 3: The color map shows the dependence of the ST gap (color code on the right) as a function of ϵ_N^{aza} and U_N^{aza} , having set $\epsilon_N^{py} = -13$ eV. White lines at $\Delta E_{ST} = 0$ mark the boundary between the ST inverted region (dark blue) and the normal region (light blue and yellow). Red and green dots indicate the parameters relevant to 2T-N and 2T-7N, respectively. Calculation done at the PPP CISDT level with model parameters defined in the main text.

2T-7N, with its six aza nitrogens on the rim, is precisely on the verge of the ST inverted region (green dot on the map). This last observation contrasts with the results of many all-electron calculations that typically predict a more pronounced ST inversion in 2T-7N than in 2T-N..^{2-4,6,30,67,68} Of course, PPP model is not an exact model, and results are affected by the specific choice of the molecular geometry and model parameters (see ESI, Section S2 for specific illustrative cases). However, it is important to stress that all-electron calculations mentioned above all rely on up to double excitations. In these approaches, large uncertainties on $\pi \rightarrow \pi^*$ transition energies are estimated (e.g., ~ 0.11 eV for CCSD and ~ 0.17 eV for ADC(2)),⁶⁹ making the error on ΔE_{ST} of the same order of magnitude as the calculated gap

itself.

Indeed, we are in the position to address the reliability of commonly adopted all-electron quantum chemical calculations, while validating at the same time our approach. As for 2T-N and 2T-7N, all-electron calculations are typically truncated to single and double electronic CIs. Among them, we mention CIS(D), an approach that accounts for single CI, and introduces double CI perturbatively, spin-component-scaled second order coupled cluster (SCS-CC2), spin-component-scaled second order algebraic diagrammatic construction (SCS-ADC(2)),^{2,3,7,56} double-hybrid DFT⁶⁸ and DFT multireference configuration interaction (DFT-MRCI).⁷⁰ All these approaches confirm the important role played by double CI to properly describe the ST inversion. However, the fast increase in the number of multiple CI precludes higher order studies using all-electron wavefunction- or density-based approaches. The PPP model, with its minimal representation of π -electrons, lends itself quite naturally to explore higher order CIs. In Fig. 4 a and b, we show the total weight of single (S), double (D), triple (T) and quadruple (Q) CIs in the lowest seven eigenstates of 2T-N and 2T-7N. For both molecules, as expected, single and triple CI marginally contribute to the ground state (according to the Brillouin theorem in fact single and triple CI are not directly mixed to the ground state configuration, and marginally contribute to it via their indirect mixing to double CI). As for excited states, in both molecules, double and triple CIs enter the picture with a similar and sizable weight. This is a strong indication that a reliable approach to the excited states of these molecules should include at least triple CIs. On the other hand, the weight of quadruple CIs stays very small, suggesting that fourth and higher order CIs can be safely neglected.

Truncating the expansion at the CISD level, the ground state is stabilized, but the stabilization of the excited states as due to the triple CI contribution is missing. As a result, CISD transition energies are overestimated, a spurious effect that is fixed when triple excitations are accounted for (see panels c and d). This result is fully confirmed by the comparison with RS exact diagonalization transition energies, that perfectly match with




Figure 4: Relative weights of single (S), double (D), triple (T), quadruple (Q) excitations for the first seven singlet and triplet eigenstates of 2T-N (panel a) and 2T-7N (panel b) calculated at the CISDTQ level, together with the excitation energies of the first few electronic states of 2T-N (panel c) and 2T-7N (panel d) at different levels of theory. Relevant results from exact diagonalization in the real space basis (full-CI) are shown as colored dots in panels c and d. Model parameters defined in the main text.

CISDT results. However, reaching triple CI in all-electron quantum chemical calculations is extremely demanding. Most often CIS(D) approaches, where double CIs are introduced perturbatively, seem to provide reasonable results as for transition energies. However, caution is needed since the wavefunctions stay at the CIS level. On the other hand, CIS(D) introduces




Figure 5: The effect of ϵ_N^{py} on the energy levels of 2T-4N. Panel a: Absorption spectra calculated for different ϵ_N^{py} values (same color code used in panel b). The crosses mark the position of the first excited (dark) state. Spectra are normalized with respect to the maximum absorbance of the most intense spectrum. Panel b: Calculated singlet-triplet energy gaps. Panel c: Frontier Hartree-Fock MOs for two different ϵ_N^{py} values. Relevant HF HOMO-LUMO exchange energy K_{H-L} and energy gaps ΔE_{H-L} are also reported. Other parameters are defined in the main text. Results obtained at the PPP CISDT level.

perturbative corrections ranging from ~ 1 eV to ~ 2 eV on the singlet states of 2T-N and 2T-7N (see ESI, Sec. S3, for ab initio CIS(D) results), shedding doubts on the reliability of a perturbative treatment of these two molecules.

Fig. 4 tells us another interesting story. In either 2T-N and 2T-7N, the singlet and triplet excited states with MR character, $^1A'_2$ and $^3A'_2$, respectively, have the normal order (positive ST gap) at the CIS level, but, as soon as correlations enter into play, from CISD up, their order is inverted, with the triplet MR state ($^3A'_2$) lying at energy higher than the relevant singlet ($^1A'_2$). However, there is another pair of states, $^3E'$ and $^1E'$, whose energy is not too far. In particular, in the case of 2T-7N, the two triplets $^3A'_2$ and $^3E'$ invert their energies. In this situation, while the relative energy of the two MR state would still give a negative ST gap, the presence of another non-MR triplet at low energy makes the ST inversion critical.




Figure 6: Decorating the 2T-N rim with an increasing number of aza nitrogen atoms while changing the pyrrole nitrogen site energy. Calculated absorption spectra (panels a and d), ST gap (panels b and e) and HF frontier orbitals (panels c and f) are shown for 2T-3N and 2T-4N'. In panels a and d, spectra are normalized with respect to the maximum absorbance of the most intense spectrum, while the band relevant to the first excited singlet state is magnified by 50. In panels c and f, we report the relevant HOMO-LUMO exchange energies, K_{H-L} , and gaps, ΔE_{H-L} . Calculations done at PPP CISDT level for the same parameters used in Fig. 4.

3.2 ST inversion: an exploratory study

PPP is a reliable and computationally inexpensive approach to π -conjugated molecules that, easily allowing to explore a large number of structures, helps to shed light on the physics of

the ST inversion. We start our analysis with 2T-4N, in Fig. 5, a molecule having the same D_{3h} symmetry as 2T-N and 2T-7N, but a qualitatively different behavior as for the ST gap. As in 2T-N and 2T-7N, the lowest energy transition is dark. The allowed transitions move first to the red when the orbital on the central N atom is stabilized (ϵ_N^{py} going from -11 eV to -13 eV), and then blue-shift for further stabilization of the central N orbital (-14 eV $> \epsilon_N^{py} > -15$ eV), as shown in Fig. 5a. More interestingly, in this case, at variance with either 2T-N or 2T-7N, if the central nitrogen orbital is stabilized, i.e. ϵ_N^{py} is made more negative, the ST gap becomes positive (see panel b) and the HOMO/LUMO nature is switched (see panel c). Indeed, for 2T-4N, the HOMO and LUMO show a tiny overlap, thus leading to a fairly large exchange energy $K_{H-L} \sim -0.1$ eV, reducing the MR character of the low-lying excitation. As a result, the ST inversion region is strongly suppressed, and the calculated ST gap is only marginally negative.




Figure 7: The role of central connectivity on the ST gap. Panel a: Calculated ST gap as a function of the number of broken central bonds for both 2T-N (black curve) and 2T-7N (red curve). Calculations done at the PPP CISDT level. Panel b: Frontier HF MOs are shown for the two annulenes obtained by breaking all three central bonds in 2T-N and 2T-7N. Relevant HOMO-LUMO exchange energies and gaps are also shown. Same model parameters as in Fig. 4.

The molecules discussed so far, 2T-N, 2T-7N and 2T-4N, all belong to the D_{3h} point group

and the question arises if the ST inversion is somehow related to the molecular symmetry. This is an important point, since in these three molecules the lowest excited singlet is a dark (i.e., optically forbidden) state. These molecules are therefore very weakly fluorescent, so that, in spite of the ST inversion, they are of little interest for OLED applications. Upon lowering the symmetry, the S_1 state can acquire oscillator strength, and, if ST inversion is maintained, extremely promising fluorescent dyes for OLED with 100% internal quantum efficiency could be devised.

To address this issue, in Fig. 6, we decorated the external rim of 2T-N with a variable number of N atoms. The two molecules, 2T-3N and 2T-4N' (structural isomer of the system in Fig. 5), have C_{2v} and C_S symmetries, respectively, so that degenerate states are not possible. Comparing with Fig. 2 panel a, as relevant to 2T-N, all spectral features move to the blue upon increasing the number of N atoms in the external rim. Most importantly, the lowest energy transition, strictly forbidden in either 2T-N or 2T-7N, acquires sizable intensity. Quite interestingly, despite the lack of complementarity between the HOMO and LUMO, the exchange energy is fairly small when the electron-withdrawing nature of the central N atom is large. As a result, a region with ST inversion is obtained for both molecules at large negative ϵ_N^{py} . Indeed, the best descriptor for ST inversion is not the complementarity of the frontier orbitals, but rather the value of the HOMO-LUMO exchange energy.

In a recent work,⁷¹ the simultaneous presence of the C_2 axis and σ_v plane was suggested as essential to ST inversion, thus making C_{2v} the lowest symmetry point group of molecules breaking the Hund's multiplicity rule. Indeed, results on 2T-4N' in Fig. 6 show that comparatively large and negative ΔE_{ST} values can be achieved even in a system where the only symmetry element is the molecular plane, but where the non-complementary frontier orbitals do minimize the HOMO-LUMO exchange integral.

Breaking some of the bonds that connect the central N atom in 2T-N and 2T-7N to the rim, as shown in Fig. 7, is another way to reduce the molecular symmetry. Quite interestingly, breaking bonds in 2T-N quickly drives it from the region with inverted ST

order to the normal region, while the opposite occurs for 2T-7N. As expected, the lowest-lying singlet state acquires sizable intensity when one or two bonds are broken and the symmetry of the system reduced. Of course when all bonds are broken the D_{3h} symmetry is recovered and the lowest energy states are dark (see ESI, Sec. S4).

Possibly, the most surprising result from Fig. 7 is the observation that just the 2T-7N rim, without the central N atom, has the largest negative ST gap, suggesting that the triangulene structure itself is not really needed to observe the ST inversion. On the other hand, coming back to our prototypical dyes, 2T-N and 2T-7N in Fig. 2, it is clear that the ST inversion is favored when the on-site energy on the central N is more negative, i.e. in systems where the two electrons from the central N are marginally involved in the rim physics. Fig. 7 panel b shows the HF frontier orbitals calculated for the cyclododecahexaene and hexaazacyclododecahexaene molecules, corresponding to the rims of the 2T-N and 2T-7N molecules. In the cyclododecahexaene the lack of spatial separation between HOMO and LUMO is reflected in a large exchange energy: the molecules (a simple cyclic polyene) does not qualify as an MR-CT dye. On the other hand, the frontier orbitals of the hexaazacyclododecahexaene variant, with their small K_{H-L} value, confirm its MR-CT nature.

3.3 ST inversion: a fresh view of spin polarization

Figures 6 and 7 suggest that the observation of an inverted ST gap is not intrinsically related to the symmetry of the molecule, nor to the presence of the connectivity of *triangulene-like* structures, rather it is amplified in $(CN)_x$ rings. $(CN)_x$ rings are a very useful playground to understand the phenomenon of dynamical spin polarization, pivotal to ST inversion. Dynamical spin polarization was early discussed^{11–15,72} in the framework of π -electron models demonstrating very clearly that electron correlations govern ST inversion, but that at least double excitations are needed to capture the phenomenon. To shed light on the elusive dynamical spin polarization issue, we performed calculations on $(CN)_x$ rings, putting the $2x$ atoms in the vertices of regular polygons with bond lengths equal to 1.4 Å. Fig. 8 shows




Figure 8: Results for rings with different numbers of aza nitrogen atoms. Upper panel: ST gap values. The red curve refers to PPP CISDT results. Dashed blue line refers to results from exact diagonalization in the RS basis (full-CI). Lower panel: HF HOMO-LUMO gap for the different aza-doped rings. Same parameters used in Fig. 4.

the evolution of the singlet triplet gap and of the HF HOMO-LUMO gap for rings with a variable number of atoms from 4 to 12. A couple of observations are in order. As expected, triple-CI results are basically equivalent to full-CI results, at variance with double-CI results (see ESI, Fig. 6S). The adopted idealized geometry is not realistic for all structures, but for the triazine molecule (i.e., $x = 3$) whose equilibrium geometry corresponds to a (almost) regular hexagon.⁷³ For this specific molecule, it is therefore interesting to compare the PPP ST gap value, ~ 0.65 eV, with available results from all-electron calculations, pointing to a positive ST gap, ~ 0.9 eV. The agreement is far from perfect, but definitely acceptable for a semiempirical approach.

The most important result of Fig. 8 is the almost perfect correlation between the HF HOMO-LUMO gap and the ST gap. The $4n/4n+2$ alternancy of the HOMO-LUMO gap is a clear remnant of the well-known alternancy of cyclic polyenes. Indeed, in cyclic polyenes, either at the Hückel or HF level, the HOMO-LUMO gap, sizable for $4n+2$ rings, vanishes for $4n$ rings. In $(\text{CN})_x$ rings a splitting of the HOMO-LUMO gap is found also in $4n$ rings, but it stays lower than for $4n+2$ rings (see ESI, Fig. 5S). In all cases, the HOMO-LUMO gap reduces upon increasing the ring dimension.

We are now in the position to understand the correlation between the two gaps in Fig. 8. The lowest excited singlet and triplet states in our closed shell systems are dominated by configurations where one electron is transferred from the HOMO to the LUMO. The exchange energy, that in most systems is responsible for triplet states lying lower than singlets, is negligible in these alternating rings (see ESI, Section S6) and the typically tiny effect of dynamical spin polarization may enter into play inverting the ST gap. Specifically, the singlet excited state is stabilized by the interaction with the doubly excited state, where both electrons reside on the LUMO. This stabilization is not possible for the corresponding triplet states. Fig. 8, showing an impressive correlation between the size of the HOMO-LUMO gap and the ST gap, neatly confirms this view: the singlet stabilization is larger the smaller is the HOMO-LUMO gap and the only ring with positive ST gap is triazine $(\text{CN})_3$, with the largest HOMO-LUMO gap. The picture is further confirmed by our results for 2T-N and 2T-7N, whose ST gap, ~ -0.2 and 0 eV, respectively, can be related to a much smaller HOMO-LUMO gap (~ 6.9 eV) for 2T-N than for 2T-7N (~ 9.8 eV). In the end, despite the adopted idealized geometries, the rings in Fig. 8 show how a molecular edge with alternating electron-withdrawing and electron-donor groups is a key ingredient for the ST inversion.

4 Conclusions

The inversion of the lowest excited singlet and triplet states involves breaking the Hund's rule and deserves some explanation. Typically, triplets have lower energy than corresponding singlets because of the exchange energy. Accordingly, the first requirement for ST inversion is a small exchange energy between HOMO and LUMO. To this end, a possible strategy consists in having non-overlapping HOMO and LUMO, so that the two orbitals have sizable weight on two different subsets of atoms. Systems with disconnected HOMO and LUMO belong to the large and interesting family of MR-CT dyes. However, our results show quite explicitly that the disjoint nature of the HOMO and LUMO is not a strict prerequisite for ST inversion. A more reliable indication of the propensity to the breaking of the Hund's rule is recognized in the HOMO-LUMO exchange energy itself.

To observe ST inversion, spin polarization is needed to bring the energy of the lowest excited singlet below the energy of the lowest triplet in systems with small HOMO-LUMO exchange energy. Spin polarization is driven by electron correlations, and cannot be modeled in single-CI approaches. Accordingly, TD-DFT, arguably the most widely applied quantum chemical approach to excited states in molecular systems, fails to describe the ST inversion that requires at least double excitations. By exploiting the PPP model, we investigated the role of higher order CIs, showing that reliable results on inverted ST systems can only be obtained if at least triple excitations are accounted for (see Fig. 4).

We studied several triangulene-based molecules to grasp the physics of the ST inversion. Starting from the highly symmetric and very weakly fluorescent 2T-N, 2T-7N (see Fig. 2) and 2T-4N (see Fig. 5), we moved to fluorescent 2T-3N and 2T-4N' (see Fig. 6), obtained by decorating the edge of 2T-N with two and three N atoms, respectively. In all cases, we found non-negligible regions of the parameter space showing ST inversion. We demonstrated that neither the presence of C_2 axis and σ_v plane, nor the very specific connectivity of triangulenes are actually needed for ST inversion.

The key ingredient for a negative singlet-triplet gap is a molecular edge with alternating

electron-donor (D) and electron-acceptor (A) groups. In 2T-N, where the rim is made of 12 carbon atoms, the central nitrogen creates an alternating pattern of electron poor and electron rich sites on the rim. Accordingly, upon gradually breaking the bonds that connect the central nitrogen to the edge until the central nitrogen is completely decoupled from the rim, one recovers a cyclic polyene structure, then losing the ST inversion (cf. Fig. 7). In 2T-7N, instead, the aza nitrogens along the rim create by themselves an alternating pattern and the central nitrogen atom is not actually needed to observe ST inversion. Rather, in 2T-7N the central N atom is detrimental to ST inversion, putting 2T-7N in the difficult region where determining the sign of the ST gap is tricky, as it fluctuates depending on the specific choice of model parameters (see Fig. 3). In the PPP π -electron only picture, large $(\text{CN})_x$ rings in an idealized planar and regular geometry are the most promising structures for ST inversion.

Triple-CI models, needed to properly deal with ST inversion, are computationally very demanding as to make their exploitation in explorative studies prohibitive. However, our analysis demonstrates that good indicators for ST inversion can be already obtained at the HF level. Specifically, a computationally inexpensive (ground state) calculation of the molecular MOs, gives access to both the HOMO-LUMO gap and to the relevant exchange integral. Systems with small exchange integrals and small gaps are promising for ST inversions and should be considered for additional theoretical analysis and/or for experimental work. An easy-to-use (PPP-based) computational tool is made publicly available (see ESI, Sec. S6) to offer the synthetic chemist a very easy and computationally inexpensive approach to identify the most promising molecular structures with inverted ST gap.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgement

We acknowledge financial support from PNRR MUR project ECS-00000033-ECOSISTER. This work benefited from the equipment and framework of the COMP-R Initiative, funded by the *Departments of Excellence* program of the Italian Ministry for University and Research (MUR, 2023-2027) and from the support of the HPC (High Performance Computing) facility at the University of Parma. F.D.M. position was co-funded by the European Union - PON Research and Innovation 2014-2020.

Supporting Information Available

Electronic Supplementary Information (ESI) available: Sec. S1 contains the derivation of the PPP model on the MO basis; Sec. S2 collects additional results obtained for different model parameters values as well as different molecular geometries (i.e., angles and bond lengths); Sec. S3 reports ab initio CIS(D) results for 2T-N and 2T-7N; Sec. S4 collects optical spectra calculated for the systems in Fig. 7; Sec. S5 contains some additional results for $(\text{CN})_x$ rings; Sec. S6 contains the HOMO-LUMO exchange integral expression on the MO and AO basis. An easy-to-use (PPP-based) computational tool is made publicly available for the calculation of K_{H-L} .

References

- (1) Aizawa, N.; Pu, Y.-J.; Harabuchi, Y.; Nihonyanagi, A.; Ibuka, R.; Inuzuka, H.; Dhara, B.; Koyama, Y.; Nakayama, K.-i.; Maeda, S.; Araoka, F.; Miyajima, D. Delayed fluorescence from inverted singlet and triplet excited states. *Nature* **2022**, *609*, 502–506.
- (2) Ehrmaier, J.; Rabe, E. J.; Pristash, S. R.; Corp, K. L.; Schlenker, C. W.;

- Sobolewski, A. L.; Domcke, W. Singlet–triplet inversion in heptazine and in polymeric carbon nitrides. *The Journal of Physical Chemistry A* **2019**, *123*, 8099–8108.
- (3) Ricci, G.; San-Fabián, E.; Olivier, Y.; Sancho-García, J. C. Singlet-Triplet Excited-State Inversion in Heptazine and Related Molecules: Assessment of TD-DFT and *ab initio* Methods. *ChemPhysChem* **2021**, *22*, 553–560.
- (4) Pollice, R.; Friederich, P.; Lavigne, C.; dos Passos Gomes, G.; Aspuru-Guzik, A. Organic molecules with inverted gaps between first excited singlet and triplet states and appreciable fluorescence rates. *Matter* **2021**, *4*, 1654–1682.
- (5) Bhattacharyya, K. Can TDDFT render the electronic excited states ordering of Azine derivative? A closer investigation with DLPNO-STEOM-CCSD. *Chemical Physics Letters* **2021**, *779*, 138827.
- (6) Ghosh, S.; Bhattacharyya, K. Origin of the Failure of Density Functional Theories in Predicting Inverted Singlet–Triplet Gaps. *The Journal of Physical Chemistry A* **2022**, *126*, 1378–1385.
- (7) Sobolewski, A. L.; Domcke, W. Are heptazine-based organic light-emitting diode chromophores thermally activated delayed fluorescence or inverted singlet–triplet systems? *The Journal of Physical Chemistry Letters* **2021**, *12*, 6852–6860.
- (8) Sanz-Rodrigo, J.; Ricci, G.; Olivier, Y.; Sancho-Garcia, J.-C. Negative singlet–triplet excitation energy gap in triangle-shaped molecular emitters for efficient triplet harvesting. *The Journal of Physical Chemistry A* **2021**, *125*, 513–522.
- (9) Hall, D.; Sancho-García, J. C.; Pershin, A.; Ricci, G.; Beljonne, D.; Zysman-Colman, E.; Olivier, Y. Modeling of Multiresonant Thermally Activated Delayed Fluorescence Emitters Properly Accounting for Electron Correlation Is Key! *Journal of Chemical Theory and Computation* **2022**, *18*, 4903–4918.

- (10) Monino, E.; Loos, P.-F. Connections and performances of Green's function methods for charged and neutral excitations. **2023**, <https://doi.org/10.48550/arXiv.2305.16959>.
- (11) Kollmar, H.; Staemmler, V. A theoretical study of the structure of cyclobutadiene. *Journal of the American Chemical Society* **1977**, *99*, 3583–3587.
- (12) Kollmar, H.; Staemmler, V. Violation of Hund's rule by spin polarization in molecules. *Theoretica Chimica Acta* **1978**, *48*, 223–239.
- (13) Koseki, S.; Nakajima, T.; Toyota, A. Violation of Hund's multiplicity rule in the electronically excited states of conjugated hydrocarbons. *Canadian Journal of Chemistry* **1985**, *63*, 1572–1579.
- (14) Toyota, A.; Nakajima, T. Violation of Hund's multiplicity rule in the lowest excited singlet–triplet pairs of cyclic bicalicene and its higher homologues. *J. Chem. Soc., Perkin Trans. 2* **1986**, 1731–1734.
- (15) Toyota, A. Violation of Hund's rule in the lowest excited singlet-triplet pairs of dicyclohepta[cd,gh]pentalene and dicyclopenta[ef,kl]heptalene. *Theoretica Chimica Acta* **1988**, *74*, 209–217.
- (16) Shizu, K.; Kaji, H. Comprehensive understanding of multiple resonance thermally activated delayed fluorescence through quantum chemistry calculations. *Communications Chemistry* **2022**, *5*.
- (17) Eng, J.; Penfold, T. J. Open questions on the photophysics of thermally activated delayed fluorescence. *Communications Chemistry* **2021**, *4*.
- (18) Kim, I.; Cho, K. H.; Jeon, S. O.; Son, W.-J.; Kim, D.; Rhee, Y. M.; Jang, I.; Choi, H.; Kim, D. S. Three States Involving Vibronic Resonance is a Key to Enhancing Reverse Intersystem Crossing Dynamics of an Organoboron-Based Ultrapure Blue Emitter. *JACS Au* **2021**, *1*, 987–997.

- (19) Northey, T.; Penfold, T. The intersystem crossing mechanism of an ultrapure blue organoboron emitter. *Organic Electronics* **2018**, *59*, 45–48.
- (20) Won, T.; Nakayama, K.-i.; Aizawa, N. Inverted singlet–triplet emitters for organic light-emitting diodes. *Chemical Physics Reviews* **2023**, *4*, 021310.
- (21) Li, J.; Zhang, Q.; Nomura, H.; Miyazaki, H.; Adachi, C. Thermally activated delayed fluorescence from $^3n\pi^*$ to $^1n\pi^*$ up-conversion and its application to organic light-emitting diodes. *Applied Physics Letters* **2014**, *105*, 013301.
- (22) Wu, X.; Su, B.-K.; Chen, D.-G.; Liu, D.; Wu, C.-C.; Huang, Z.-X.; Lin, T.-C.; Wu, C.-H.; Zhu, M.; Li, E. Y.; Hung, W.-Y.; Zhu, W.; Chou, P.-T. The role of host–guest interactions in organic emitters employing MR-TADF. *Nature Photonics* **2021**, *15*, 780–786.
- (23) Jiang, H.; Cao, Y.; Yang, Q.; Xian, L.; Tao, Y.; Chen, R.; Huang, W. Organic Resonance Materials: Molecular Design, Photophysical Properties, and Optoelectronic Applications. *The Journal of Physical Chemistry Letters* **2020**, *11*, 7739–7754.
- (24) Suresh, S. M.; Hall, D.; Beljonne, D.; Olivier, Y.; Zysman-Colman, E. Multiresonant Thermally Activated Delayed Fluorescence Emitters Based on Heteroatom-Doped Nanographenes: Recent Advances and Prospects for Organic Light-Emitting Diodes. *Advanced Functional Materials* **2020**, *30*, 1908677.
- (25) Liu, Y.; Xiao, X.; Ran, Y.; Bin, Z.; You, J. Molecular design of thermally activated delayed fluorescent emitters for narrowband orange–red OLEDs boosted by a cyano-functionalization strategy. *Chemical Science* **2021**, *12*, 9408–9412.
- (26) Wu, S.; Li, W.; Yoshida, K.; Hall, D.; Suresh, S. M.; Sayner, T.; Gong, J.; Beljonne, D.; Olivier, Y.; Samuel, I. D. W.; Zysman-Colman, E. Excited-State Modulation in Donor-Substituted Multiresonant Thermally Activated Delayed Fluorescence Emitters. *ACS Applied Materials & Interfaces* **2022**, *14*, 22341–22352.

- (27) Li, J.; Li, Z.; Liu, H.; Gong, H.; Zhang, J.; Li, X.; Wang, Y.; Guo, Q. Down-conversion-induced delayed fluorescence via an inverted singlet-triplet channel. *Dyes and Pigments* **2022**, *203*, 110366.
- (28) Gao, H.; Shen, S.; Qin, Y.; Liu, G.; Gao, T.; Dong, X.; Pang, Z.; Xie, X.; Wang, P.; Wang, Y. Ultrapure Blue Thermally Activated Delayed Fluorescence (TADF) Emitters Based on Rigid Sulfur/Oxygen-Bridged Triarylboron Acceptor: MR TADF and D–A TADF. *The Journal of Physical Chemistry Letters* **2022**, *13*, 7561–7567.
- (29) Sano, Y.; Shintani, T.; Hayakawa, M.; Oda, S.; Kondo, M.; Matsushita, T.; Hatakeyama, T. One-Shot Construction of BN-Embedded Heptadecacene Framework Exhibiting Ultra-narrowband Green Thermally Activated Delayed Fluorescence. *Journal of the American Chemical Society* **2023**, *145*, 11504–11511.
- (30) Jin, J.; Duan, C.; Jiang, H.; Tao, P.; Xu, H.; Wong, W.-Y. Integrating Asymmetric O-B-N Unit in Multi-Resonance Thermally Activated Delayed Fluorescence Emitters towards High-Performance Deep-Blue Organic Light-Emitting Diodes. *Angewandte Chemie International Edition* **2023**, *62*, 18.
- (31) Wu, S.; Zhang, L.; Wang, J.; Kumar Gupta, A.; Samuel, I. D. W.; Zysman-Colman, E. Merging Boron and Carbonyl based MR-TADF Emitter Designs to Achieve High Performance Pure Blue OLEDs. *Angewandte Chemie International Edition* *n/a*, e202305182.
- (32) de Silva, P. Inverted singlet–triplet gaps and their relevance to thermally activated delayed fluorescence. *The Journal of Physical Chemistry Letters* **2019**, *10*, 5674–5679.
- (33) Pershin, A.; Hall, D.; Lemaire, V.; Sancho-Garcia, J.-C.; Muccioli, L.; Zysman-Colman, E.; Beljonne, D.; Olivier, Y. Highly emissive excitons with reduced exchange energy in thermally activated delayed fluorescent molecules. *Nature Communications* **2019**, *10*, 597.

- (34) Pratik, S. M.; Coropceanu, V.; Brédas, J.-L. Purely Organic Emitters for Multiresonant Thermally Activated Delay Fluorescence: Design of Highly Efficient Sulfur and Selenium Derivatives. *ACS Materials Letters* **2022**, *4*, 440–447.
- (35) Pratik, S. M.; Coropceanu, V.; Brédas, J.-L. Enhancement of Thermally Activated Delayed Fluorescence (TADF) in Multi-Resonant Emitters via Control of Chalcogen Atom Embedding. *Chemistry of Materials* **2022**, *34*, 8022–8030.
- (36) Pariser, R.; Parr, R. G. A Semi-Empirical Theory of the Electronic Spectra and Electronic Structure of Complex Unsaturated Molecules. I. *The Journal of Chemical Physics* **1953**, *21*, 466–471.
- (37) Pariser, R.; Parr, R. G. A Semi-Empirical Theory of the Electronic Spectra and Electronic Structure of Complex Unsaturated Molecules. II. *The Journal of Chemical Physics* **1953**, *21*, 767–776.
- (38) Pople, J. A. Electron interaction in unsaturated hydrocarbons. *Transactions of the Faraday Society* **1953**, *49*, 1375.
- (39) Pariser, R. Theory of the Electronic Spectra and Structure of the Polyacenes and of Alternant Hydrocarbons. *The Journal of Chemical Physics* **1956**, *24*, 250–268.
- (40) Pariser, R. Electronic Spectrum and Structure of Azulene. *The Journal of Chemical Physics* **1956**, *25*, 1112–1116.
- (41) Favini, G.; Vandoni, I.; Simonetta, M. Calculation of electronic spectra of aza-benzenes and aza-naphthalenes by the Pariser-Parr-Pople method. *Theoretica Chimica Acta* **1965**, *3*, 45–58.
- (42) Albert, I. D. L.; Ramasesha, S.; Das, P. K. Properties of some low-lying electronic states in polymethineimines and poly(2,3-diazabutadienes). *Physical Review B* **1991**, *43*, 7013–7019.

- (43) Mukhopadhyay, S.; Topham, B. J.; Soos, Z. G.; Ramasesha, S. Neutral and Charged Excited States in Polar Organic Films: Origin of Unusual Electroluminescence in Tri-*p*-tolylamine-Based Hole Conductors. *The Journal of Physical Chemistry A* **2008**, *112*, 7271–7279.
- (44) Kumar, M.; Pati, Y. A.; Ramasesha, S. A density matrix renormalization group method study of optical properties of porphines and metalloporphines. *The Journal of Chemical Physics* **2012**, *136*, 014112.
- (45) Thomas, S.; Pati, Y.; Ramasesha, S. Linear and nonlinear optical properties of expanded porphyrins: A DMRG study. *The Journal of Physical Chemistry A* **2013**, *117*, 7804–7809.
- (46) Aryanpour, K.; Shukla, A.; Mazumdar, S. Electron correlations and two-photon states in polycyclic aromatic hydrocarbon molecules: A peculiar role of geometry. *The Journal of Chemical Physics* **2014**, *140*, 104301.
- (47) Bhattacharyya, P.; Rai, D. K.; Shukla, A. Pariser–Parr–Pople Model Based Configuration-Interaction Study of Linear Optical Absorption in Lower-Symmetry Polycyclic Aromatic Hydrocarbon Molecules. *The Journal of Physical Chemistry C* **2020**, *124*, 14297–14305.
- (48) Soos, Z. G.; Ramasesha, S.; Galvão, D. S. Band to correlated crossover in alternating Hubbard and Pariser-Parr-Pople chains: Nature of the lowest singlet excitation of conjugated polymers. *Physical Review Letters* **1993**, *71*, 1609–1612.
- (49) Goli, V. M. L. D. P.; Prodhan, S.; Mazumdar, S.; Ramasesha, S. Correlated electronic properties of some graphene nanoribbons: A DMRG study. *Physical Review B* **2016**, *94*, 035139.
- (50) Valentine, D. J.; Manawadu, D.; Barford, W. Higher-energy triplet-pair states in

- polyenes and their role in intramolecular singlet fission. *Physical Review B* **2020**, *102*, 125107.
- (51) Barford, W. Theory of the dark state of polyenes and carotenoids. *Physical Review B* **2022**, *106*, 035201.
- (52) Ohno, K. Some remarks on the Pariser-Parr-Pople method. *Theoretica chimica acta* **1964**, *2*, 219–227.
- (53) Soos, Z. G.; Ramasesha, S. Valence-bond theory of linear Hubbard and Pariser-Parr-Pople models. *Physical Review B* **1984**, *29*, 5410–5422.
- (54) Ramasesha, S. Electron-electron interactions in polyacetylene. *Journal of Chemical Sciences* **1986**, *96*, 509–521.
- (55) Lehoucq, R.; Sorensen, D.; Yang, C. *ARPACK Users' Guide*; Society for Industrial and Applied Mathematics, 1998.
- (56) Pios, S.; Huang, X.; Sobolewski, A. L.; Domcke, W. Triangular boron carbon nitrides: An unexplored family of chromophores with unique properties for photocatalysis and optoelectronics. *Physical Chemistry Chemical Physics* **2021**, *23*, 12968–12975.
- (57) Leupin, W.; Wirz, J. Low-lying electronically excited states of cycl[3.3.3]azine, a bridged 12.πi.-perimeter. *Journal of the American Chemical Society* **1980**, *102*, 6068–6075.
- (58) Halpern, A. M.; Rossman, M. A.; Hosmane, R. S.; Leonard, N. J. Photophysics of the $S_1 \longleftrightarrow S_0$ transition in tri-s-triazine. *The Journal of Physical Chemistry* **1984**, *88*, 4324–4326.
- (59) Prodhan, S.; Soos, Z. G.; Ramasesha, S. Model for triplet state engineering in organic light emitting diodes. *The Journal of Chemical Physics* **2014**, *140*, 214313.
- (60) Nishimoto, K.; Forster, L. S. SCFMO calculations of heteroatomic systems with the variable? approximation. *Theoretica Chimica Acta* **1966**, *4*, 155–165.

- (61) Michl, J.; Koutecky, J.; Becker, R. S.; Earhart, C. E. A note on the parameters for heteroatoms in Pariser-Parr-Pople (PPP) calculations. *Theoretica Chimica Acta* **1970**, *19*, 92–97.
- (62) Hinze, J.; Beveridge, D. L. Parametrization of semiempirical π -electron molecular orbital calculations. π Systems containing carbon, nitrogen, oxygen, and fluorine. *Journal of the American Chemical Society* **1971**, *93*, 3107–3114.
- (63) Zahradník, R.; Tesařová, I.; Pancíř, J. Experimental and theoretical (HMO and LCI-SCF) study of singlet-triplet transitions in conjugated hydrocarbons and their derivatives. *Collection of Czechoslovak Chemical Communications* **1971**, *36*, 2867–2880.
- (64) Griffiths, J. Practical aspects of colour prediction of organic dye molecules. *Dyes and Pigments* **1982**, *3*, 211–233.
- (65) Grossjean, M. F.; Tavan, P. Wavelength regulation in bacteriorhodopsin and halorhodopsin: A Pariser–Parr–Pople multireference double excitation configuration interaction study of retinal dyes. *The Journal of Chemical Physics* **1988**, *88*, 4884–4896.
- (66) Albert, I. D. L.; Das, P. K.; Ramasesha, S. Optical nonlinearities in symmetric cyanine dyes and related systems. *Journal of the Optical Society of America B* **1993**, *10*, 1365.
- (67) Sancho-García, J.-C.; San-Fabián, E. Organic Emitters Showing Excited-States Energy Inversion: An Assessment of MC-PDFT and Correlation Energy Functionals Beyond TD-DFT. *Computation* **2022**, *10*, 13.
- (68) Sancho-García, J. C.; Brémond, E.; Ricci, G.; Pérez-Jiménez, A. J.; Olivier, Y.; Adamo, C. Violation of Hund’s rule in molecules: Predicting the excited-state energy inversion by TD-DFT with double-hybrid methods. *The Journal of Chemical Physics* **2022**, *156*, 034105.

- (69) Loos, P.-F.; Lipparini, F.; Boggio-Pasqua, M.; Scemama, A.; Jacquemin, D. A Mountaineering Strategy to Excited States: Highly Accurate Energies and Benchmarks for Medium Sized Molecules. *Journal of Chemical Theory and Computation* **2020**, *16*, 1711–1741.
- (70) Dinkelbach, F.; Bracker, M.; Kleinschmidt, M.; Marian, C. M. Large Inverted Singlet–Triplet Energy Gaps Are Not Always Favorable for Triplet Harvesting: Vibronic Coupling Drives the (Reverse) Intersystem Crossing in Heptazine Derivatives. *The Journal of Physical Chemistry A* **2021**, *125*, 10044–10051.
- (71) Ricci, G.; Sancho-García, J.-C.; Olivier, Y. Establishing design strategies for emissive materials with an inverted singlet–triplet energy gap (INVEST): a computational perspective on how symmetry rules the interplay between triplet harvesting and light emission. *Journal of Materials Chemistry C* **2022**, *10*, 12680–12698.
- (72) Borden, W. T. Can a square or effectively square singlet be the ground state of cyclobutadiene. *Journal of the American Chemical Society* **1975**, *97*, 5968–5970.
- (73) VÉril, M.; Scemama, A.; Caffarel, M.; Lipparini, F.; Boggio-Pasqua, M.; Jacquemin, D.; Loos, P.-F. scpQUESTDB/scp : A database of highly accurate excitation energies for the electronic structure community. *WIREs Computational Molecular Science* **2021**, *11*, 5.