

Halogen Bonding and Mechanochemistry Combined: Synthesis, Characterization, and Application of *N*-Iodosaccharin Pyridine Complexes

Christian Schumacher,^{††a,b} Khai-Nghi Truong,^{†§a} Jas S. Ward,^{†a} Anssi Rajala,^a Elias Lassila,^a Carsten Bolm,^b Kari Rissanen^{*a}

Halogen-bonded complexes are utilized across a myriad of synthetic chemistry fields, with halogen(I) complexes such as Barluenga's reagent ubiquitous with respect to halogenation reactions. Especially the preparation of Barluenga's reagent requires the use of heavy metal salts and vast amounts of chlorinated solvents. In line with a more modern, environmentally-conscious ethos, halogen bonded adducts and a halogen(I) complex similar to Barluenga's reagent based on *N*-iodosaccharin were prepared by mechanochemical processes for the first time. The general absence of solvents or the use of vanishingly small amounts of ethyl acetate in a liquid-assisted grinding approach during the mechanochemical preparations, enabled the homoleptic [(DMAP)–I–(DMAP)]⁺ iodine(I) complex to be synthesized. The as prepared mechanochemically materials were used in the iodination of antipyrine, demonstrating their potential use as surrogates for Barluenga's reagent in both solution and solid-state synthesis.

Introduction

Noncovalent interactions are an unambiguously important aspect in nature, as strikingly demonstrated by proteins and the double helix structure of the DNA.^{1–3} Hence, noncovalent interactions are extensively studied in the field of natural and material sciences.^{2,4,5} Out of the plethora of these interactions, the concept of σ -hole bonding gained vast interest in the last years,^{6–10} halogen bonding being the most prominent ambassador. Already discovered in the 19th century and albeit not named so,¹¹ halogen bonding was a main aspect in the Nobel prize awarded works of Mulliken on the chemical bond (1966) and fundamental to the conformational studies of Hassel (1969).^{11–20} Since then, the concept of halogen bonding made its way into various fields,¹¹ e.g., supramolecular chemistry,^{21–26} crystal engineering,^{27–30} liquid crystals,^{31–35} polymers,^{36–39} gels,^{40–44} biomolecular systems,^{45–49} or organocatalysis.^{50–56}

Nowadays, a halogen bond (XB) is defined as the net attractive interaction between the electrophilic region of a halogen, the halogen bond donor, and a Lewis base, the halogen bond acceptor.⁵⁷ The nature of this attraction is discussed to be mostly based on dispersion,^{58–63} charge transfer,^{64–68} and electrostatic interactions.^{69–73} Spanning a broad range of interaction energies,¹¹ it ranges from weak but distinctive interactions between quinuclidine and iodobenzene [or (iodoethynyl)benzenes (ca. -1.1 kJ·mol⁻¹)]^{74,75} to strong interactions found in *N*-halosaccharin pyridine *N*-oxide complexes (up to -120.3 kJ·mol⁻¹).⁷⁶ Additionally, XBs offer distinct characteristics, such as high directionality (interaction angle of $\sim 180^\circ$), short contacts between donor and acceptor [shorter than the sum of the van der Waals (vdW) radii], and tuneability.^{11,57} As it presents comparable features, XB is often referred to as the "(long lost) brother" of hydrogen bonding (HB),⁷⁷ endowing it the potential to become as widely applicable.^{11,50–56}

Considering these similarities, Fourmigué, Espinosa, and coworkers rose the question, if halogen bonded adducts based on a 1:1 stoichiometry between XB donor and acceptor are co-crystals or salts (Figure 1a).⁷⁸ As a matter of fact, so far, a salt

formation was only observed, when dihalogens were reacted with pyridines (and thiones), leading to *N*-iodopyridinium salts, bonded to the halide anion by a XB.^{79–83}

In the Fourmigué's study it was reasoned that the position of the iodine atom in these systems ($D-I\cdots A \rightleftharpoons [D]\cdots[I-A]^+$) would define the degree of ionicity. To favor strong halogen bonds, electron-donating substituents were attached to the C4-position of the pyridines increasing the electron density on the sp² nitrogen making it a stronger XB acceptor.^{78,84} A "close-to-neutral" co-crystal for halogen bonded complexes between *N*-iodosuccinimide and pyridine derivatives was found, whereas a "close-to-ionic" nature was observed, when *N*-iodosaccharin (NISac), a much stronger XB donor,⁷⁶ was used. The predominant ionic character is well-founded by a much shorter distance between the iodine and the sp² nitrogen atom of the pyridine derivative than the nitrogen atom of the saccharin part. Compared to the "close-to-ionic" nature, a genuine ionic





Figure 1. (a) Schematic representation of the co-crystal and "close-to-ionic" coordination mode in NISac based XB complexes. (b) Molecular structure of Barluenga's reagent (CCDC 202334; thermal ellipsoids shown at 50% probability level; distances are given in Å).^{94–101} (c) Possible asymmetric and symmetric coordination modes of a three-center-four-electron halogen bond.

form by charge conservation is found in symmetrical trihalide anions and in *bis*(pyridine)iodine(I) cations.^{85–93} One of the most prominent examples is *bis*(pyridine)iodine(I) tetrafluoroborate, Barluenga's reagent (Figure 1b).^{94–101} Since its reemergence in 1985,⁹⁴ it (and analogous complexes) have found a myriad of applications in organic synthesis for oxidations, or as versatile electrophilic halogenation reagents,^{94–101} e.g., with alkenes, alkynes, and aromatics.^{102–110} Due to the three-center four-electron bond, the [N–I–N]⁺ iodine(I) complexes differ from the "classical XB" by manifesting a strong simultaneous interaction (≥ -100 kJ·mol⁻¹)^{76,111} between two electron donor moieties and the electrophilic iodine atom. This three-center-four-electron halogen bond should not be referred to as a "coordinative halogen bond" anymore,^{86–91,112} simply because halogen(I) complexes and coordinative metal complexes behave differently. Being a three-center complex, an asymmetric or symmetric coordination mode for the iodine(I) is possible (Figure 1c). Therefore, such complexes have been analyzed in detail by the groups of Rissanen and Erdélyi in the solid-state, as well as in solution. In these studies, a highly symmetric coordination mode was found, regardless of the counteranion, substituent effects, or the solvent.^{85–91,111} Only when two different pyridines, covalently bonded by an aryldiyl linker, viz. a clamp ligand, is used, an asymmetric halogen bond was formed locating the iodine(I) closer to the more electron-rich pyridine.¹¹³ Then, in 2020, the first unrestrained, asymmetric halogen-bonded iodine(I) complexes were reported in the solid-state by Ward *et al.*¹¹⁴ However, scrambling between asymmetric and symmetric complexes in solution occurred, demonstrating the lability and reactivity of these complexes in solution.^{115,116}

Usually, *bis*(pyridine)halogen(I) complexes like Barluenga's reagent are prepared in a two-step Ag(I) to I(I) cation exchange reaction.^{117–122} Initially, the corresponding Ag(I) complex is formed, which then reacts with molecular iodine to form the desired iodine(I) complex in a metathesis reaction. Not only does this approach involve the use of heavy and expensive metal salts (originally, even HgO has been applied)⁹⁴ but also generates a tremendous amount of by-products (*i.e.*, precipitated AgI or AgBr) while using an extensive volume of chlorinated solvent.¹¹⁷ Motivated by the description of a "close-to-ionic" coordination mode in NISac based XB complexes,⁷⁸ the preparation of *bis*(pyridine)iodine(I) complexes by an exchange reaction of the saccharinato anion with another pyridine-based XB acceptor (Scheme 1) was targeted.¹²³ Not only would this approach avoid the use of heavy metals but hopefully lead to a straightforward synthesis of symmetric iodine(I) complexes making use of the entatic iodine atom in the XB complex.



Scheme 1. Schematic representation of the intended exchange reaction in XB complexes with a "close-to-ionic" halogen bond to prepare symmetric and asymmetric iodine(I) complexes.

Results and Discussion

Preparation of NISac-based XB complexes in solution

The investigation began via the preparation of four halogen bonded complexes between NISac (**1**) and pyridine derivatives **2**. 4-Dimethylaminopyridine (**2a**, DMAP), 4-pyrrolidinopyridine (**2b**, PPY), 4-morpholinopyridine (**2c**, MPY), and 4-piperidinopyridine (**2d**, PiPY) were selected to prepare the corresponding XB complexes **3a** – **3d** that would offer a "close-to ionic" XB based on the results of Fourmigué's study. After slow evaporating solutions containing equal amounts of XB donor **1** and the selected pyridine derivative **2**, crystals suitable for single crystal X-ray diffraction (SCXRD) analysis were obtained and compared to literature structures where available.¹²⁴ The corresponding X-ray crystal structures of the desired XB complexes **3a** – **3d** are shown in Figure 2.

In all cases XB complexes with a 1:1 stoichiometry and close to linear coordination geometry were formed with a very short N–I···N' halogen bond. Even though the distance between donor and acceptor does not linearly correlate with the interaction strength, the normalized contact distance R_{XB} of the prepared XB complexes was calculated to compare them with each other as these systems are very similar in their scaffolds. The R_{XB} is defined as the distance between the donor and the acceptor, d_{XB} [Å], and is divided by the sum of vdW radii [Å] of the donor (X) and the acceptor (B) [$R_{XB} = d_{XB}/(X_{vdW} + B_{vdW})$].^{86,125–128} Using 1.55 Å for N and 1.98 Å for I,^{125–128} the complexes **3a** and **3d** gave a value of 0.63, and a value of 0.64 for **3b** and **3c**. A more detailed analysis of the crystal structures revealed possible secondary interactions (see Supporting Information, Figure S3–S10). For **3a** π -stacking is present, with centroid-to-centroid distances of *ca.* 3.71 Å (between the aromatic rings of DMAP and the *N*-iodosaccharin). The secondary interactions are present in each complex and their R_{XB} value variation is negligible, indicating that crystal packing of very strong *N*-






Figure 2. X-ray crystal structures of XB complexes between NISac and (a) DMAP (**3a**), (b) PPY (**3b**), (c) MPY (**3c**), (d) PiPY (**3d**), and corresponding average reduction ratios R_{XB} (thermal ellipsoids shown at 50% probability level; distances are given in Å).




Scheme 2. Synthesis of XB complexes **3** between NISac (**1**) and 4-substituted pyridine derivatives **2** by precipitation. All reactions were performed on a 0.32 mmol scale. NISac (**1**, 100 mg, 0.32 mmol, 1.0 equiv.), pyridine derivative **2** (0.32 mmol, 1.0 equiv.), both dissolved in 1.5 mL EtOAc, then combined. The XB complexes **3a** – **3d** were isolated by filtration.

iodosaccharin-pyridine XB complexes does not have an impact of the N–I···N' halogen bond itself. These observations are in accordance with the work of Fourmigué and Espinosa, who observed the same secondary interactions for a different polymorph of XB complex **3a** as well.⁷⁸ As the XB complexes **3a** – **3d** constituted the foundation of further studies, a robust and scalable synthetic protocol for their preparation was devised. During initial crystallization attempts, colorless precipitates were always found when EtOAc was used as solvent. The further analysis by SCXRD (single crystal X-ray diffraction) identified them as the desired XB complexes **3a** – **3d**. Therefore, the synthesis of these XB complexes was attempted by a precipitation procedure (Scheme 2). First, equimolar amounts of XB donor **1** and the chosen acceptor **2** were dissolved separately in EtOAc, then combined and the desired XB complex precipitated immediately. Through use of this protocol, the desired complexes were obtained in high yields ranging from 81% for **3b** to 91% for **3c**. Being first performed on a scale of few milligrams, the protocol was effortlessly scaled up to a quantity of 100 mg NISac for the complexes **3a**, **3c**, and **3d**. In addition, a 1 mmol batch yielded XB complex **3a** in 87% demonstrating the robustness of the developed protocol. Only in case of PPY, solubility problems might interfere with the standard protocol. A potential water contamination of the pyridine derivative results in protonation and therefore in a lower solubility. In this case, pyridine **2b** was dissolved in DCM, combined with a solution of **1** in EtOAc, and kept overnight for precipitation. If freshly prepared pyridine **2b** was used, the solubility increased, and the standard protocol applied.

Preparation of the symmetric and asymmetric iodine(I) complexes by crystallization

After obtaining sufficient quantities of **3a** – **3d**, the hypothesis if a strongly polarized “entatic” iodine atom allows for an exchange reaction (Scheme 3) was tested. The exchange reaction of the *N*-saccharinato part of the complex **3a** – **3d** with another XB acceptor **2** would lead to a metal-free synthesis of symmetric or asymmetric iodine(I) complexes **4a** – **4cd**, similar to Barluenga’s reagent (Scheme 3).



Symmetric:

4a: R¹ = R² = –NMe₂

4b: R¹ = R² = –N(CH₂)₄

4c: R¹ = R² = –N(CH₂)₄O

4d: R¹ = R² = –N(CH₂)₅

Asymmetric:

4ab: R¹ = –NMe₂, R² = –N(CH₂)₄

4ac: R¹ = –NMe₂, R² = –N(CH₂)₄O

4ad: R¹ = –NMe₂, R² = –N(CH₂)₅

4bc: R¹ = –N(CH₂)₄, R² = –N(CH₂)₄O

4bd: R¹ = –N(CH₂)₄, R² = –N(CH₂)₅

4cd: R¹ = –N(CH₂)₄O, R² = –N(CH₂)₅

Scheme 3. Preparation of symmetric and asymmetric iodine(I) complexes **4a** – **4cd** by an exchange reaction between XB complexes **3** and the corresponding pyridines **2**.

First, as a proof-of-concept, the crystallization of *bis*(4-dimethylaminopyridine)iodine(I) complex **4a** was performed. Equal amounts of DMAP and **3a** were dissolved in different solvents and kept at ambient temperature for co-crystallization by slow evaporation. After a few days, crystals suitable for SCXRD analysis were formed in acetonitrile. The crystal structure confirmed the desired iodine(I) complex **4a** (**4a_1**, Figure 3a) with a symmetric N–I bond lengths, 2.25(1) and 2.25(1) Å, of the [N–I–N]⁺ moiety and the N–I–N angle of 178.0(5)°, very similar to those reported for Barluenga’s reagent (Figure 1b). The *N*-saccharinato anion does not manifest any interactions with the iodine(I) cation. When the same reaction was done in DCM, a slow evaporation of a DCM solution resulted in a different polymorph of **4a**, namely **4a_2**. The geometry of the three-center-four-electron halogen bond in **4a_2** is very similar, with N–I–N angle of 177.67(11)° and N–I bond lengths of 2.248(3) and 2.250(3) Å. Despite the success in the crystallization of the desired complex **4a**, it was realized that co-crystallization under even slightly altered conditions can lead to a different composition.^{115,116} For instance, using DCM as solvent, the desired *bis*(2a)iodine(I) cation was formed. However, the counter anion is not the expected NSac[–]. Instead, the linear triiodide was found resulting from a decomposition of the in-situ formed **4a** (SI, Figure S30–S31). Besides the desired complexes, the used starting materials **2a** (in protonated form) or **3a** tend to crystallize out from the solution, yielding a complex mixture (SI, Figure S23–S29). Motivated by the presence of crystals of symmetric iodine(I) complex **4a** through the co-crystallization method, the crystallization of the symmetric iodine(I) complexes **4b** – **d** was attempted. Unfortunately, the crystallization of complex **4b** from **3b** proved to be difficult. In all attempts, only decomposition products, the **2b**H-saccharinato salt, or the XB complex **3b** was found (SI, Figure S27–S29). Therefore, the crystal structure of **4b** remains unknown. The desired symmetric [2c–I–2c]⁺ complex **4c** was successfully crystallized at –20 °C (Figure 3b) using pyridine **2c** and XB complex **3c** as the starting materials in acetone. The N–I bond lengths of 2.26(1) and 2.26(1) Å, and the N–I–N angle of 179.4(4)° was found for **4c_1** (acetone solvate, SI Figure S17). Two additional structures for **4c** were obtained, **4c_2** (non-solvated, SI Figure S18) and **4c_3** (chloroform solvate, SI Figure S19), with similar XB geometries to **4c_1**, with N–I bond lengths of 2.26(1) Å (**4c_2**) and 2.256(7)/2.251(7) Å (**4c_3**), and N–I–N angles of 180° (**4c_2**) and 179.4(3)° (**4c_3**). A similar behavior was found for complex **4d** (Figure 3c) which crystallized from DCM at 4 °C using pyridine **2d** and XB adduct **3d** as starting




Figure 3. X-ray crystal structures of symmetric iodine(I) complexes **4**: (a) [I(DMAP)₂]⁺ (**4a_1**), (b) [I(MPY)₂]⁺ (**4c_1**), (c) [I(PiPY)₂]⁺ (**4d**) (thermal ellipsoids shown at 50% probability level; anions, minor disordered positions, and solvates omitted for clarity; distances are given in Å).

materials. It shows N–I bond lengths of 2.241(3) and 2.248(4) Å, with a N–I–N angle of 177.7(1)° and crystallizes as a DCM solvate (SI, Figure S21). In the three prepared symmetric complexes of **4c** no secondary interaction on the [N–I–N]⁺ cation was observed. The slight asymmetry in the XB bond length for complexes **4c** and **4d** is very likely caused by crystal packing, especially as two of these compounds crystallized as solvates. The preparation of asymmetric iodine(I) complexes **4ab** – **4cd** was then attempted, knowing about the existence of similar complexes.^{114–116} As the preparation of the symmetric iodine(I) complexes **4a**, **4c**, and **4d** was possible by the exchange reaction between the *N*-saccharinato anion and a pyridine derivative **2**, the same procedure for the preparation of the asymmetric complexes was used. Equal amounts of the chosen XB complex **3** and the pyridine derivative **2** in the crystallization were used. Out of the six possible asymmetric iodine(I) complexes **4ab** – **4cd**, the presence of four complexes (Figure 4) was demonstrated by the analysis of the corresponding crystalline materials that were formed by co-crystallization. Slow cooling at 4 °C of a DCM solution over 15 hours resulted in single crystals of **4ab** (Figure 4a, SI, Figure S14). The asymmetric iodine(I) complex **4ab** exhibits an N–I–N angle of 178.6(3)° and an asymmetric N–I bond lengths. The distance for N(**2a**)–I is 2.258(9) Å and for I–N(**2b**) 2.240(9) Å. This result was expected as the pyrrolidino group is slightly more electron-donating than the dimethylamino group. As a result, the electron density on the sp² nitrogen atom of **2b** is higher, making it stronger XB acceptor. Starting from XB complex **3c** and **2a** or vice versa (XB complex **3a** and pyridine **2c**), the asymmetric complex **4ac** was crystallized from acetone, or MeCN respectively, at 4 °C (Figure 4b). Consequently, successful crystallization cannot be attributed solely to a difference in the relative (already alike) Lewis base character of the pyridines used, since otherwise only the symmetrical complex of the stronger Lewis basic pyridine is to be expected. Therefore, it is reasoned that the association constants plays an important role, too. In a similar way, complex **4bd** was prepared from **3d** and **2b** using DCM as solvent at 4 °C

(Figure 4c). For both iodine(I) complexes, the N–I–N angle of 178.4(3)° (**4ac**) and 177.4(2)° (**4bd**) was found. On the other hand, unexpected symmetric [N–I–N]⁺ moiety was found with N–I bond lengths of 2.234(5) Å and 2.254(5) Å for **4ac** and **4bd**, respectively. It should be noted that both **4ac** and **4bd** demonstrated symmetry-based disorder of **2a/2c** (in **4ac**) and **2b/2d** (in **4bd**), and therefore their individual N–I bond lengths were indistinguishable from one another. The fourth asymmetric iodine(I) complex **4cd** was prepared by crystallization at –20 °C from **3c** and **2d** in chloroform. It showed the expected asymmetric [N–I–N]⁺ moiety with N–I–N angle of 178.9(6)° and N–I lengths of 2.25(1) Å towards **2c** and 2.28(2) Å for **2d**. The implementation of oxygen in the six-membered ring in the morpholino group makes it a slightly stronger electron-donating group than the piperidino group resulting in the distinct difference in bond length towards the iodine. The complexes **4ad** and **4bc** were not obtained upon crystallization in any of the methods attempted, as the starting materials or decomposition products crystallized preferentially. Other products, such as [**2bH**]⁺ and the [**3cH**]⁺saccharinato salts crystallized out in the attempt to prepare complex **4bc**, and the [**2dH**]₃ salt for **4ad**. As undesired products the symmetric iodine(I) complex **4c** was observed in attempts to prepare the asymmetric complex **4cd** from **2c** and **3d**. Similar results were obtained in an attempt to crystallize complex **4bc** from **2b** and **3c**. As asymmetric iodine(I) complexes are also highly reactive and only a few examples had been crystallized,^{114–116} the presence of undesired side products was expected (SI, Figure S23–S32). This is underlined by the fact, that all obtained crystal structures of the complexes **4ab**, **4ac**, **4db** and **4cd** were obtained at lowered temperatures and [2H]saccharinato salts





Figure 4. X-ray crystal structures of asymmetric iodine(I) complexes **4**: (a) [I(DMAP)(PPY)]⁺ (**4ab**), (b) [I(DMAP)(MPY)]⁺ (**4ac**), (c) [I(PPY)(PiPY)]⁺ (**4bd**), (d) [I(MPY)(PiPY)]⁺ (**4cd**) (thermal ellipsoids shown at 50% probability level; anions, minor disordered positions, and solvates omitted for clarity; distances are given in Å).

were detected as side products. Especially the combination of their structural unstableness in solution and the described co-occurrence with side products made further characterization, e.g., by NMR, difficult, and SCXRD remained the only characterization method for the asymmetric complexes.

Mechanochemical synthesis of the NISac XB complexes

The crystallization of the iodine(I) complexes **4** in solution was successful but proved to be difficult due to the observation of numerous side products. Therefore, a solid-state synthetic approach was considered to be a promising alternative. Due to the absence of solvent, the solid-state properties of NISac complexes **3** are unaltered and would allow a direct synthesis of Barluenga-type iodine(I) complexes **4**. When successful, the solid-state synthesis could be used for the preparation of larger quantities of the desired complexes and promote their application as electrophilic iodinating agents, whilst also promoting an environmentally-friendly methodology. For this purpose, a mechanochemical approach using mortar and pestle for triturating the starting materials was adapted.^{129–131} The concept of mechanochemistry is defined as the induction of a chemical reaction by the direct absorption of mechanical energy.¹³² Due to the absence of solvent, shorter reaction times, cleaner reactions,^{129–131,133–139} and access to products that are inaccessible by other means,^{140–147} mechanochemistry enjoys increasing popularity in numerous fields,^{129,134,135,137,139} e.g., organic synthesis,^{130,133,138,140,145} supramolecular chemistry,¹⁴⁸ material sciences,^{141,142,144,146} catalysis,^{136,149} crystal engineering,^{150–156} or coordination chemistry,^{143,157} just to name a few. Due to the high popularity and numerous advantages IUPAC has nominated it as one of ten innovations with the potential to change the world,¹⁵⁸ and has already been successfully applied in the contexts of co-crystallization and halogen bonding.^{148,150–156,159,160}

The co-crystal formation between XB donor **1** and acceptor **2a** as a model reaction was selected (Scheme 4). To avoid any influence of the solvent, a focus was made on *ex-situ* analysis using IR spectroscopy and PXRD (powder X-ray diffraction) for the optimization and compared the spectra to the complexes obtained from solution. As a first attempt, equal amounts of NISac and DMAP were mixed in a vial using a spatula. Just by this gentle mixing the desired XB complex, **3a** was not formed according to IR analysis (SI, Figure S44). As a next try equal amounts of DMAP and NISac were ground in a mortar. Already after 1 minute of grinding the formation of complex **3a** could be detected, together with unreacted starting materials, by IR (SI, Figure S45). A total grinding time of 15 min was necessary to quantitatively convert the starting materials to the desired product according to IR (Figure 5), however, the sample was amorphous and could not be analyzed by PXRD.



Scheme 4. Mechanochemical synthesis of XB complexes **3** from NISac (**1**) and pyridine derivatives **2** using EtOAc for a liquid-assisted grinding (LAG) approach.

A similar behavior was found for the formation of complexes **3b** – **3d**, out of which **3d** required longer grinding time due to the slower conversion reaction (SI, Figure S46–S50). The slower conversion to **3d** compared to the fast reaction in solution was probed by analyzing the XB donor NISac more deeply. The IR spectrum of NISac (SI, Figure S33) showed broad bands in the region from 3600 to 3200 cm^{-1} which could be attributed to OH vibrations. This observation agrees well with a NISac hydrate, which was indeed confirmed by elemental analysis and comparison to the self-prepared sample of **1**. Moreover, a search in the CSD revealed that the reported crystal structures of NISac occur either as a halogen bonded hydrate or a solvate (THF).¹⁰⁰ In addition, attempts to prepare non-hydrated or non-solvated NISac crystals failed. Based on these findings, it was assumed that the hydrate was hampering the reactivity of NISac in the solid state. Therefore, an investigation of a positive influence of small quantities of solvent with hopes to enhance the reactivity of NISac by a liquid-assisted grinding (LAG) was done.^{149,157,161–164} By adding a single drop of EtOAc prior to grinding of NISac and DMAP for 1 min, conversion to the complex **3a** was observed, and IR (Figure 5) indicated its quantitative formation after 5 cycles of adding 1 drop of EtOAc and grinding the mixture for 1 min (Scheme 4). Moreover, the freshly prepared XB complex **3a** was analyzed by PXRD. Even though the mechanochemically prepared **3a** had an amorphous character, the PXRD pattern is in good agreement with a simulated pattern obtained from a SCXRD analysis of solution-prepared pure sample of **3a** (SI, Figure S74). Altering the procedure by adding immediately 3 drops at the beginning and grinding the mixture for 3 min or reducing the amount of solvent by just adding one drop of solvent did not prove to be as efficient (SI, Figure S51–53). Subsequently, the protocol was extended to the preparation of **3b** – **3d** and successfully confirmed their presence by IR and PXRD analysis (SI, Figure S54–57, S75–77). The milling time for **3c** and **3d** had to be extended to 6 min as traces of starting materials could still be detected in the IR spectra after shorter grinding times. The PXRD analysis of the mechanochemically prepared complexes **3a** – **3d** confirmed them to be pure as the perfectly fit with the simulated PXRD pattern based on SCXRD structures of the solution prepared pure **3a** – **3d** (SI, Figure S575–77). The successful mechanochemical preparation of the NISac complexes **3** is further verified by their successful subsequent crystallization and confirmation of their identities from SCXRD studies. In addition to EtOAc as the LAG solvent, MeCN was also tested as an alternative. However, the decomposition reaction




Figure 5. Comparison of IR spectra around 1800-400 cm^{-1} obtained in the mechanochemical synthesis of XB complex **3a**. Black: XB complex **3a** from solution. Red: Neat grinding equal amounts of DMAP and NISac for 15 min. Blue: LAG (EtOAc) equal amounts of DMAP and NISac for 5 min.


was visually observed for **2d** after only 2 min, as the reaction mixture turned from the usual colorless powder to a brown slurry. The decomposition could also be seen by IR analysis (SI, Figure S57).

As a proof-of-concept the developed protocol was transferred to a planetary ball mill, which would simplify up-scaling of the synthesis, making it more reproducible as it would be performed under well-controlled conditions. The experiment was executed for **3a** on a 0.2 mmol scale. An agate vessel, 3 agate balls (10 mm in \varnothing), and 400 rpm frequency were used. After ball-milling of equal amounts of **1** and **2a** in an agate vessel for 5 cycles of 1 min and with addition of one drop of EtOAc prior to each cycle, **3a** was quantitatively obtained according to IR. Even though the solution approach is quite straightforwardly up-scalable too, the benefits of the mechanochemical protocol are substantial as only minute amounts of the LAG solvent is used.

Mechanochemical preparation of the iodine(I) complexes

Having demonstrated the successful utilization of a mechanochemical protocol for the synthesis of NISac complexes **3**, the preparation of symmetric and asymmetric iodine (I) complexes **4** was attempted. The preparation of symmetric iodine(I) complex **4a** was selected as the model target. Equal amounts of DMAP and pure, solution-prepared solid **3a** were ground under neat conditions using an agate mortar. The IR analysis of the product indicated the formation of **4a** after 10 min grinding. The desired complex **4a** could already be detected after 2 min grinding (SI, Figure S59–S60). LAG using EtOAc was then applied to probe the possibility of reducing the reaction time. The LAG experiment consisted of 5 cycles by adding a drop of solvent and grinding equal amounts of starting materials **2a** and **3a** in an agate mortar (Scheme 5, route A). Not only did the reaction quantitatively produce the desired **4a** after 5 min grinding, but it also allowed to use other solvents such as DCM, CHCl_3 , acetone, or MeCN (SI, Figure S61–S62). To further utilize the mechanochemistry in two subsequent reactions, the above LAG protocol was first applied to the synthesis of **3a**, and then continued by adding an equivalent amount of DMAP to the as-obtained **3a** to convert it to the symmetric iodine(I) complex **4a**. The IR and PXRD analysis confirmed (SI, Figure S63–S65, S78–S81) the quantitative conversion (Scheme 5, route B).

This result suggested that **4a** could be obtained mechanochemically through a one-step (concerted or domino-type) reaction, by simply grinding 2 equivalents of DMAP and



Scheme 5. Investigated mechanochemical routes towards iodine(I) complex **4a**.

1 equivalent of NISac using a LAG approach with EtOAc in a mortar (Scheme 5, route C). The IR and PXRD (Figure 6) were very satisfying by revealing the quantitative formation of **4a** in just 5 min. Dissolving **4a**, from the above reaction, in DCM and leaving it to slowly evaporate led to single crystals suitable for SCXRD analysis, which confirmed the identity of **4a**.

After establishing the above, fast and robust, one-step protocol, the preparation of other symmetric iodine(I) complexes **4c–d** was carried out using the two above-mentioned LAG protocols. The first is based on grinding equal amounts of the solution-prepared **3c** and **3d** and the matching pyridines **2c** and **2d** (Scheme 5, route A), whereas the second protocol uses the one-pot synthesis by grinding NISac and 2 equivalents of the chosen pyridines **2c** and **2d** (Scheme 5, route C). For both the **2c-3c** and **2d-4d** reactions, the IR and PXRD analysis revealed the presence of the desired products **4c** and **4d**. As expected, the products were partially amorphous. Nonetheless a good match for the simulated PXRD patterns and those from as-prepared **4c** and **4d** was found (SI, Figure S66–S70, S82–S83). Moreover, the products prepared by two-step or one-step LAG methods show almost equal spectra. The same observations were made for the attempted synthesis of complex **4b**. However, as any crystallization attempt was not successful at this stage and the molecular structure remains unknown so far, the extension of the protocols for the synthesis of product **4b** is still to be confirmed. Only the comparison between the IR spectra of used starting materials and the prepared sample differ, indicating the presence of a new species (SI, Figure S71–S72, S84).

As with the mortar-based mechanochemical preparation of XB complex **3a**, subsequent transfer of the protocol to a planetary ball mill, with a focus on the preparation of symmetric iodine(I) complex **4a** as a proof-of-concept target took place. The same mechanochemical conditions already implemented for the proof-of-concept preparation of **3a** in the planetary ball mill were utilized. Only the molar ratios were changed, with 1 equivalent of NISac and 2 equivalents of DMAP being milled under the aforementioned LAG conditions using EtOAc. The





Figure 6. Comparison of PXRD patterns obtained in the mechanochemical preparation of symmetric iodine(I) complex **4a**.³³ Black: Simulated PXRD pattern of symmetric iodine(I) complex **4a**. Red: route A (**3a** and DMAP for 5 min under LAG (EtOAc) conditions). Blue: route B (two step synthesis under LAG (EtOAc) conditions). Green: route C (grinding one equivalent of NISac and two equivalents of DMAP for 5 min under LAG (EtOAc) conditions).

effective preparation was shown by IR analysis, and it could effortlessly be scaled up to a 1 mmol approach.

Attempts were made to extend the mechanochemical methodology to the synthesis of asymmetric iodine(I) complexes **4ab** – **4cd** by grinding equal amounts of XB complex **3** and another pyridine **2**, however, the systems seemed to undergo scrambling which made their analysis difficult. Thus, the synthesis of asymmetric iodine(I) complexes under mechanochemical conditions was not investigated further.

Iodination of antipyrine

Being able to prepare sufficiently large quantities of the materials **3a** and **4a**, an attempt to use them as surrogates for Barluenga's reagent in proof-of-concept electrophilic iodination reaction was carried out. Literature reports that the iodination of tertiary enamines using Barluenga's reagent is feasible,¹⁶⁵ whereas other methods are described to be difficult.^{165–169} Motivated by these reports, the α -iodination of antipyrine (**5**) to 4-iodoantipyrine (**6**) was selected as the model reaction (Scheme 6).^{165,170} Antipyrine derivatives are known to be biologically active as they exhibit antipyretic, analgesic and anti-inflammatory properties^{165–169} and are used for medicinal screening processes as well.^{171–178} Therefore, iodo-derivative **6** could be a useful building block for other biologically active antipyrine derivatives since the iodination opens a door for further structural diversity.^{165,178} Initially, the reported solution iodination protocol was followed as a baseline reaction.^{165,170} Equimolar amounts of antipyrine and the chosen iodination reagent were stirred in DCM for 1 h at ambient temperature. A parallel mechanochemical protocol was devised, which used a planetary ball mill operating at 400 rpm. The freshly prepared Barluenga's reagent *bis*(pyridine)iodine(I) tetrafluoroborate was used as a benchmark solution reaction (Table 1, entry 1b), which gave **6** in an excellent 91% yield. Using 10 agate balls (5 mm in \varnothing) the same yield (Table 1, entry 1c) was obtained for the mechanochemical protocol, instead using 3 agate balls of 10 mm in diameter yield was increased to 96% (Table 1, entry 1d). Using NISac-based **3a** a very good yield of 87% for both the solution and mechanochemical protocols (Table 1, entry 2) was achieved. The symmetric iodine(I) complex **4a** resulted in a good yield of 80% for solution and 76% in the ball mill, respectively (Table 1, entry 3b and 3c). However, when the mechanochemical reaction was performed on a 1 mmol scale, product **6** was obtained in very good yield of 89% (Table 1, entry 3e). The SCXRD determined unit cell confirmed the product to be **6**.¹⁷⁰ The above results confirm that both the charge-neutral



Scheme 6. Iodination of antipyrine.

Table 1. Results of the iodination of antipyrine using different iodinating reagents.^a

Entry	Reagent	Solution ^b	Ball Mill ^c
1	Barluenga's reagent	91	91 (96) ^d
2	3a	87	87
3	4a	80	76 (89) ^e
4	[(DMAP) ₂ I]BF ₄	85	83
5	NISac	85	93
6	I ₂	89	49

(a) All reactions were performed on a 0.2 mmol scale. **5** (37.6 mg, 0.2 mmol), "I⁺" source (0.2 mmol). Yields after column chromatography. (b) DCM (3 mL), 1 h at RT. (c) Agate vessel, 12 mL, 10 agate balls (5 mm in \varnothing), 400 rpm, 4x15 min, reversing rotation every cycle. (d) 3 balls (10 mm in \varnothing). (e) Performed on a 1 mmol scale.

NISac and cationic iodine(I) complexes can be used for iodination reactions and achieve yields similar to Barluenga's reagent. Even though the yield is slightly lower when made mechanochemically, the advantage in using **3a** and **4a** is clearly based on their very simple and metal-free synthesis compared to preparations that utilize Barluenga's reagent. In addition, 59% of DMAP and 11% of saccharin could be recovered under unoptimized conditions in a 1 mmol approach. This allows saccharin to be recovered and converted back to NISac, which would provide an avenue to a continuous recovery and usage of the starting materials. Consequently, the as prepared mechanochemical materials **3a** and **4a** represent suitable surrogates for Barluenga's reagent. The influence of the ligand modulation of the Barluenga's reagent was studied by using the DMAP derivative of Barluenga's reagent. A good yield of 85% and 83% for both protocols (Table 1, entry 4) was achieved for the solution and mechanochemical strategies, respectively. As these results are similar to the ones using the reagents **3a** and **4a**, it was reasoned that the influence and the nature of the ligand of the cationic iodine(I) complex was negligible. Finally, NISac and elemental iodine were tested (Table 1, entry 5 and 6). In contrast to the literature (*vide supra*) we found the iodination to take place. However, as this puts the use of Barluenga's reagent for the iodination of antipyrine into question, it does not interfere with the fact that the mechanochemically NISac based materials **3a** and **4a** present potential substitutes to Barlueng's reagent. Not only is the performance similar but they also evade the disadvantages of preparing Barluenga's reagent such as chlorinated solvents and heavy metals.

Conclusions

In summary, the current study reports the synthesis of NISac-based XB complexes **3a** – **3d** and the presence of symmetric and asymmetric cationic iodine(I) complexes **4a** – **4d**, both in solution and under mechanochemical conditions. The studied systems were based on NISac (**1**) as a strong halogen bond donor, and electron-rich pyridine derivatives **2** as halogen bond acceptors. Molecular structures of XB complexes **3** manifested very strong halogen bonds. An innovative solution protocol enabled us to prepare symmetric iodine(I) complexes **4** by a substitution of the saccharinato part of the XB complex by a stronger neutral XB acceptor. Seven complexes were successfully identified by exchanging the NSac anion by another pyridine derivative **2**, and were analyzed by SCXRD. However,

the preparation proved to be difficult due to the reactivity of the produced Barluenga-type complexes, manifested by the observation of several decomposition products and ligand scrambling events. As mechanochemistry had not been previously applied in the synthesis of these types of XB complexes, mechanochemical protocols for the synthesis of XB complexes **3** and **4** were developed, with the Liquid Assisted Grinding (LAG) method especially proving to be highly successful. In the preparation of symmetric iodine(I) complexes **4** the LAG method proved to be superior in the synthesis of large quantities of the desired complex **4a**, endearing itself to future commercial uses. Access to large quantities allowed survey **3a** and **4a** as electrophilic iodination reagents in comparison to Barluenga's reagent. Comparable good yields as found with Barluenga's reagent were obtained both in solution and under mechanochemical conditions, such that materials **3a** and **4a** proved to be suitable alternatives to Barluenga's reagent. The developed mechanochemical protocols do not involve heavy metal salts and use only minute amounts of environmentally benign solvents, allowing recovery and reuse of the starting materials. Overall, this makes the mechanochemical methodology developed herein a highly attractive novel strategy for the large-scale synthesis and implementation of iodinating reagents.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

C.S. thanks the Verband der Chemischen Industrie for providing a Kekulé scholarship and the German Academic Exchange Service for a DAAD Scholarship. K.R. and J.S.W. gratefully acknowledge the Academy of Finland (grant numbers: 351121 (KR) and 356187 (JSW)) for funding.

^a University of Jyväskylä, Department of Chemistry, Surfontie 9 B, 40014 Jyväskylä, Finland. Email: kari.t.rissanen@juv.fi

^b Institute of Organic Chemistry, RWTH Aachen University, Landoltweg 1, 52074 Aachen, Germany.

[†] Present address: InnoSyn B.V. Urmonderbaan 22, 6167 RD Geleen, The Netherlands.

[§] Present address: Rigaku Europe SE, Hugenottenallee 167, 63263 Neu-Isenburg, Germany

[‡] These authors contributed equally.

Notes and references

- R. R. Knowles and E. N. Jacobsen, Attractive Noncovalent Interactions in Asymmetric Catalysis: Links between Enzymes and Small Molecule Catalysts, *Proc. Natl. Acad. Sci.*, 2010, **107**, 20678–20685.
- D. Herschlag and M. M. Pinney, Hydrogen Bonds: Simple after All?, *Biochemistry*, 2018, **57**, 3338–3352.
- J. D. Watson and F. H. C. Crick, Molecular Structure of Nucleic Acids: A Structure for Deoxyribose Nucleic Acid, *Nature*, 1953, **171**, 737–738.
- P. Hobza and J. Řezáč, Introduction: Noncovalent Interactions, *Chem. Rev.*, 2016, **116**, 4911–4912.
- I. Alkorta; J. Elguero and A. Frontera, Not Only Hydrogen Bonds: Other Noncovalent Interactions, *Crystals*. 2020, p 180.
- P. Politzer, Book Review of Halogen Bonding: Fundamentals and Applications. Structure and Bonding, 126, *J. Am. Chem. Soc.*, 2008, **130**, 10446–10447.
- H. Wang; W. Wang and W. J. Jin, σ -Hole Bond vs π -Hole Bond: A Comparison Based on Halogen Bond, *Chem. Rev.*, 2016, **116**, 5072–5104.
- L. Vogel; P. Wonner and S. M. Huber, Chalcogen Bonding: An Overview, *Angew. Chem. Int. Ed.*, 2019, **58**, 1880–1891.
- P. Politzer; J. S. Murray and T. Clark, Halogen Bonding and Other σ -Hole Interactions: A Perspective, *Phys. Chem. Chem. Phys.*, 2013, **15**, 11178–11189.
- A. Docker; C. H. Guthrie; H. Kuhn and P. D. Beer, Modulating Chalcogen Bonding and Halogen Bonding Sigma-Hole Donor Atom Potency and Selectivity for Halide Anion Recognition, *Angew. Chem. Int. Ed.*, 2021, **60**, 21973–21978.
- G. Cavallo; P. Metrangolo; R. Milani; T. Pilati; A. Priimagi; G. Resnati and G. Terraneo, The Halogen Bond, *Chem. Rev.*, 2016, **116**, 2478–2601.
- R. S. Mulliken, Structures of Complexes Formed by Halogen Molecules with Aromatic and with Oxygenated Solvents¹, *J. Am. Chem. Soc.*, 1950, **72**, 600–608.
- R. S. Mulliken, Molecular Compounds and Their Spectra. II, *J. Am. Chem. Soc.*, 1952, **74**, 811–824.
- R. S. Mulliken, Molecular Compounds and Their Spectra. III. The Interaction of Electron Donors and Acceptors, *J. Phys. Chem.*, 1952, **56**, 801–822.
- J. R. Platt, 1966 Nobel Laureate in Chemistry: Robert S. Mulliken, *Science (80-)*, 1966, **154**, 745–747.
- O. Hassel, Structural Aspects of Interatomic Charge-Transfer Bonding, *Science (80-)*, 1970, **170**, 497–502.
- O. Hassel and J. Hvoslef, The Structure of Bromine 1,4-Dioxanate, *Acta Chem. Scand.*, 1954, **8**, 873–873.
- O. Hassel and C. Rømming, Direct Structural Evidence for Weak Charge-Transfer Bonds in Solids Containing Chemically Saturated Molecules, *Q. Rev. Chem. Soc.*, 1962, **16**, 1–18.
- O. Hassel and K. O. Strømme, Structure of the Crystalline Compound Benzene-Bromine (1:1), *Acta Chem. Scand.*, 1958, **12**, 1146.
- O. Hassel and K. O. Strømme, Crystal Structure of the Addition Compound Benzene-Chlorine (1:1), *Acta Chem. Scand.*, 1959, **13**, 1781–1786.
- A. De Santis; A. Forni; R. Liantonio; P. Metrangolo; T. Pilati and G. Resnati, N \cdots Br Halogen Bonding: One-Dimensional Infinite Chains through the Self-Assembly of Dibromotetrafluorobenzenes with Dipyrindyl Derivatives, *Chem. – A Eur. J.*, 2003, **9**, 3974–3983.
- L. C. Gilday; S. W. Robinson; T. A. Barendt; M. J. Langton; B. R. Mullaney and P. D. Beer, Halogen Bonding in Supramolecular Chemistry, *Chem. Rev.*, 2015, **115**, 7118–

- 7195.
- (23) L. C. Gilday and P. D. Beer, Halogen- and Hydrogen-Bonding Catenanes for Halide-Anion Recognition, *Chem. – A Eur. J.*, 2014, **20**, 8379–8385.
- (24) O. Dumele; N. Trapp and F. Diederich, Halogen Bonding Molecular Capsules, *Angew. Chem. Int. Ed.*, 2015, **54**, 12339–12344.
- (25) N. K. Beyeh; F. Pan and K. Rissanen, A Halogen-Bonded Dimeric Resorcinarene Capsule, *Angew. Chem. Int. Ed.*, 2015, **54**, 7303–7307.
- (26) C.-Z. Liu; S. Koppireddi; H. Wang; D.-W. Zhang and Z.-T. Li, Halogen Bonding-Driven Formation of Supramolecular Macrocycles and Double Helix, *Chinese Chem. Lett.*, 2019, **30**, 953–956.
- (27) A. Mukherjee; S. Tothadi and G. R. Desiraju, Halogen Bonds in Crystal Engineering: Like Hydrogen Bonds yet Different, *Acc. Chem. Res.*, 2014, **47**, 2514–2524.
- (28) J. Teyssandier; K. S. Mali and S. De Feyter, Halogen Bonding in Two-Dimensional Crystal Engineering, *ChemistryOpen*, 2020, **9**, 225–241.
- (29) R. B. Walsh; C. W. Padgett; P. Metrangolo; G. Resnati; T. W. Hanks and W. T. Pennington, Crystal Engineering through Halogen Bonding: Complexes of Nitrogen Heterocycles with Organic Iodides, *Cryst. Growth Des.*, 2001, **1**, 165–175.
- (30) G. Mínguez Espallargas; F. Zordan; L. Arroyo Marín; H. Adams; K. Shankland; J. van de Streek and L. Brammer, Rational Modification of the Hierarchy of Intermolecular Interactions in Molecular Crystal Structures by Using Tunable Halogen Bonds, *Chem. – A Eur. J.*, 2009, **15**, 7554–7568.
- (31) H. L. Nguyen; P. N. Horton; M. B. Hursthouse; A. C. Legon and D. W. Bruce, Halogen Bonding: A New Interaction for Liquid Crystal Formation, *J. Am. Chem. Soc.*, 2004, **126**, 16–17.
- (32) M. Saccone; M. Spengler; M. Pflötscher; K. Kuntze; M. Virkki; C. Wölper; R. Gehrke; G. Jansen; P. Metrangolo; A. Priimagi and M. Giese, Photoresponsive Halogen-Bonded Liquid Crystals: The Role of Aromatic Fluorine Substitution, *Chem. Mater.*, 2019, **31**, 462–470.
- (33) D. Devadiga and T. N. Ahipa, An Up-to-Date Review on Halogen-Bonded Liquid Crystals, *J. Mol. Liq.*, 2021, **333**, 115961.
- (34) F. Fernandez-Palacio; M. Poutanen; M. Saccone; A. Siiskonen; G. Terraneo; G. Resnati; O. Ikkala; P. Metrangolo and A. Priimagi, Efficient Light-Induced Phase Transitions in Halogen-Bonded Liquid Crystals, *Chem. Mater.*, 2016, **28**, 8314–8321.
- (35) G. Cavallo; G. Terraneo; A. Monfredini; M. Saccone; A. Priimagi; T. Pilati; G. Resnati; P. Metrangolo and D. W. Bruce, Superfluorinated Ionic Liquid Crystals Based on Supramolecular, Halogen-Bonded Anions, *Angew. Chem. Int. Ed.*, 2016, **55**, 6300–6304.
- (36) R. Kampes; S. Zechel; M. D. Hager and U. S. Schubert, Halogen Bonding in Polymer Science: Towards New Smart Materials, *Chem. Sci.*, 2021, **12**, 9275–9286.
- (37) A. Priimagi; G. Cavallo; A. Forni; M. Gorynsztejn–Leben; M. Kaivola; P. Metrangolo; R. Milani; A. Shishido; T. Pilati; G. Resnati and G. Terraneo, Halogen Bonding versus Hydrogen Bonding in Driving Self-Assembly and Performance of Light-Responsive Supramolecular Polymers, *Adv. Funct. Mater.*, 2012, **22**, 2572–2579.
- (38) A. Vanderkooy and M. S. Taylor, Solution-Phase Self-Assembly of Complementary Halogen Bonding Polymers, *J. Am. Chem. Soc.*, 2015, **137**, 5080–5086.
- (39) G. Berger; J. Soubhye and F. Meyer, Halogen Bonding in Polymer Science: From Crystal Engineering to Functional Supramolecular Polymers and Materials, *Polym. Chem.*, 2015, **6**, 3559–3580.
- (40) L. Meazza; J. A. Foster; K. Fücke; P. Metrangolo; G. Resnati and J. W. Steed, Halogen-Bonding-Triggered Supramolecular Gel Formation, *Nat. Chem.*, 2013, **5**, 42–47.
- (41) G. Berger; P. Frangville and F. Meyer, Halogen Bonding for Molecular Recognition: New Developments in Materials and Biological Sciences, *Chem. Commun.*, 2020, **56**, 4970–4981.
- (42) A. Pizzi; L. Lascialfari; N. Demitri; A. Bertolani; D. Maiolo; E. Carretti and P. Metrangolo, Halogen Bonding Modulates Hydrogel Formation from Fmoc Amino Acids, *CrystEngComm*, 2017, **19**, 1870–1874.
- (43) S. Bhattacharjee and S. Bhattacharya, Remarkable Role of C–I⋯N Halogen Bonding in Thixotropic ‘Halo’Gel Formation, *Langmuir*, 2016, **32**, 4270–4277.
- (44) X. Tong; X. Zhao; Y. Qiu; H. Wang; Y. Liao and X. Xie, Intrinsically Visible Light-Responsive Liquid Crystalline Physical Gels Driven by a Halogen Bond, *Langmuir*, 2020, **36**, 11873–11879.
- (45) R. Wilcken; X. Liu; M. O. Zimmermann; T. J. Rutherford; A. R. Fersht; A. C. Joerger and F. M. Boeckler, Halogen-Enriched Fragment Libraries as Leads for Drug Rescue of Mutant P53, *J. Am. Chem. Soc.*, 2012, **134**, 6810–6818.
- (46) M. R. Scholfield; C. M. Vander Zanden; M. Carter and P. S. Ho, Halogen Bonding (X-Bonding): A Biological Perspective, *Protein Sci.*, 2013, **22**, 139–152.
- (47) R. Wilcken; M. O. Zimmermann; A. Lange; A. C. Joerger and F. M. Boeckler, Principles and Applications of Halogen Bonding in Medicinal Chemistry and Chemical Biology, *J. Med. Chem.*, 2013, **56**, 1363–1388.
- (48) L. A. Hardegger; B. Kuhn; B. Spinnler; L. Anselm; R. Ecabert; M. Stihle; B. Gsell; R. Thoma; J. Diez; J. Benz; J.-M. Plancher; G. Hartmann; D. W. Banner; W. Haap and F. Diederich, Systematic Investigation of Halogen Bonding in Protein–Ligand Interactions, *Angew. Chem. Int. Ed.*, 2011, **50**, 314–318.
- (49) J. Heidrich; L. E. Sperl and F. M. Boeckler, Embracing the Diversity of Halogen Bonding Motifs in Fragment-Based Drug Discovery—Construction of a Diversity-Optimized Halogen-Enriched Fragment Library, *Frontiers in Chemistry*. 2019, pp 1–9. <https://www.frontiersin.org/articles/10.3389/fchem.2019.00009>.
- (50) A. P. Bruckmann Miguel A; Bolm, Carsten, Organocatalysis through Halogen-Bond Activation, *Synlett*, 2008, **2008**, 900–902.
- (51) R. L. Sutar and S. M. Huber, Catalysis of Organic Reactions through Halogen Bonding, *ACS Catal.*, 2019, **9**, 9622–9639.

- (52) A. Dreger; P. Wonner; E. Engelage; S. M. Walter; R. Stoll and S. M. Huber, A Halogen-Bonding-Catalysed Nazarov Cyclisation Reaction, *Chem. Commun.*, 2019, **55**, 8262–8265.
- (53) Y. T. Kobayashi Yoshiji, Reactions Catalyzed by 2-Halogenated Azolium Salts: From Halogen-Bond Donors to Brønsted-Acidic Salts, *Synlett*, 2020, **31**, 772–783.
- (54) D. Bulfield and S. M. Huber, Halogen Bonding in Organic Synthesis and Organocatalysis, *Chem. – A Eur. J.*, 2016, **22**, 14434–14450.
- (55) S. Kuwano; T. Suzuki; Y. Hosaka and T. Arai, A Chiral Organic Base Catalyst with Halogen-Bonding-Donor Functionality: Asymmetric Mannich Reactions of Malononitrile with N-Boc Aldimines and Ketimines, *Chem. Commun.*, 2018, **54**, 3847–3850.
- (56) S. H. Jungbauer and S. M. Huber, Cationic Multidentate Halogen-Bond Donors in Halide Abstraction Organocatalysis: Catalyst Optimization by Preorganization, *J. Am. Chem. Soc.*, 2015, **137**, 12110–12120.
- (57) G. R. Desiraju; P. S. Ho; L. Kloo; A. C. Legon; R. Marquardt; P. Metrangolo; P. Politzer; G. Resnati and K. Rissanen, Definition of the Halogen Bond (IUPAC Recommendations 2013), *Pure Appl. Chem.*, 2013, **85**, 1711–1713.
- (58) P. Politzer and J. S. Murray, Halogen Bonding: An Interim Discussion, *ChemPhysChem*, 2013, **14**, 278–294.
- (59) F. F. Awwadi; R. D. Willett; K. A. Peterson and B. Twamley, The Nature of Halogen···Halogen Synthons: Crystallographic and Theoretical Studies, *Chem. – A Eur. J.*, 2006, **12**, 8952–8960.
- (60) K. E. Riley; J. S. Murray; P. Politzer; M. C. Concha and P. Hobza, Br···O Complexes as Probes of Factors Affecting Halogen Bonding: Interactions of Bromobenzenes and Bromopyrimidines with Acetone, *J. Chem. Theory Comput.*, 2009, **5**, 155–163.
- (61) T. Clark, σ -Holes, *WIREs Comput. Mol. Sci.*, 2013, **3**, 13–20.
- (62) K. E. Riley and P. Hobza, Investigations into the Nature of Halogen Bonding Including Symmetry Adapted Perturbation Theory Analyses, *J. Chem. Theory Comput.*, 2008, **4**, 232–242.
- (63) K. E. Riley; J. S. Murray; J. Fanfrlík; J. Řezáč; R. J. Solá; M. C. Concha; F. M. Ramos and P. Politzer, Halogen Bond Tunability II: The Varying Roles of Electrostatic and Dispersion Contributions to Attraction in Halogen Bonds, *J. Mol. Model.*, 2013, **19**, 4651–4659.
- (64) H. A. Bent, Structural Chemistry of Donor-Acceptor Interactions, *Chem. Rev.*, 1968, **68**, 587–648.
- (65) A. E. Reed; F. Weinhold; R. Weiss and J. Macheleid, Nature of the Contact Ion Pair Trichloromethyl-Chloride (CCl₃+Cl⁻). A Theoretical Study, *J. Phys. Chem.*, 1985, **89**, 2688–2694.
- (66) J. Řezáč and A. de la Lande, On the Role of Charge Transfer in Halogen Bonding, *Phys. Chem. Chem. Phys.*, 2017, **19**, 791–803.
- (67) B. Inscoc; H. Rathnayake and Y. Mo, Role of Charge Transfer in Halogen Bonding, *J. Phys. Chem. A*, 2021, **125**, 2944–2953.
- (68) C. Wang; D. Danovich; Y. Mo and S. Shaik, On The Nature of the Halogen Bond, *J. Chem. Theory Comput.*, 2014, **10**, 3726–3737.
- (69) P. Politzer; J. S. Murray and M. C. Concha, Halogen Bonding and the Design of New Materials: Organic Bromides, Chlorides and Perhaps Even Fluorides as Donors, *J. Mol. Model.*, 2007, **13**, 643–650.
- (70) P. Politzer; P. Lane; M. C. Concha; Y. Ma and J. S. Murray, An Overview of Halogen Bonding, *J. Mol. Model.*, 2007, **13**, 305–311.
- (71) P. Politzer; J. S. Murray and T. Clark, Halogen Bonding: An Electrostatically-Driven Highly Directional Noncovalent Interaction, *Phys. Chem. Chem. Phys.*, 2010, **12**, 7748–7757.
- (72) J. M. Holthoff; E. Engelage; R. Weiss and S. M. Huber, “Anti-Electrostatic” Halogen Bonding, *Angew. Chem. Int. Ed.*, 2020, **59**, 11150–11157.
- (73) T. Brinck and A. N. Borrfors, Electrostatics and Polarization Determine the Strength of the Halogen Bond: A Red Card for Charge Transfer, *J. Mol. Model.*, 2019, **25**, 125.
- (74) F. Otte; J. Kleinheider; W. Hiller; R. Wang; U. Englert and C. Strohmann, Weak yet Decisive: Molecular Halogen Bond and Competing Weak Interactions of Iodobenzene and Quinuclidine, *J. Am. Chem. Soc.*, 2021, **143**, 4133–4137.
- (75) O. Dumele; D. Wu; N. Trapp; N. Goroff and F. Diederich, Halogen Bonding of (Iodoethynyl)Benzene Derivatives in Solution, *Org. Lett.*, 2014, **16**, 4722–4725.
- (76) R. Puttreddy; J. M. Rautiainen; T. Mäkelä and K. Rissanen, Strong N–X···O–N Halogen Bonds: A Comprehensive Study on N-Halosaccharin Pyridine N-Oxide Complexes, *Angew. Chem. Int. Ed.*, 2019, **58**, 18610–18618.
- (77) K. Rissanen, Halogen Bonded Supramolecular Complexes and Networks, *CrystEngComm*, 2008, **10**, 1107–1113.
- (78) O. Makhotkina; J. Liefbrig; O. Jeannin; M. Fourmigué; E. Aubert and E. Espinosa, Cocrystal or Salt: Solid State-Controlled Iodine Shift in Crystalline Halogen-Bonded Systems, *Cryst. Growth Des.*, 2015, **15**, 3464–3473.
- (79) C. Ouvrard; J.-Y. Le Questel; M. Berthelot and C. Laurence, Halogen-Bond Geometry: A Crystallographic Database Investigation of Dihalogen Complexes, *Acta Crystallogr. Sect. B*, 2003, **59**, 512–526.
- (80) M. C. Aragoni; M. Arca; F. A. Devillanova; F. Isaia; V. Lippolis; A. Mancini; L. Pala; A. M. Z. Slawin and J. D. Woollins, First Example of an Infinite Polybromide 2D-Network, *Chem. Commun.*, 2003, No. 17, 2226–2227.
- (81) A. S. Batsanov; J. A. K. Howard; A. P. Lightfoot; S. J. R. Twiddle and A. Whiting, Stereoselective Chloro-Deboronation Reactions Induced by Substituted Pyridine–Iodine Chloride Complexes, *Eur. J. Org. Chem.*, 2005, **2005**, 1876–1883.
- (82) C. Rømming, Refinement of the Crystal Structure of the Charge Transfer Compound Pyridine-Iodomono-chloride, *Acta Chem. Scand.*, 1972, **26**, 1555–1560.
- (83) M. C. Aragoni; M. Arca; S. J. Coles; F. A. Devillanova; M. B. Hursthouse; S. L. Coles (née Huth); F. Isaia; V. Lippolis and A. Mancini, CT-Adduct vs. Pyridinium Polyhalide Salt Formation in the Reactions between Polypyridyl Donors and Dihalogens: Reactivity of 1,4-Di-(3'-Pyridylethynyl)Benzene towards Br₂ and I₂, *CrystEngComm*, 2011, **13**, 6319–6322.
- (84) J.-Y. Le Questel; C. Laurence and J. Graton, Halogen-Bond

- Interactions: A Crystallographic Basicity Scale towards Iodoorganic Compounds, *CrystEngComm*, 2013, **15**, 3212–3221.
- (85) G. C. Pimentel, The Bonding of Trihalide and Bifluoride Ions by the Molecular Orbital Method, *J. Chem. Phys.*, 1951, **19**, 446–448.
- (86) L. Koskinen; P. Hirva; E. Kalenius; S. Jääskeläinen; K. Rissanen and M. Haukka, Halogen Bonds with Coordinative Nature: Halogen Bonding in a S–I⁺–S Iodonium Complex, *CrystEngComm*, 2015, **17**, 1231–1236.
- (87) M. Bedin; A. Karim; M. Reitti; A.-C. C. Carlsson; F. Topić; M. Cetina; F. Pan; V. Havel; F. Al-Ameri; V. Sindelar; K. Rissanen; J. Gräfenstein and M. Erdélyi, Counterion Influence on the N–I–N Halogen Bond, *Chem. Sci.*, 2015, **6**, 3746–3756.
- (88) S. B. Hakkert and M. Erdélyi, Halogen Bond Symmetry: The N–X–N Bond, *J. Phys. Org. Chem.*, 2015, **28**, 226–233.
- (89) L. Turunen and M. Erdélyi, Halogen Bonds of Halonium Ions, *Chem. Soc. Rev.*, 2020, **49**, 2688–2700.
- (90) J. S. Ward; K.-N. Truong; M. Erdélyi and K. Rissanen, 1.12 - Halogen-Bonded Halogen(I) Ion Complexes., In *Comprehensive Inorganic Chemistry III*; Reedijk, J., Poeppelemeier, K. R. B. T.-C. I. C. I. I. I. (Third E., Eds.; Elsevier: Oxford, 2023; pp 586–601.
- (91) A.-C. C. Carlsson; K. Mehmeti; M. Uhrbom; A. Karim; M. Bedin; R. Puttreddy; R. Kleinmaier; A. A. Neverov; B. Nekoueishahraki; J. Gräfenstein; K. Rissanen and M. Erdélyi, Substituent Effects on the [N–I–N] + Halogen Bond, *J. Am. Chem. Soc.*, 2016, **138**, 9853–9863.
- (92) L. P. Wolters and F. M. Bickelhaupt, Halogen Bonding versus Hydrogen Bonding: A Molecular Orbital Perspective, *ChemistryOpen*, 2012, **1**, 96–105.
- (93) F. Jiménez-Grávalos; M. Gallegos; Á. Martín Pendás and A. S. Novikov, Challenging the Electrostatic σ -Hole Picture of Halogen Bonding Using Minimal Models and the Interacting Quantum Atoms Approach, *J. Comput. Chem.*, 2021, **42**, 676–687.
- (94) J. Barluenga; J. M. González; P. J. Campos and G. Asensio, I(Py)₂BF₄, a New Reagent in Organic Synthesis: General Method for the 1,2-Iodofunctionalization of Olefins, *Angew. Chem. Int. Ed.*, 1985, **24**, 319–320.
- (95) J. Barluenga; H. Vázquez-Villa; A. Ballesteros and J. M. González, Cyclization of Carbonyl Groups onto Alkynes upon Reaction with IPy₂BF₄ and Their Trapping with Nucleophiles: A Versatile Trigger for Assembling Oxygen Heterocycles, *J. Am. Chem. Soc.*, 2003, **125**, 9028–9029.
- (96) J. Barluenga; F. González-Bobes; M. C. Murguía; S. R. Ananthoju and J. M. González, Bis(Pyridine)Iodonium Tetrafluoroborate (IPy₂BF₄): A Versatile Oxidizing Reagent, *Chem. – A Eur. J.*, 2004, **10**, 4206–4213.
- (97) J. Barluenga, Transferring Iodine: More than a Simple Functional Group Exchange in Organic Synthesis, *Pure Appl. Chem.*, 1999, **71**, 431–436.
- (98) Y. Kim; E. J. Mckinley; K. E. Christensen; N. H. Rees and A. L. Thompson, Toward the Understanding of Modulation in Molecular Materials: Barluenga's Reagent and Its Analogues., *Cryst. Growth Des.*, 2014, **14**, 6294–6301.
- (99) C. Alvarez-Rua; S. Garcia-Granda; A. Ballesteros; F. Gonzalez-Bobes and J. M. Gonzalez, Bis(Pyridine)Iodonium(I) Tetrafluoroborate, *Acta Crystallogr. Sect. E*, 2002, **58**, o1381–o1383.
- (100) C. R. Groom; I. J. Bruno; M. P. Lightfoot and S. C. Ward, The Cambridge Structural Database, *Acta Crystallogr. Sect. B*, 2016, **72**, 171–179.
- (101) C. F. Macrae; I. J. Bruno; J. A. Chisholm; P. R. Edgington; P. McCabe; E. Pidcock; L. Rodriguez-Monge; R. Taylor; J. van de Streek and P. A. Wood, Mercury CSD 2.0 - New Features for the Visualization and Investigation of Crystal Structures, *J. Appl. Crystallogr.*, 2008, **41**, 466–470.
- (102) T. Okitsu; S. Yumitate; K. Sato; Y. In and A. Wada, Substituent Effect of Bis(Pyridines)Iodonium Complexes as Iodinating Reagents: Control of the Iodocyclization/Oxidation Process, *Chem. – A Eur. J.*, 2013, **19**, 4992–4996.
- (103) R. D. Evans and J. H. Schauble, Syntheses of α -Iodocarbonyl Compounds Using Bis(Sym-Collidine)Iodine(I) Tetrafluoroborate/Dimethyl Sulfoxide, *Synth.*, 1986, **1986**, 727–730.
- (104) J. C. López; P. Bernal-Albert; C. Uriel; S. Valverde and A. M. Gómez, IPy₂BF₄/HF-Pyridine: A New Combination of Reagents for the Transformation of Partially Unprotected Thioglycosides and n-Pentenyl Glycosides to Glycosyl Fluorides, *J. Org. Chem.*, 2007, **72**, 10268–10271.
- (105) J. C. López; C. Uriel; A. Guillamón-Martín; S. Valverde and A. M. Gómez, IPy₂BF₄-Mediated Transformation of n-Pentenyl Glycosides to Glycosyl Fluorides: A New Pair of Semiorthogonal Glycosyl Donors, *Org. Lett.*, 2007, **9**, 2759–2762.
- (106) J. W. Lown and A. V Joshua, Stereochemistry and Regiochemistry of the Addition of Iodonium Nitrate to Alkenes, *J. Chem. Soc. Perkin Trans. 1*, 1973, No. 0, 2680–2687.
- (107) J. W. Lown and A. V Joshua, Electrophilic Additions of Iodonium Nitrate to Unsaturated Substrates, *Can. J. Chem.*, 1977, **55**, 122–130.
- (108) J. W. Lown and A. V Joshua, Electrophilic Additions of Bromonium Nitrate to Unsaturated Substrates, *Can. J. Chem.*, 1977, **55**, 508–521.
- (109) M. Bigioni; P. Ganis; A. Panunzi; F. Ruffo; C. Salvatore and A. Vito, Electrophilic Attack of [I(Py)₂]⁺(NO₃⁻) on Three-Coordinate Pt(0) Precursors: Synthesis and in Vitro Antitumor Activity of Water-Soluble, Five-Coordinate Pt(II) Complexes, *Eur. J. Inorg. Chem.*, 2000, **2000**, 1717–1721.
- (110) M. Vilaró; J. Nieto; J. R. La Parra; M. R. Almeida; A. Ballesteros; A. Planas; G. Arsequell and G. Valencia, Tuning Transthyretin Amyloidosis Inhibition Properties of Iododifluoride by Combinatorial Engineering of the Nonsalicylic Ring Substitutions, *ACS Comb. Sci.*, 2015, **17**, 32–38.
- (111) A.-C. C. Carlsson; J. Gräfenstein; A. Budnjo; J. L. Laurila; J. Bergquist; A. Karim; R. Kleinmaier; U. Brath and M. Erdélyi, Symmetric Halogen Bonding Is Preferred in Solution, *J. Am. Chem. Soc.*, 2012, **134**, 5706–5715.
- (112) D. C. Georgiou; P. Butler; E. C. Browne; D. J. D. Wilson and J. L. Dutton, On the Bonding in Bis-Pyridine Iodonium Cations*, *Aust. J. Chem.*, 2013, **66**, 1179–1188.

- (113) S. Lindblad; K. Mehmeti; A. X. Veiga; B. Nekoueshahraki; J. Gräfenstein and M. Erdélyi, Halogen Bond Asymmetry in Solution, *J. Am. Chem. Soc.*, 2018, **140**, 13503–13513.
- (114) J. S. Ward; G. Fiorini; A. Frontera and K. Rissanen, Asymmetric [N–I–N]⁺ Halonium Complexes, *Chem. Commun.*, 2020, **56**, 8428–8431.
- (115) D. von der Heiden; K. Rissanen and M. Erdélyi, Asymmetric [N–I–N]⁺ Halonium Complexes in Solution?, *Chem. Commun.*, 2020, **56**, 14431–14434.
- (116) S. Yu and J. S. Ward, Ligand Exchange among Iodine(I) Complexes, *Dalton Trans.*, 2022, **51**, 4668–4674.
- (117) J. M. Chalker; A. L. Thompson; B. G. Davis; N. Zaware and P. Wipf, Safe and Scalable Preparation of Barluenga's Reagent: (Bis(Pyridine) Iodonium(I) Tetrafluoroborate), *Organic Syntheses*. 2010, pp 288–298. <https://www.scopus.com/inward/record.uri?eid=2-s2.0-79952547604&doi=10.15227%2Forgsyn.087.0288&partnerID=40&md5=cc28947db137eecd7fd469341c28a9f2>.
- (118) J. S. Ward; A. Frontera and K. Rissanen, Iodonium Complexes of the Tertiary Amines Quinuclidine and 1-Ethylpiperidine, *Dalton Trans.*, 2021, **50**, 8297–8301.
- (119) J. S. Ward; R. M. Gomila; A. Frontera and K. Rissanen, Iodine(I) Complexes Incorporating Sterically Bulky 2-Substituted Pyridines, *RSC Adv.*, 2022, **12**, 8674–8682.
- (120) J. S. Ward, The Solid-State Hierarchy and Iodination Potential of [Bis(3-Acetaminopyridine)Iodine(I)]PF₆, *CrystEngComm*, 2022, **24**, 7029–7033.
- (121) K. Rissanen and J. S. Ward, Iodine(I) and Silver(I) Complexes Incorporating 3-Substituted Pyridines, *ACS Omega*, 2023, **8**, 24064–24071.
- (122) J. S. Ward; E. I. Sievänen and K. Rissanen, Solid-State NMR Spectroscopy of Iodine(I) Complexes, *Chem. – An Asian J.*, 2023, **18**, e202201203.
- (123) C. Wang; D. Danovich; H. Chen and S. Shaik, Oriented External Electric Fields: Tweezers and Catalysts for Reactivity in Halogen-Bond Complexes, *J. Am. Chem. Soc.*, 2019, **141**, 7122–7136.
- (124) M. Mattila; K. Rissanen and J. S. Ward, Chiral Carbonyl Hypiodites, *Chem. Commun.*, 2023, **59**, 4648–4651.
- (125) A. Bondi, Van Der Waals Volumes and Radii, *J. Phys. Chem.*, 1964, **68**, 441–451.
- (126) J. P. M. Lommerse; A. J. Stone; R. Taylor and F. H. Allen, The Nature and Geometry of Intermolecular Interactions between Halogens and Oxygen or Nitrogen, *J. Am. Chem. Soc.*, 1996, **118**, 3108–3116.
- (127) L. Brammer; E. A. Bruton and P. Sherwood, Understanding the Behavior of Halogens as Hydrogen Bond Acceptors, *Cryst. Growth Des.*, 2001, **1**, 277–290.
- (128) F. Zordan; L. Brammer and P. Sherwood, Supramolecular Chemistry of Halogens: Complementary Features of Inorganic (M–X) and Organic (C–X') Halogens Applied to M–X···X'–C Halogen Bond Formation, *J. Am. Chem. Soc.*, 2005, **127**, 5979–5989.
- (129) S. L. James and T. Friščić, Mechanochemistry, *Chem. Soc. Rev.*, 2013, **42**, 7494–7496.
- (130) C. Oliver Kappe; J. Mack and C. Bolm, Enabling Techniques for Organic Synthesis, *J. Org. Chem.*, 2021, **86**, 14242–14244.
- (131) C. Bolm and J. G. Hernández, Mechanochemistry of Gaseous Reactants, *Angew. Chem. Int. Ed.*, 2019, **58**, 3285–3299.
- (132) K. Horie; M. Barón; R. B. Fox; J. He; M. Hess; J. Kahovec; T. Kitayama; P. Kubisa; E. Maréchal; W. Mormann; R. F. T. Stepto; D. Tabak; J. Vohlídal; E. S. Wilks and W. J. Work, Definitions of Terms Relating to Reactions of Polymers and to Functional Polymeric Materials (IUPAC Recommendations 2003), 2004, **76**, 889–906.
- (133) T. Friščić; C. Mottillo and H. M. Titi, Mechanochemistry for Synthesis, *Angew. Chem. Int. Ed.*, 2020, **59**, 1018–1029.
- (134) S. L. James; C. J. Adams; C. Bolm; D. Braga; P. Collier; T. Friščić; F. Grepioni; K. D. M. Harris; G. Hyett; W. Jones; A. Krebs; J. Mack; L. Maini; A. G. Orpen; I. P. Parkin; W. C. Shearouse; J. W. Steed and D. C. Waddell, Mechanochemistry: Opportunities for New and Cleaner Synthesis, *Chem. Soc. Rev.*, 2012, **41**, 413–447.
- (135) J. L. Howard; Q. Cao and D. L. Browne, Mechanochemistry as an Emerging Tool for Molecular Synthesis: What Can It Offer?, *Chem. Sci.*, 2018, **9**, 3080–3094.
- (136) W. Pickhardt; S. Grätz and L. Borchardt, Direct Mechanocatalysis: Using Milling Balls as Catalysts, *Chem. – A Eur. J.*, 2020, **26**, 12903–12911.
- (137) K. J. Ardila-Fierro and J. G. Hernández, Sustainability Assessment of Mechanochemistry by Using the Twelve Principles of Green Chemistry, *ChemSusChem*, 2021, **14**, 2145–2162.
- (138) C. G. Avila-Ortiz and E. Juaristi, Novel Methodologies for Chemical Activation in Organic Synthesis under Solvent-Free Reaction Conditions, *Molecules*. 2020, pp 1–25.
- (139) J.-L. Do and T. Friščić, Mechanochemistry: A Force of Synthesis, *ACS Cent. Sci.*, 2017, **3**, 13–19.
- (140) L. Konnert; B. Reneaud; R. M. de Figueiredo; J.-M. Campagne; F. Lamaty; J. Martinez and E. Colacino, Mechanochemical Preparation of Hydantoins from Amine Esters: Application to the Synthesis of the Antiepileptic Drug Phenytoin, *J. Org. Chem.*, 2014, **79**, 10132–10142.
- (141) G.-W. Wang; K. Komatsu; Y. Murata and M. Shiro, Synthesis and X-Ray Structure of Dumb-Bell-Shaped C₁₂₀, *Nature*, 1997, **387**, 583–586.
- (142) K. Komatsu; G.-W. Wang; Y. Murata; T. Tanaka; K. Fujiwara; K. Yamamoto and M. Saunders, Mechanochemical Synthesis and Characterization of the Fullerene Dimer C₁₂₀, *J. Org. Chem.*, 1998, **63**, 9358–9366.
- (143) J. G. Hernández and C. Bolm, Altering Product Selectivity by Mechanochemistry, *J. Org. Chem.*, 2017, **82**, 4007–4019.
- (144) S.-E. Zhu; F. Li and G.-W. Wang, Mechanochemistry of Fullerenes and Related Materials, *Chem. Soc. Rev.*, 2013, **42**, 7535–7570.
- (145) G.-W. Wang, Mechanochemical Organic Synthesis, *Chem. Soc. Rev.*, 2013, **42**, 7668–7700.
- (146) G.-W. Wang, Fullerene Mechanochemistry: Serendipitous Discovery of Dumb-Bell-Shaped C₁₂₀ and Beyond, *Chinese J. Chem.*, 2021, **39**, 1797–1803.
- (147) A. O. Patil; D. Y. Curtin and I. C. Paul, Solid-State Formation of Quinhydrone from Their Components. Use of Solid-Solid Reactions to Prepare Compounds Not Accessible from Solution, *J. Am. Chem. Soc.*, 1984, **106**, 348–353.

- (148) T. Friščić, Supramolecular Concepts and New Techniques in Mechanochemistry: Cocrystals, Cages, Rotaxanes, Open Metal–Organic Frameworks, *Chem. Soc. Rev.*, 2012, **41**, 3493–3510.
- (149) P. Ying; J. Yu and W. Su, Liquid-Assisted Grinding Mechanochemistry in the Synthesis of Pharmaceuticals, *Adv. Synth. Catal.*, 2021, **363**, 1246–1271.
- (150) Y. Xu; J. Viger-Gravel; I. Korobkov and D. L. Bryce, Mechanochemical Production of Halogen-Bonded Solids Featuring P=O···I–C Motifs and Characterization via X-Ray Diffraction, Solid-State Multinuclear Magnetic Resonance, and Density Functional Theory, *J. Phys. Chem. C*, 2015, **119**, 27104–27117.
- (151) Y. Xu; L. Champion; B. Gabidullin and D. L. Bryce, A Kinetic Study of Mechanochemical Halogen Bond Formation by in Situ ³¹P Solid-State NMR Spectroscopy, *Chem. Commun.*, 2017, **53**, 9930–9933.
- (152) D. Cinčić; T. Friščić and W. Jones, A Stepwise Mechanism for the Mechanochemical Synthesis of Halogen-Bonded Cocrystal Architectures, *J. Am. Chem. Soc.*, 2008, **130**, 7524–7525.
- (153) M. Karimi-Jafari; L. Padrela; G. M. Walker and D. M. Croker, Creating Cocrystals: A Review of Pharmaceutical Cocrystal Preparation Routes and Applications, *Cryst. Growth Des.*, 2018, **18**, 6370–6387.
- (154) D. Braga; L. Maini and F. Grepioni, Mechanochemical Preparation of Co-Crystals, *Chem. Soc. Rev.*, 2013, **42**, 7638–7648.
- (155) T. Carstens; D. A. Haynes and V. J. Smith, Cocrystals: Solution, Mechanochemistry, and Sublimation, *Cryst. Growth Des.*, 2020, **20**, 1139–1149.
- (156) D. Hasa; E. Miniussi and W. Jones, Mechanochemical Synthesis of Multicomponent Crystals: One Liquid for One Polymorph? A Myth to Dispel, *Cryst. Growth Des.*, 2016, **16**, 4582–4588.
- (157) T. Friščić; D. G. Reid; I. Halasz; R. S. Stein; R. E. Dinnebier and M. J. Duer, Ion- and Liquid-Assisted Grinding: Improved Mechanochemical Synthesis of Metal–Organic Frameworks Reveals Salt Inclusion and Anion Templating, *Angew. Chem. Int. Ed.*, 2010, **49**, 712–715.
- (158) F. Gomollón-Bel, Ten Chemical Innovations That Will Change Our World: IUPAC Identifies Emerging Technologies in Chemistry with Potential to Make Our Planet More Sustainable, *Chem. Int.*, 2019, **41**, 12–17.
- (159) D. Hasa and W. Jones, Screening for New Pharmaceutical Solid Forms Using Mechanochemistry: A Practical Guide, *Adv. Drug Deliv. Rev.*, 2017, **117**, 147–161.
- (160) C. Schumacher; H. Fergen; R. Puttreddy; K.-N. Truong; T. Rinesch; K. Rissanen and C. Bolm, N-(2,3,5,6-Tetrafluoropyridyl)Sulfoximines: Synthesis, X-Ray Crystallography, and Halogen Bonding, *Org. Chem. Front.*, 2020, **7**, 3896–3906.
- (161) S. Karki; T. Friščić; W. Jones and W. D. S. Motherwell, Screening for Pharmaceutical Cocrystal Hydrates via Neat and Liquid-Assisted Grinding, *Mol. Pharm.*, 2007, **4**, 347–354.
- (162) J. L. Howard; Y. Sagatov; L. Repousseau; C. Schotten and D. L. Browne, Controlling Reactivity through Liquid Assisted Grinding: The Curious Case of Mechanochemical Fluorination, *Green Chem.*, 2017, **19**, 2798–2802.
- (163) N. Shan; F. Toda and W. Jones, Mechanochemistry and Co-Crystal Formation: Effect of Solvent on Reaction Kinetics, *Chem. Commun.*, 2002, No. 20, 2372–2373.
- (164) V. S. Pfennig; R. C. Villella; J. Nikodemus and C. Bolm, Mechanochemical Grignard Reactions with Gaseous CO₂ and Sodium Methyl Carbonate**, *Angew. Chem. Int. Ed.*, 2022, **61**, e202116514.
- (165) P. J. Campos; J. Arranz and M. A. Rodríguez, α -Iodination of Enaminones with Bis(Pyridine)Iodonium(I) Tetrafluoroborate, *Tetrahedron Lett.*, 1997, **38**, 8397–8400.
- (166) I. Jirkovsky, Studies on Enaminoketones, *Can. J. Chem.*, 1974, **52**, 55–65.
- (167) G. G. Habermehl; I. Wippermann; H. C. Krebs and S. tom Dieck, Halogenierung Und Acyloxylierung von N-Cyclohex-2-En-1-on-3-Yl-Aminosäuren/Halogenation and Acyloxylation of N-Cyclohex-2-En-1-on-3-Yl-Amino Acids, *Zeitschrift für Naturforsch. B*, 1991, **46**, 1415–1420.
- (168) K. Matsuo; S. Ishida and Y. Takuno, Reaction of Enaminones with Benzyltrimethylammonium Dichloroiodate, *Chem. Pharm. Bull. (Tokyo)*, 1994, **42**, 1149–1150.
- (169) C. P. Kordik and A. B. Reitz, Unexpected α -Chlorination of Tertiary Enaminones Using Benzyltrimethylammonium Dichloroiodate, *J. Org. Chem.*, 1996, **61**, 5644–5645.
- (170) L. M. E. Wilson; K. Rissanen and J. S. Ward, Iodination of Antipyrine with [N–I–N]⁺ and Carbonyl Hypoiodite Iodine(i) Complexes, *New J. Chem.*, 2023, **47**, 2978–2982.
- (171) A. N. Evstropov; V. E. Yavorovskaya; E. S. Vorob'ev; Z. P. Khudonogova; L. N. Gritsenko; E. V. Shmidt; S. G. Medvedeva; V. D. Filimonov; T. P. Prishchep and A. S. Saratikov, Synthesis and Antiviral Activity of Antipyrine Derivatives, *Pharm. Chem. J.*, 1992, **26**, 426–430.
- (172) O. Sakurada; C. Kennedy; J. Jehle; J. D. Brown; G. L. Carbin and L. Sokoloff, Measurement of Local Cerebral Blood Flow with Iodo [¹⁴C] Antipyrine, *Am. J. Physiol. Circ. Physiol.*, 1978, **234**, H59–H66.
- (173) O. Z. Chi; C. Hunter; X. Liu and H. R. Weiss, The Effects of Isoflurane Pretreatment on Cerebral Blood Flow, Capillary Permeability, and Oxygen Consumption in Focal Cerebral Ischemia in Rats, *Anesth. Analg.*, 2010, **110**.
- (174) D. P. Holschneider; J.-M. I. Maarek; J. Yang; J. Harimoto and O. U. Scremin, Functional Brain Mapping in Freely Moving Rats during Treadmill Walking, *J. Cereb. Blood Flow Metab.*, 2003, **23**, 925–932.
- (175) G. M. Tozer; V. E. Prise and V. J. Cunningham, Quantitative Estimation of Tissue Blood Flow Rate BT - Angiogenesis Protocols: Second Edition, In *Methods in Molecular Biology*, vol 467; Murray, C., Martin, S., Eds.; Humana Press: Totowa, NJ, 2009; pp 271–286.
- (176) Z. Tao; S. M. Gomha; M. G. Badrey; T. T. El-Idreesy and T. M. A. Eldebss, Novel 4-Heteroaryl-antipyrines: Synthesis, Molecular Docking, and Evaluation as Potential Anti-breast Cancer Agents, *J. Heterocycl. Chem.*, 2018, **55**, 2408–2416.
- (177) F. A. Dehmel Mohamed; Knochel, Paul, Preparation of New Magnesiated Functionalized Imidazoles and Antipyrines via an Iodine-Magnesium Exchange, *Synlett*, 2000, **2000**, 345–

346.

- (178) E. V Shchegolkov; Y. V Burgart; D. A. Matsneva; S. S. Borisevich; R. A. Kadyrova; I. R. Orshanskaya; V. V Zarubaev and V. I. Saloutin, Polyfluoroalkylated Antipyrines in Pd-Catalyzed Transformations, *RSC Adv.*, 2021, **11**, 35174–35181.