

Intracellular delivery of functional proteins with DNA-protein nanogels-lipids complex

Marina Mariconti^{1,‡}, Laurie Dechamboux², Marion Heckmann², Julien Gros¹, Mathieu Morel¹, Virginie Escriou², Damien Baigl¹, Céline Hoffmann^{2,*§} and Sergii Rudiuk^{1,*§}

¹*PASTEUR, UMR8640, Department of Chemistry, PSL University, Sorbonne Université, CNRS, Ecole Normale Supérieure, Paris 75005, France*

²*Université Paris Cité, CNRS, INSERM, UTCBS, F-75006 Paris, France*

*To whom correspondence may be addressed. Email: sergii.rudiuk@ens.psl.eu or celine.hoffmann@parisdescartes.fr

§ Equal contribution

‡ Current address: ClinSearch, Malakoff, France

Abstract

Using functional proteins for therapeutic purposes due to their high selectivity and/or catalytic properties can enable the control of various cellular processes, however, the transport of active proteins inside living cells remains a major challenge. In contrast, intracellular delivery of nucleic acids has become a routine method for a number of applications in gene therapy, genome-editing or immunization. Here we report a functionalizable platform constituting of DNA-protein nanogel carriers crosslinked through streptavidin-biotin or streptactin-biotin interactions and demonstrate its applicability for intracellular delivery of active proteins. We show that the nanogels can be loaded with active proteins bearing either biotin, streptavidin or strep-tag, and the resulting functionalized nanogels can be delivered into living cells after complexation with cationic lipid vectors. We use this approach for delivery of alkaline phosphatase enzyme which is shown to keep its catalytic activity after internalization by mouse melanoma B16 cells, as demonstrated by DDAO-phosphate assay. The resulting functionalized nanogels have dimensions of the order of 100 nm, contain around 100 enzyme molecules and are shown to be transfectable at low lipid concentrations (charge ratio $R_{\pm} = 0.75$). This ensures low toxicity of our system, which in combination with high local enzyme concentration ($\sim 100 \mu\text{M}$) underlines potential interest of this nanopatform for biomedical applications.

Introduction

Due to their highly specific biological activities and intrinsic biocompatibility, delivery of active proteins into cells represents great interest not only for fundamental research¹ and intracellular sensing,² but also for numerous therapeutic applications.³⁻⁵ However, despite their high potential, proteins suffer from chemical and thermal degradation, kidney clearance, and difficulty to penetrate through biological membranes.⁶ In particular, enzymes present a major limitation to be sensitive to both medium and the temperature, leading to the loss of their catalytic activity in inappropriate conditions thus limiting their applicability in both enzyme replacement therapy^{7,8} and enzyme prodrug therapy.^{9,10} Another aspect requiring consideration is the localization of proteins. Indeed, while most therapeutic proteins currently on the market have an extracellular localization, the intracellular transport of active proteins aiming cytosolic targets could strongly expand their exploitation for biomedical purposes.^{11,12} Yet the transport to the cytosol is still a major challenge, mainly due to physical impediment of the cell membrane and possible protein inactivation due to endocytosis.¹³⁻¹⁵

Various vectorization systems have thus been proposed for protein protection and intracellular delivery, such as PEGylation, conjugation with cell-penetrating (CPP) or specifically engineered peptides, and formulations with liposomes or polymeric nanoparticles.^{5,16-18} DNA represents another promising protein carrier in nanomedicine,^{19,20} due to its intrinsic biocompatibility, programmability and transfectability. In fact, nucleic acids can be readily delivered into cells by using various non-viral cationic transfection agents, among which, cationic lipids represent the most involved vectors in clinical studies.²¹ In this view, recently DNA has been bound to β -galactosidase and glucose oxidase in order to both protect and deliver them into the cells by using lipofectamine as transfection agent.^{2,22} In this case individual enzyme molecules were covalently conjugated to several single-stranded DNA oligonucleotides. In another example, three luciferase molecules inside tubular DNA origami nanostructures were delivered into the cells via polyethylenimine transfection vector.²³ A modular protein delivery system was recently proposed by Ryu et al., where 3 different proteins could be bound to 3 extremities of Y-shape DNA nanostructure via DNA-binding zinc-finger protein tags.²⁴ This system was successfully applied for cytosolic delivery of therapeutic endogenous tumor suppressor PTEN (phosphatase and tension homolog).

In order to expand DNA-based protein delivery to strong local concentrations of delivered protein, here we propose to incorporate a large number of active protein molecules into three-dimensional DNA nanogel network. We have recently reported a new type of DNA nanogels where the intramolecular crosslinking was achieved not by hybridization or covalent DNA crosslinking, but by noncovalent specific interactions between a multibiotinylated DNA (bt-DNA) and streptavidin.^{25,26} Here, we show different strategies for functionalization of these nanogels with two model cargo proteins: green fluorescent protein (GFP), and alkaline phosphatase (ALP) - enzyme which has been proposed for replacement therapy against the hypophosphatasia disorder.²⁷ And we investigate the delivery of the resulting functionalized nanogels, containing up to 100 μ M local protein concentration, into cells by using lipidic nucleic acid transfection agents, known for promoting endosomal escape after internalization of lipoplexes by disrupting the membrane integrity.²⁸

Results

Formation of complexes between DNA-protein nanogels and cationic vesicles - lipoproplexes

Experimental procedure for preparation, vectorization and delivery of DNA-streptavidin nanogels (NGs) is schematically shown in Figure 1A. The scaffold of the NGs constituted of multibiotinylated 3480 base pairs (bp) DNA (bt-DNA) was prepared by PCR as previously described.²⁶ Crosslinking of the resulting bt-DNA was performed by adding streptavidin labeled with AlexaFluor633 fluorophore to give fluorescent NGs (NG-633) enabling their tracking and analysis by fluorescence. The resulting NGs had 114 ± 36 nm apparent diameter, as measured by Atomic Force Microscopy (Figure S1A). For vectorization of the NGs, cationic liposomes constituted of equimolar amounts of zwitterionic lipid DOPE (1,2-dioleoyl-sn-glycero-3-phosphoethanolamine) and the cationic lipid DMAPAP (2-3-[bis-(3-amino-propyl)-amino]-propylamino-N-ditetradecyl carbamoyl methyl-acetamide) were used.²⁹ In analogy with the formation of DNA-lipid complexes (lipoplexes), in our case the complexation of DNA-protein nanogel with lipids would give complexes containing DNA, lipids and proteins, which we named lipoproplexes (LPX). We thus investigated the interaction between NG-633 and DMAPAP/DOPE liposomes containing 5% of fluorescent DOPE labeled with fluorescein (FITC). First, the colloidal stability was assessed for the LPX-633 obtained at different charge ratios ($R^{+/-}$) by fluorescence microscopy (Figure S2). We considered the charge of the nanogels to be dependent only on the DNA component, as streptavidin was shown not to affect the zeta-potential of bt-DNA upon its folding into nanogels,²⁶ and the charge concentration of the liposomes was calculated by considering the DMAPAP to have 3 positive groups at physiological pH. It was found that aggregation of the LPX occurred at $R^{+/-}$ higher than 1.5. We thus fixed the ratio $R^{+/-} = 1.5$ and analyzed the resulting LPX by flow cytometry, a method which enables co-detection of different fluorophores within individual particles of nanometric size.³⁰ The resulting dot-plots for NGs-633 only, FITC-labeled liposomes only and LPX-633 are given in the Figure 1B. For NGs and liposomes, the dot plots demonstrate a higher fluorescence signal for AlexaFluor633 (71.96%) and FITC (75.99%) signals respectively. However, when NGs and liposomes were mixed together, a large population of particles demonstrating the signal from both fluorophores (86.34%) was detected, indicating successful complexation. The observation that the mean FITC fluorescence in the LPX decreased, while the mean AlexaFluor633 fluorescence increased can be explained either by FRET energy transfer between the fluorophores, or by change of their fluorescent properties in lipidic microenvironment,^{31,32} both further confirming binding between DNA-protein nanogels and cationic lipids leading to the efficient formation of the lipoproplexes.

lipoproplexes demonstrated corresponding fluorescence inside B16 cells, regardless of the fluorescent labeling. In contrast, only poor fluorescent signal was detected in B16 cells in the control samples where cells were treated with NG-633 in absence of liposomes or with non-conjugated AlexaFluor633-streptavidin protein in presence of liposomes. Contrary to several previous reports on intracellular delivery of DNA-based nanomaterials,³³⁻³⁵ we did not detect any internalization of our nanogels in cells in absence of the transfection agent. This difference in internalization could be due to a difference in rigidity between the highly structured tetrahedral DNA nanostructures having edges and corners³⁴ and our soft nanogels, as it has been demonstrated that the penetration through the membrane depends on the rigidity of the particle.³⁶ On the other hand, it has already been shown that the use of a transfection agent could increase the internalization of a DNA-based nanogel in cells.³⁵ Moreover, we propose that this agent can allow the nanogel to escape degradation in lysosomes. Indeed, Liang et al. showed that bare DNA tetrahedron, after endocytosis, ends up trapped in lysosomes,³⁴ whereas transfection agents, such as cationic lipids, are known to allow the exit of the endosomes.²⁸

The kinetics of lipoproplex internalization for up to 2 hours of cells incubation was analyzed. Lipoproplex samples (or controls corresponding to NGs in absence of liposomes) were incubated with cells for 30, 60 or 120 minutes, before cell fixation and widefield fluorescence imaging (Figures 2C and S4). We observed that nanogel internalization progressively increased upon incubation and reach a maximum of internalization after 2 hours of incubation (Figure S5). The toxicity of the lipoproplexes after internalization in B16 cells was then assessed with MTT assay (Figure 2D).³⁷ In this test, formazan absorbance at 570 nm is proportional to the number of viable cells, and no significant difference were observed between lipoproplexes, cationic liposomes and control, indicating no toxicity for the B16 cells (n=4, Kruskal-Wallis multiple comparison test). Moreover, the detection by confocal microscopy of the nanogels inside the cells 26 hours after their internalization (Figure S6), further confirmed their biocompatibility and stability in the cellular microenvironment.

We then investigated the fate of the nanogels after their internalization by the cells. First, we studied colocalization of FITC-labeled lipids with NG-633 at different time after their internalization by the cells. We found that 30 minutes after delivery, fluorescent signal corresponding to FITC-lipid could be colocalized with the NG-633 fluorescent signal, suggesting that NG-633 and FITC-lipid are still associated. After longer period (5 h and 26 h post incubation with FITC-LPX-633), only fluorescent signal corresponding to NG-633 is observed inside cells. (Figure S6). This shows the dissociation of lipids from nanogels followed by the loss of fluorescent signal of FITC-liposome probably due to “dilution” of lipids in the cell lipidic environment following a probable lipid exchange occurring during FITC-LPX-633 cell internalization. Indeed, during internalization of nucleic acid vectorized by cationic liposomes, it is known that a lipid exchange occurs with endosomal membrane inducing disruption of endosomal membrane and delivery of nucleic acid in cytoplasm.²⁸ In order to provide evidence of endosomal escape of our DNA-protein NG inside cells, we performed immunolabeling of endosomes and lysosomes in cells after delivery of lipoproplexes. For this purpose, anti-EEA1 and anti-LAMP1 antibodies were used for immunostaining of early endosomes and lysosomes, respectively. After 2 hours of incubation of cells with fluorescent LPX (in this experiment, we used AlexaFluor555 streptavidin for nanogel formation, resulting in LPX-555 to allow multi-channel observation of nanogels, cell nuclei and immunolabeling), we observed only rare events of colocalization of NG with EEA1 (Figures 2E and 2F). Interestingly, in the most cases the nanogels were found close to endosomal

compartments. We thus hypothesize that this observation was performed after endosomal escape. Concerning lysosomes, we observed no colocalization of NG with LAMP1 labelled compartments (Figure S7). The endosomal-lysosomal system is a dynamic process where intracellular membranous compartment interconvert from early endosome, that mature to late endosomes and then fused into lysosomes.³⁸ As lysosome appears to be the “last” part of this pathway, we can assume that if NG are rarely observed inside early endosomes, it is expected that NG inside lysosome may be even more rare or even impossible to observe.




Figure 2. Internalization of DNA-protein nanogels mediated by cationic liposomes. A) Scheme of experimental procedure for delivery of lipoproplexes into cells. B) Confocal microscopy observation of B16 cells treated with AlexaFluor633-labeled nanogels in absence (NG-633) and in presence (LPX-633) of cationic lipids after 2h of incubation. Grey color corresponds to actin labeled with phalloidin ($\lambda_{EX}= 488$ nm and $\lambda_{Em}= 495-550$ nm), blue to the nucleus stained with DAPI ($\lambda_{EX}= 405$ nm and $\lambda_{Em}= 415-450$ nm), and red to nanogels containing AlexaFluor633-labeled streptavidin ($\lambda_{EX}= 638$ nm and $\lambda_{Em}= 645-700$ nm). Orthogonal views are given for the cells treated with LPX-633 (z-depth = 3 μ m). C) Widefield fluorescence microscopy images of B16 cells incubated with LPX-633 during different times ($\lambda_{EX}= 578 \pm 11$ nm and $\lambda_{Em}= 641 \pm 38$ nm). D) MTT cell viability assay. Formazan absorbance for B16 cells after 2 hours of incubation with different concentrations of lipoproplexes or liposomes. Circles show data points, and bars show the

mean values \pm sd. 4 independent tests, statistic Kruskal-Wallis multiple comparison test: n.s. E-F) Confocal microscopy analysis of B16 cell with EEA1 immunolabeling after LPX-555 delivery. E) Composite image and the separate channels of AlexaFluor555-labeled nanogels (NG-555, colored in yellow λ_{Ex} = 555 nm and λ_{Em} = 645-700 nm), EEA1 (colored in magenta, λ_{Ex} = 638 nm and λ_{Em} = 645-700 nm) and nuclei (DAPI staining, colored in blue, λ_{Ex} = 405 nm and λ_{Em} = 415-450 nm). F) Zoom on the selected region from (E) and orthogonal views along the white lines (z -depth = 2.29 μ m). Arrows :nanogels colocalized with EEA1, arrowheads: nanogel not colocalized/close to EEA1.

Intracellular delivery of GFP-loaded lipopropoxes

We then explored the possibility of loading biotinylated cargo proteins into already formed NGs. Biotinylated green fluorescent protein (btGFP) was used as model protein due to its fluorescent properties. The preparation of btGFP-loaded nanogels (btGFP-NGs) was conducted in two steps: i) DNA-protein nanogels were formed with AlexaFluor633-functionalized streptavidin in the conditions described above, and ii) btGFPs was added and bound to remaining available streptavidin sites on the nanogels (Figure 3A). By AFM, the size of the resulting GFP-functionalized nanogels was measured to be 130 ± 51 nm (Figure S1B). The NGs were then characterized by flow cytometry (Figures 3B left and S8A). Colocalization of the btGFP with AlexaFluor633 fluorescence in the NGs demonstrates successful formation of GFP-functionalized NGs (btGFP-NGs-633), as 72.04% of fluorescent events were observed in the upper-right quadrant of the dot plot. The NGs were then complexed with non-fluorescent cationic liposomes in order to form the functionalized lipopropoxes btGFP-LPX-633. Flow cytometry (Figure 3B right and S8B) demonstrated that the formation of the lipopropoxes did not interfere with GFP functionality, since the signal of functional GFP protein was still co-detected with AlexaFluor633 from the nanogels after adding the liposomes, and the percentage of colocalized fluorescence events in the upper-right quadrant of the dot plot stayed quite unaffected (74.28%).

B16 cells were then incubated with the btGFP-LPX-633 and observed by confocal microscopy. Figure 3C shows the presence of fluorescent nanogels in the cells only when they have been complexed with the liposomes. Even though the fluorescence signal of the GFP was weaker, at least 76 ± 8 % (for over 400 events) of the nanogels demonstrated colocalization of AlexaFluor633 red fluorescent signal from streptavidin with the green fluorescence from the GFP (see example of composite image and orthogonal views in the Figure 3D), in agreement with the flow cytometry analysis. We typically observed 7.7 ± 4.5 of internalized particles per cell (for 140 cells) in a single confocal plane and, around 50 internalized nanogels per cell have been observed on one particular z -stack of 7 confocal planes shown in the Figure S9. Biotinylated GFP protein can thus been successfully delivered into the cells within DNA-streptavidin nanogels complexed with cationic liposome vector.

To extend the applicability of our method, we devised another functionalization approach by using GFP fused to twin-streptag oligopeptide sequence (stGFP).^{39,40} Twin-streptag is a 28 amino acids tag having 2 binding sequences for biotin-binding pocket of streptavidin, with a strong affinity (in the range of pM) for the engineered variant of streptavidin – streptactin. We first found that streptactin labeled with DY649 dye was also crosslinking the bt-DNA, but at higher protein concentrations (above 300 nM) than with the streptavidin-biotin system (Figure S10A). To functionalize the nanogels, we first obtained a stoichiometric complex of stGFP with streptactin and then used it for crosslinking of bt-DNA (Figure 3E and Figure S10B). We found that even more stGFP-streptactin complex was

necessary in this case to produce compact nanogels (around 1000 nM). We then formed the lipoproplexes with the stGFP-NG-649, incubated them to B16 cells and imaged them by confocal microscopy (Figure 3F). Here again, fluorescent spots with colocalized fluorescence from streptactin (red) and GFP fluorescence (green) were found inside the cells when they were treated with stGFP-LPX-649 (Figure 3G). These results expand the functionalizability of DNA-streptavidin (or DNA-streptactin) nanogels and demonstrate that nanogel-based approach for intracellular protein delivery can also be applicable for cloned proteins bearing not biotin, but specific oligopeptide tags.




Figure 3. Delivery of lipoproplexes functionalized with GFP. A) Scheme of post-functionalization of NG-633 with biotinylated GFP (btGFP) and preparation of the corresponding lipoproplexes (btGFP-LPX-633). B) Flow cytometry dot plots for btGFP-NG-633 (left) and btGFP-LPX-633 (right). Red lines separate the dot plots into 4 quadrants. Blue numbers give the percentage of observed events in the upper right and lower right quadrants. C) Transmission and confocal microscopy images of B16 cells incubated either with btGFP-NG-633 (bottom) or with btGFP-LPX-633 (top). Cells are imaged in transmission light and overlaid with fluorescence signal from confocal observations. Green color corresponds to GFP cargo protein (λ_{EX} = 488 nm and λ_{Em} = 495-550 nm), cyan to the

nucleus stained with DAPI ($\lambda_{\text{Ex}} = 405$ nm and $\lambda_{\text{Em}} = 415\text{-}450$ nm), and red to nanogels containing AlexaFluor633-labeled streptavidin ($\lambda_{\text{Ex}} = 638$ nm and $\lambda_{\text{Em}} = 645\text{-}700$ nm). D) Higher magnification confocal microscopy observation of the sample from (C) with orthogonal views ($z\text{-depth} = 5.79$ μm). E) Scheme of preparation of GFP-functionalized streptactin-DNA nanogels (stGFP-NG-649) based on streptag-streptactin binding, and preparation of the corresponding lipoproplexes (stGFP-LPX-649). F) Confocal microscopy observation of B16 cells incubated with stGFP-LPX-649. Green color corresponds to GFP cargo protein ($\lambda_{\text{Ex}} = 488$ nm and $\lambda_{\text{Em}} = 495\text{-}550$ nm), cyan to the nucleus stained with DAPI ($\lambda_{\text{Ex}} = 405$ nm and $\lambda_{\text{Em}} = 415\text{-}450$ nm), magenta to the tubulin labeled with antibody directed against mouse tubulin, labelled with Alexa-Fluor555 ($\lambda_{\text{Ex}} = 552$ nm and $\lambda_{\text{Em}} = 584\text{-}613$ nm), and red to nanogels containing DY649-labeled streptavidin ($\lambda_{\text{Ex}} = 638$ nm and $\lambda_{\text{Em}} = 645\text{-}700$ nm). G) Higher magnification confocal microscopy observation of the sample from (F) with orthogonal views ($z\text{-depth} = 2.65$ μm).

Intracellular delivery of enzyme-loaded lipoproplexes

We finally investigated whether the lipoproplexes could be used as carriers for an active enzyme, and if the enzyme would keep its catalytic activity after their delivery into the cells. For this purpose, bt-DNA was directly crosslinked in one step with commercially available streptavidin conjugated to alkaline phosphatase (ALP) enzyme (Figure 4A) ²⁶ to give functionalized nanogels (ALP-NGs). The formation of ALP-NGs was characterized by AFM imaging (Figures S11) and the concentration of streptavidin-ALP conjugate of 15 nM (for 1 μM of bt-DNA in phosphate groups) was selected for further experiments, giving an apparent diameter of 133 ± 41 nm (Figure S1C). This ratio of streptavidin-ALP to bt-DNA corresponds to up to 100 enzymes per nanogel. The complexation of non-fluorescent ALP-NGs with FITC-labeled liposomes was then confirmed by flow cytometry (Figure S12). We observed that complexation with functionalized nanogels led to increase of mean fluorescence signal distribution of liposomes from the charge ratio $R_{\pm} = 0.75$ (i.e. 0.5 μM liposomes for 2 μM bt-DNA). Such a low R_{\pm} was thus sufficient for the formation of ALP-loaded lipoproplexes (ALP-LPX).

We then checked if ALP enzyme kept its catalytic activity within the lipoproplexes *in vitro*. For the assay, 9H-(1,3-dichloro-9,9-dimethylacridin-2-one-7-yl) phosphate, diammonium salt (DDAO-P) fluorogenic substrate was used to follow the hydrolysis of its phosphate moieties by alkaline phosphatase via appearance of far-red fluorescence emission of the resulting DDAO. We found that incorporation of the ALP into the nanogels only slightly decreased the catalytic activity of the enzyme to 88% (Figure S13). Adding liposomes to the ALP alone did not affect its activity, whereas within ALP-loaded lipoproplexes the enzyme conserved 79% of its initial catalytic activity, as compared to the activity of the native ALP-streptavidin conjugate. The ALP enzyme thus stayed available to the DDAO-P substrate and catalytically active even after loading into the DNA nanogels and their complexation with cationic lipids.

Finally, delivery of ALP-functionalized lipoproplexes (ALP-LPX) into B16 cells was investigated (Figure 4B). The cells were treated with ALP-LPX for 2 hours and washed to remove any remaining nanogels. The efficiency of delivery could not be assessed for the non-fluorescent NGs, instead we directly characterized the enzymatic activity of ALP inside the cells. For this purpose, DDAO-P substrate was added into the cell medium, and after 40 minutes of incubation the substrate was removed from the medium, cells were rinsed and intracellular fluorescence of the hydrolyzed DDAO was analyzed either by flow cytometry after detaching (Trypsin-EDTA) and fixing the cells

(Figure 4C), or by fluorescence microscopy after rinsing and fixing the cells onto culture plate (Figure 4D,E, S14 and S15). We found that even in absence of added ALP, the B16 cells were able to slightly hydrolyze the DDAO-P substrate giving a basal level of alkaline phosphatase activity.⁴¹ Flow cytometry showed no effect of adding enzyme alone and enzyme in presence of liposomes on the fluorescence of the cells. The distribution of fluorescence of hydrolyzed substrate slightly increased when cells were incubated with ALP-functionalized nanogels (ALP-NG) without liposomes, but the increase in fluorescence was much stronger when the cells were incubated with ALP-LPX (Figure 4E). Figure S15 shows fluorescence distributions for 3 independent experiments, and Figure 4D gives the average fluorescence value of the triplicated ALP delivery experiments normalized by the basal level of ALP activity inside the B16 cells treated with phosphate buffer only. We systematically observed that fluorescence intensity corresponding to the hydrolyzed substrate inside the cells treated with ALP-LPX, was higher than for the controls containing phosphate buffer only, free enzyme (free ALP), ALP with liposomes or ALP-functionalized nanogels without liposomes, with significative difference for ALP-LPX vs ALP+liposomes and ALP-LPX vs. ALP alone (Kruskal-Wallis Test and Dunn's multiple comparison test). In an additional control, we compared ALP enzymatic activity inside cells incubated with functionalized and non-functionalized LPX. Figure S16 demonstrates that no enhancement of ALP enzymatic activity after delivery of the enzyme-free LPX was observed, indicating that the increased enzymatic activity in ALP-LPX was due to the catalytic activity of delivered enzyme, and not a result of increasing the basal expression of alkaline phosphatase due to the interaction of lipopropoxes with cellular environment. All these results show that the ALP has been efficiently delivered inside the cells in its active state via functionalized lipopropoxes. To our knowledge, it is a first example of efficient delivery of a large local number of active enzymes.




Figure 4. Delivery of ALP-functionalized lipopropoxes and enzymatic activity assay in the cells. A) Preparation of ALP-loaded lipopropoxes (ALP-LPX), B) Schematic representation of the experimental procedure for ALP delivery and measurement of its activity inside the cells. C) Flow cytometry distribution of hydrolyzed DDAO substrate inside B16 cells treated with ALP-LPX and in control samples. The distribution for ALP alone control is not visible, as it has practically the same shape as the one for ALP+liposomes control. D) Mean \pm sd for normalized fluorescence intensities of hydrolyzed DDAO. White bar shows intrinsic ALP activity (normalized to 100 %) for cells incubated with the phosphate buffer (in absence of ALP). Black circles show the data points. (n=3 independent experiments, Kruskal-Wallis Test and Dunn's multiple comparison test, $p=0,0412$ ALP-LPX vs ALP+liposomes and $p=0,0689$ for ALP-LPX vs. ALP alone). E) Example of combined transmission and fluorescence microscopy images of hydrolyzed DDAO substrate after treatment of B16 cells with enzyme alone (ALP) or enzyme loaded into DNA nanogels (ALP-NGs) in presence and in absence of liposomes.

Discussion

We reported a new vector-carrier-cargo platform called lipopropox, schematically represented in the Figure S17, and demonstrated its ability to transport active proteins through the cell membrane. For this purpose, we used DNA nanogels cross-linked via streptavidin-biotin interactions. This platform enables loading of the protein cargo into

DNA carrier in several ways (i.e. using streptavidinated, biotinylated or strep-tagged cargo proteins) thus opening the possibility of large number of functionalities that can be potentially introduced into a single nanogel. We used cationic liposomes for the vectorization of DNA-protein NGs and renamed the resulting complex as “lipoproplex”, which evokes the complex between DNA and liposomes (i.e. lipoplex) but is intended to emphasize the importance of the protein component in our DNA-protein nanogels. Three-dimensional structure of the nanogels demands lower concentration of cationic liposomes for complexation (down to $R_{+/-} = 0.75$), which in combination with intrinsic biocompatibility of DNA nanoscaffold results in non-toxic protein delivery system as demonstrated by the MTT assay. For a proof of concept, here we investigated delivery of two proteins: biotinylated fluorescent protein GFP and the enzyme alkaline phosphatase conjugated to streptavidin. In both cases, the complexation with cationic liposomes and subsequent internalization by the cells did not affect the respective properties of the proteins: GFP retained its fluorescence emission whereas the ALP kept its efficiency to hydrolyze the DDAO-P substrate. Both strategies thus proved to be adapted for the nanogels functionalization.

By considering that all streptavidin-ALP conjugates enter the nanogels and each NG consists of one DNA scaffold molecule, in our conditions (15 nM of strep-ALP for 1 μ M of bt-DNA in phosphates) each nanogel can contain up to 100 loaded enzyme molecules. By assuming the nanogels to have spherical morphology and 130 nm diameter, the local concentration of the loaded protein can be evaluated to around 130 μ M. High number of potentially loadable proteins can be useful for delivery into cells of multifunctional nanogels containing a large number of different proteins, and the high local concentration of the loaded proteins is promising for biomedical applications. This approach can thus be of particular interest for co-delivery of several active enzymes at controlled stoichiometry. Co-transfection of genes coding for different enzymes was suggested for cancer therapy^{42,43} and treatment of metabolic diseases.⁴⁴ But it was shown in some cases to result in the expression of only one or the other enzyme rather than the desired dual expression of both enzymes,⁴⁵ because of the competition for the use of intracellular resources available for gene expression.⁴⁶ Direct delivery of NGs containing defined enzymatic composition avoids the co-expression step and can thus overcome this issue. Moreover, high local enzyme concentrations within the nanogels can be of particular interest for creating or restoring complex metabolic pathways requiring close spatial positioning of several catalytic processes within an enzymatic cascade.

Conclusions

In conclusion, we presented a new DNA-protein nanogel platform for the transport of active loaded proteins into living cells. The loaded proteins have been efficiently internalized by the cells while keeping their specific properties, which underlines potential therapeutic interest of lipoproplexes for delivery and co-delivery into the cells of active enzymes. We anticipate that DNA-protein nanogels could be employed as nanoscaffold for the transport of other biological and chemical entities, not only by incorporating them into the nanogel by streptavidin-biotin binding, but also by orthogonal binding into DNA double helix (i.e. intercalative drugs, such as doxorubicin).²⁴ Moreover, the commercial availability of biotinylated, streptavidinated or strep-tagged recombinant proteins, nanoparticles, aptameric sequences or antibodies for the specific recognition of target molecules, opens up the possibility of using nanogels as universal scaffolds with a high degree of functionalizability that not only could encapsulate a number of desired features in a single nanosized material, but also could protect the incorporated

entities from biological microenvironment. All these results underline great potential of using DNA transfection agents for soft DNA nanomaterials delivery.

References

- (1) Kusmierz, C. D.; Bujold, K. E.; Callmann, C. E.; Mirkin, C. A. Defining the Design Parameters for in Vivo Enzyme Delivery Through Protein Spherical Nucleic Acids. *ACS Cent. Sci.* **2020**, *6* (5), 815–822. <https://doi.org/10.1021/acscentsci.0c00313>.
- (2) Samanta, D.; Ebrahimi, S. B.; Kusmierz, C. D.; Cheng, H. F.; Mirkin, C. A. Protein Spherical Nucleic Acids for Live-Cell Chemical Analysis. *J. Am. Chem. Soc.* **2020**, *142* (31), 13350–13355. <https://doi.org/10.1021/jacs.0c06866>.
- (3) Kintzing, J. R.; Filsinger Interrante, M. V.; Cochran, J. R. Emerging Strategies for Developing Next-Generation Protein Therapeutics for Cancer Treatment. *Trends Pharmacol. Sci.* **2016**, *37* (12), 993–1008. <https://doi.org/10.1016/j.tips.2016.10.005>.
- (4) Dimitrov, D. S. Therapeutic Proteins. In *Physiology & behavior*; 2012; Vol. 176, pp 1–26. https://doi.org/10.1007/978-1-61779-921-1_1.
- (5) Dean, S. N.; Turner, K. B.; Medintz, I. L.; Walper, S. A. Targeting and Delivery of Therapeutic Enzymes. *Ther. Deliv.* **2017**, *8* (7), 577–595. <https://doi.org/10.4155/tde-2017-0020>.
- (6) Maximov, V.; Reukov, V.; Vertegel, A. A. Targeted Delivery of Therapeutic Enzymes. *J. Drug Deliv. Sci. Technol.* **2009**, *19* (5), 311–320. [https://doi.org/10.1016/S1773-2247\(09\)50066-4](https://doi.org/10.1016/S1773-2247(09)50066-4).
- (7) Desnick, R. J.; Schuchman, E. H. Enzyme Replacement Therapy for Lysosomal Diseases: Lessons from 20 Years of Experience and Remaining Challenges. *Annu. Rev. Genomics Hum. Genet.* **2012**, *13* (1), 307–335. <https://doi.org/10.1146/annurev-genom-090711-163739>.
- (8) Concolino, D.; Deodato, F.; Parini, R. Enzyme Replacement Therapy: Efficacy and Limitations. *Ital. J. Pediatr.* **2018**, *44* (S2), 120. <https://doi.org/10.1186/s13052-018-0562-1>.
- (9) Penet, M.-F.; Chen, Z.; Li, C.; Winnard, P. T.; Bhujwala, Z. M. Prodrug Enzymes and Their Applications in Image-Guided Therapy of Cancer: Tracking Prodrug Enzymes to Minimize Collateral Damage. *Drug Deliv. Transl. Res.* **2012**, *2* (1), 22–30. <https://doi.org/10.1007/s13346-011-0052-0>.
- (10) Yata, V. K.; Banerjee, S.; Ghosh, S. S. Folic Acid Conjugated-Bio Polymeric Nanocarriers: Synthesis, Characterization and In Vitro Delivery of Prodrug Converting Enzyme. *Adv. Sci. Eng. Med.* **2014**, *6* (4), 388–392. <https://doi.org/10.1166/ asem.2014.1508>.
- (11) Raman, V.; Van Dessel, N.; Hall, C. L.; Wetherby, V. E.; Whitney, S. A.; Kolewe, E. L.; Bloom, S. M. K.; Sharma, A.; Hardy, J. A.; Bollen, M.; Van Eynde, A.; Forbes, N. S. Intracellular Delivery of Protein Drugs with an Autonomously Lysing Bacterial System Reduces Tumor Growth and Metastases. *Nat. Commun.* **2021**, *12* (1), 6116. <https://doi.org/10.1038/s41467-021-26367-9>.
- (12) Tian, Y.; Tirrell, M. V.; LaBelle, J. L. Harnessing the Therapeutic Potential of Biomacromolecules through Intracellular Delivery of Nucleic Acids, Peptides, and Proteins. *Adv. Healthc. Mater.* **2022**, *11* (12), 2102600. <https://doi.org/10.1002/adhm.202102600>.

- (13) Goswami, R.; Jeon, T.; Nagaraj, H.; Zhai, S.; Rotello, V. M. Accessing Intracellular Targets through Nanocarrier-Mediated Cytosolic Protein Delivery. *Trends Pharmacol. Sci.* **2020**, *41* (10), 743–754. <https://doi.org/10.1016/j.tips.2020.08.005>.
- (14) Miersch, S.; Sidhu, S. S. Intracellular Targeting with Engineered Proteins. *F1000Research* **2016**, *5*, 1947. <https://doi.org/10.12688/f1000research.8915.1>.
- (15) Fu, A.; Tang, R.; Hardie, J.; Farkas, M. E.; Rotello, V. M. Promises and Pitfalls of Intracellular Delivery of Proteins. *Bioconjug. Chem.* **2014**, *25* (9), 1602–1608. <https://doi.org/10.1021/bc500320j>.
- (16) Pisal, D. S.; Kosloski, M. P.; Balu-Iyer, S. V. Delivery of Therapeutic Proteins. *J. Pharm. Sci.* **2010**, *99* (6), 2557–2575. <https://doi.org/10.1002/jps.22054>.
- (17) Oba, M.; Tanaka, M. Intracellular Internalization Mechanism of Protein Transfection Reagents. *Biol. Pharm. Bull.* **2012**, *35* (7), 1064–1068. <https://doi.org/10.1248/bpb.b12-00001>.
- (18) Dhankher, A.; Lv, W.; Studstill, W. T.; Champion, J. A. Coiled Coil Exposure and Histidine Tags Drive Function of an Intracellular Protein Drug Carrier. *J. Control. Release* **2021**, *339* (March), 248–258. <https://doi.org/10.1016/j.jconrel.2021.09.026>.
- (19) Li, Y.; Champion, J. A. Self-Assembling Nanocarriers from Engineered Proteins: Design, Functionalization, and Application for Drug Delivery. *Adv. Drug Deliv. Rev.* **2022**, *189*, 114462. <https://doi.org/10.1016/j.addr.2022.114462>.
- (20) He, L.; Mu, J.; Gang, O.; Chen, X. Rationally Programming Nanomaterials with DNA for Biomedical Applications. *Adv. Sci.* **2021**, *8* (8), 2003775. <https://doi.org/10.1002/advs.202003775>.
- (21) Mendonça, M. C. P.; Kont, A.; Kowalski, P. S.; O'Driscoll, C. M. Design of Lipid-Based Nanoparticles for Delivery of Therapeutic Nucleic Acids. *Drug Discov. Today* **2023**, *28* (3), 103505. <https://doi.org/10.1016/j.drudis.2023.103505>.
- (22) Brodin, J. D.; Sprangers, A. J.; McMillan, J. R.; Mirkin, C. A. DNA-Mediated Cellular Delivery of Functional Enzymes. *J. Am. Chem. Soc.* **2015**, *137* (47), 14838–14841. <https://doi.org/10.1021/jacs.5b09711>.
- (23) Ora, A.; Järvihaavisto, E.; Zhang, H.; Auvinen, H.; Santos, H. A.; Kostianen, M. A.; Linko, V. Cellular Delivery of Enzyme-Loaded DNA Origami. *Chem. Commun.* **2016**, *52* (98), 14161–14164. <https://doi.org/10.1039/C6CC08197E>.
- (24) Ryu, Y.; Hong, C. A.; Song, Y.; Beak, J.; Seo, B. A.; Lee, J.; Kim, H.-S. Modular Protein–DNA Hybrid Nanostructures as a Drug Delivery Platform. *Nanoscale* **2020**, *12* (8), 4975–4981. <https://doi.org/10.1039/C9NR08519J>.
- (25) Zhou, L.; Morel, M.; Rudiuk, S.; Baigl, D. Intramolecularly Protein-Crosslinked DNA Gels: New Biohybrid Nanomaterials with Controllable Size and Catalytic Activity. *Small* **2017**, *13* (28), 1700706. <https://doi.org/10.1002/sml.201700706>.
- (26) Mariconti, M.; Morel, M.; Baigl, D.; Rudiuk, S. Enzymatically Active DNA-Protein Nanogels with Tunable Cross-Linking Density. *Biomacromolecules* *22* (8), 3431–3439. <https://doi.org/10.1021/acs.biomac.1c00501>.
- (27) Bowden, S. A.; Foster, B. L. Alkaline Phosphatase Replacement Therapy for Hypophosphatasia in Development and Practice. In *Advances in Experimental Medicine and Biology*; 2019; Vol. 1148, pp 279–322.

https://doi.org/10.1007/978-981-13-7709-9_13.

- (28) Mui, B.; Ahkong, Q. .; Chow, L.; Hope, M. . Membrane Perturbation and the Mechanism of Lipid-Mediated Transfer of DNA into Cells. *Biochim. Biophys. Acta - Biomembr.* **2000**, *1467* (2), 281–292. [https://doi.org/10.1016/S0005-2736\(00\)00226-1](https://doi.org/10.1016/S0005-2736(00)00226-1).
- (29) Byk, G.; Scherman, D.; Schwartz, B.; Dubertret, C. Lipopolyamines as Transfection Agents and Pharmaceutical Uses Thereof, 2001.
- (30) Arruda, D. C.; Lachagès, A.-M.; Demory, H.; Escriou, G.; Lai-Kuen, R.; Dugas, P.-Y.; Hoffmann, C.; Bessoles, S.; Sarra Bayrouse, G.; Malachias, A.; Finet, S.; Gastelois, P. L.; de Almeida Macedo, W. A.; da Silva Cunha, A.; Bigey, P.; Escriou, V. Spheroplexes: Hybrid PLGA-Cationic Lipid Nanoparticles, for in Vitro and Oral Delivery of siRNA. *J. Control. Release* **2022**, *350*, 228–243. <https://doi.org/10.1016/j.jconrel.2022.08.030>.
- (31) Hughes, L. D.; Rawle, R. J.; Boxer, S. G. Choose Your Label Wisely: Water-Soluble Fluorophores Often Interact with Lipid Bilayers. *PLoS One* **2014**, *9* (2), e87649. <https://doi.org/10.1371/journal.pone.0087649>.
- (32) Zhang, Z.; Yomo, D.; Gradinaru, C. Choosing the Right Fluorophore for Single-Molecule Fluorescence Studies in a Lipid Environment. *Biochim. Biophys. Acta - Biomembr.* **2017**, *1859* (7), 1242–1253. <https://doi.org/10.1016/j.bbamem.2017.04.001>.
- (33) Li, J.; Pei, H.; Zhu, B.; Liang, L.; Wei, M.; He, Y.; Chen, N.; Li, D.; Huang, Q.; Fan, C. Self-Assembled Multivalent DNA Nanostructures for Noninvasive Intracellular Delivery of Immunostimulatory CpG Oligonucleotides. *ACS Nano* **2011**, *5* (11), 8783–8789. <https://doi.org/10.1021/nn202774x>.
- (34) Liang, L.; Li, J.; Li, Q.; Huang, Q.; Shi, J.; Yan, H.; Fan, C. Single-Particle Tracking and Modulation of Cell Entry Pathways of a Tetrahedral DNA Nanostructure in Live Cells. *Angew. Chemie Int. Ed.* **2014**, *53* (30), 7745–7750. <https://doi.org/10.1002/anie.201403236>.
- (35) Walsh, A. S.; Yin, H.; Erben, C. M.; Wood, M. J. A.; Turberfield, A. J. DNA Cage Delivery to Mammalian Cells. *ACS Nano* **2011**, *5* (7), 5427–5432. <https://doi.org/10.1021/nn2005574>.
- (36) Wang, S.; Guo, H.; Li, Y.; Li, X. Penetration of Nanoparticles across a Lipid Bilayer: Effects of Particle Stiffness and Surface Hydrophobicity. *Nanoscale* **2019**, *11* (9), 4025–4034. <https://doi.org/10.1039/C8NR09381D>.
- (37) Mosmann, T. Rapid Colorimetric Assay for Cellular Growth and Survival: Application to Proliferation and Cytotoxicity Assays. *J. Immunol. Methods* **1983**, *65* (1–2), 55–63. [https://doi.org/10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4).
- (38) Hu, Y.-B.; Dammer, E. B.; Ren, R.-J.; Wang, G. The Endosomal-Lysosomal System: From Acidification and Cargo Sorting to Neurodegeneration. *Transl. Neurodegener.* **2015**, *4* (1), 18. <https://doi.org/10.1186/s40035-015-0041-1>.
- (39) Schmidt, T. G. M.; Koepke, J.; Frank, R.; Skerra, A. Molecular Interaction Between the Strep-Tag Affinity Peptide and Its Cognate Target, Streptavidin. *J. Mol. Biol.* **1996**, *255* (5), 753–766. <https://doi.org/10.1006/jmbi.1996.0061>.
- (40) Skerra, A.; Schmidt, T. G. . Applications of a Peptide Ligand for Streptavidin: The Strep-Tag. *Biomol. Eng.* **1999**, *16* (1–4), 79–86. [https://doi.org/10.1016/S1050-3862\(99\)00033-9](https://doi.org/10.1016/S1050-3862(99)00033-9).

- (41) Burghoff, S.; Gong, X.; Viethen, C.; Jacoby, C.; Flögel, U.; Bongardt, S.; Schorr, A.; Hippe, A.; Homey, B.; Schrader, J. Growth and Metastasis of B16-F10 Melanoma Cells Is Not Critically Dependent on Host CD73 Expression in Mice. *BMC Cancer* **2014**, *14* (1), 898. <https://doi.org/10.1186/1471-2407-14-898>.
- (42) Liu, Y.; Wang, H.; Zhao, J.; Ma, J.; Wei, L.; Wu, S.; Xie, T.; Shen, F.; Trojan, J.; Habib, N.; Anthony, D. D.; Wu, M.; Guo, Y. Enhancement of Immunogenicity of Tumor Cells by Cotransfection with Genes Encoding Antisense Insulin-like Growth Factor-1 and B7.1 Molecules. *Cancer Gene Ther.* **2000**, *7* (3), 456–465. <https://doi.org/10.1038/sj.cgt.7700137>.
- (43) Uckert, W.; Kammertöns, T.; Haack, K.; Qin, Z.; Gebert, J.; Schendel, D. J.; Blankenstein, T. Double Suicide Gene (Cytosine Deaminase and Herpes Simplex Virus Thymidine Kinase) but Not Single Gene Transfer Allows Reliable Elimination of Tumor Cells In Vivo. *Hum. Gene Ther.* **1998**, *9* (6), 855–865. <https://doi.org/10.1089/hum.1998.9.6-855>.
- (44) De Sabbata, G.; Boisgerault, F.; Guarnaccia, C.; Iaconcig, A.; Bortolussi, G.; Collaud, F.; Ronzitti, G.; Sola, M. S.; Vidal, P.; Rouillon, J.; Charles, S.; Nicastro, E.; D'Antiga, L.; Ilyinskii, P.; Mingozi, F.; Kishimoto, T. K.; Muro, A. F. Long-Term Correction of Ornithine Transcarbamylase Deficiency in Spf-Ash Mice with a Translationally Optimized AAV Vector. *Mol. Ther. - Methods Clin. Dev.* **2021**, *20* (March), 169–180. <https://doi.org/10.1016/j.omtm.2020.11.005>.
- (45) Gupta, S.; Shah, B.; Fung, C. S.; Chan, P. K.; Wakefield, D. L.; Kuhns, S.; Goudar, C. T.; Piret, J. M. Engineering Protein Glycosylation in CHO Cells to Be Highly Similar to Murine Host Cells. *Front. Bioeng. Biotechnol.* **2023**, *11* (February), 1–14. <https://doi.org/10.3389/fbioe.2023.1113994>.
- (46) Di Blasi, R.; Marbiah, M. M.; Siciliano, V.; Polizzi, K.; Ceroni, F. A Call for Caution in Analysing Mammalian Co-Transfection Experiments and Implications of Resource Competition in Data Misinterpretation. *Nat. Commun.* **2021**, *12* (1), 2545. <https://doi.org/10.1038/s41467-021-22795-9>.

Supporting Information.

The Supporting Information is available free of charge at XXX
Materials and methods, and Supplementary Figures S1–S17.

Acknowledgment

This work was supported by Agence Nationale de la Recherche (ANR) grants ANR-22-CE06-0009 and ANR-18-CE07-0001 (S.R.), ANR-18-CE18-0005-01 (C.H.) and ANR-18-CE06-0019 (D.B.). We thank the Plateforme d'Imagerie Cellulaire et Moléculaire, US25 Inserm, UAR3612 CNRS, Université Paris Cité, Faculté de Pharmacie, Paris, France for access to confocal microscopy.

Notes

The authors declare no competing financial interest.

TOC Graphic

