Regioselective Hydroamination of Unactivated Olefins with Diazirines as a Diversifiable Nitrogen Source

Qingyu Xing¹, Preeti P. Chandrachud², Khalilia Tillett¹, and Justin M. Lopchuk^{1,2,3*}

¹Department of Chemistry, University of South Florida, Tampa, FL 33620, USA.

²Drug Discovery Department, H. Lee Moffitt Cancer Center and Research Institute, 12902 Magnolia Drive, Tampa, FL 33612, USA.

³Department of Oncologic Sciences, College of Medicine, University of South Florida, Tampa, FL 33612, USA.

Abstract

Nitrogen-containing compounds, such as amines, hydrazines, and heterocycles, play an indispensable role in medicine, agriculture, and materials. Alkylated derivatives of these compounds, especially in sterically congested environments, remain a challenge to prepare. Here we report a versatile method for the regioselective hydroamination of readily available unactivated olefins with diazirines. Over fifty examples are reported, including the protecting group-free amination of fourteen different natural products. A broad functional group tolerance includes alcohols, ketones, aldehydes, and epoxides. The proximate products of these reactions are diaziridines, which, under mild conditions, are converted to primary amines,



hydrazines, and heterocycles. Five target- and diversity-oriented syntheses of pharmaceutical compounds are shown, along with the preparation of a bis-¹⁵N diazirine validated in the late-stage isotopic labeling of an RNA splicing modulator candidate.

Introduction

Amines, hydrazines, and their derivatives persist as pivotal heteroatomic species in medicinal chemistry and drug discovery, playing an indispensable role in the development of novel therapeutic agents.^{1,2} Consequently, chemists dedicate a significant portion of their time and effort to the synthesis, functionalization, and purification of nitrogen-containing compounds. Alkyl amines play a crucial role across a spectrum of disease states, including for the treatment of pain, allergies, inflammation, lipid disorders, and cancer (**Fig. 1A**).² The syntheses of these compounds can pose significant challenges, particularly in sterically congested environments or when pursuing diversity-oriented routes. While archetypical C–N bond forming reactions such as alkylations, cross-couplings, and reductive aminations dominate the toolkit of most medicinal chemistry programs, the escalating demand for rapid access into more diverse and sp³-rich chemical space necessitates the development of new reactions, especially those that allow for non-traditional retrosynthetic disconnections and the incorporation of electrophilic amine sources.

In principle, a hydroamination reaction addresses this problem by leveraging the ubiquity, structural variety, and commercial availability of olefins in conjunction with reactive nitrogen species in order to produce alkyl amines.³ However, numerous challenges exist in the development of intermolecular versions of these reactions: overcoming the activation barrier of remote olefins with good chemo- and regioselectivity, driving the reaction to completion while avoiding a large excess of either the olefin or amination reagent, and further manipulation of potentially undesired functionality retained on the nitrogen of the resulting products. In recent years, radical-based hydroaminations have become a popular way to circumvent several of these challenges.^{4,5} Since the



Figure 1: Nitrogen-containing drugs and strategies for their syntheses. A. Pharmaceuticals with a common alkylamine motif. B. Radical hydroamination approaches via metal-hydride hydrogen atom transfer. C. This work. Radical hydroamination of unactivated olefins with diazirines as a diversifiable nitrogen donor and its synthetic applications.

radical intermediates are relatively high in energy, the activation barrier can be overcome, allowing for milder reaction conditions, along with better selectivity and functional group tolerance. Nevertheless, the search for a more broadly applicable, diversifiable nitrogen source remains a critical need.

The electrophilic nitrogen sources currently used in radical hydroaminations fall into one of two categories: sp^2 or sp^3 hybridized nitrogens bearing a leaving group, or sp^2 hybridized nitrogens bonded to another nitrogen or oxygen (**Fig. 1B**). The former group contains reagents such as sulfonyl azides^{6,7} that expel the sulfonyl group to give alkyl azides and N-fluorobenzenesulfonimide (NFSI),^{8,9} resulting in alkylated benzenesulfonimides. Alkyl azides have been reported for the synthesis of secondary alkyl amines with the loss of molecular nitrogen (*10,11*). Arylalkyl azo species can be prepared from tosyl diazenes, with the tosyl serving as the leaving group (*12*). The latter group of reagents, where the N=X bond accepts a carbon radical, are exemplified by azodicarboxylates (*6,13*) (to form protected hydrazines), and electron-deficient diazo compounds (*14,15*) (to form hydrazones). Aromatic nitro groups are also effective nitrogen sources, delivering hindered secondary anilines (*16,17*). While each of these approaches offers a distinct pathway into alkyl-substituted amines via C–N bond formation, all bear extra functionality that may or may not be desirable in the final products making further diversification challenging. Alternatively, diazirines have been shown to engage in both radical and electrophilic amination processes to afford diaziridines that serve as isolable masked amines and hydrazines (*18,19*). They are readily cleaved under mild conditions to the corresponding free amines or hydrazines with concomitant recovery of the ketone backbone, which is then recycled into the diazirine reagent synthesis. The diaziridines

can also be used directly in a variety of one-pot/telescoped syntheses of heterocycles (e.g. azoles, pyrroles, aziridines, phthalazinones), obviating the need for the often-troublesome purification of highly polar free amines and hydrazines. Here we report the practical, regioselective hydroamination of unactivated olefins with diazirines (**Fig. 1C**). Over fifty examples demonstrate the broad scope and functional group tolerance of this reaction. Synthetic applications include target- and diversity-oriented syntheses of numerous pharmaceuticals and clinical candidates. Finally, we report the synthesis of bis-¹⁵N diazirine **1**, using ¹⁵NH₃ as the sole source of the isotopic label, and its deployment in the synthesis of a ¹⁵N-labeled RNA splicing modulator.

Reaction Development and Optimization

Our initial forays into the development of diversifiable amination reagents centered around the use of diazirines, which are historically known for their utility as carbene precursors in chemical biology (20). Instead, we recently demonstrated their use in transition metal-catalyzed and photochemical decarboxylative aminations. While many carboxylic acids and their derivatives are commercial and widely available, this is far less true of the tertiary acids that would enable access to the alkyl amines required for the pharmaceuticals displayed in Figure 1A. Furthermore, synthetic access into these tertiary acids, especially when remote from other functional groups, is not straightforward. To overcome this hurdle, we turned our attention to the use of olefins in concert with diazirine 1. Substituted olefins are abundantly available, in both commercial building blocks and natural products, and are simple to prepare in the laboratory via robust chemical processes (e.g. alkylation, cross- or ring-closing metathesis, Wittig, etc.).

Carreira's elegant hydrohydrazination work served as inspiration for the immediate starting point due to the mild cobalt-catalyzed conditions and good regioselectivity that were reported (6). Indeed, upon initial reaction of diazirine **1** with 4-phenylbut-1-ene, low yields of the corresponding diaziridine were observed. While this served as confirmation that the C–N bond formed, extensive optimization of solvent, catalyst, and hydride source proved ineffective for raising the yield. However, inspired by Nojima, di-*t*butyl peroxide was added to facilitate the formation



Figure 2. Optimization of the reaction. ^aReactions were conducted with 2 (0.1 mmol), 1 (0.15 mmol, 1.5 eq), PhSiH₃ (0.1 mmol, 1 eq), cat-1 (0.005 mmol, 0.05 eq), DCE (0.4 mL) and IPA (0.1 mL), 40 °C for 20 h under argon atmosphere. ^bIsolated yield. ^cReaction conducted with 2 (0.1 mmol), 1 (0.15 mmol, 1.5 eq), PhSiH₃ (0.1 mmol, 1 eq), Mn(dpm)₃ (0.005 mmol, 0.05 eq), DCE (0.4 mL) and IPA (0.1 mL), 0 °C for 2 h under argon atmosphere. ^dMarkovnikov/anti-Markovnikov ratio. ^eReaction run for 30 h. ^f5.5 M solution in decane (dried over 4Å molecular sieves). ^gReaction run under air atmosphere with no protection from light. TES = triethylsilane, PHMS = polymethylhydrosiloxane.

of the cobalt hydride (*21*). To our delight, this proved the key to unlocking the reaction (see SI for proposed mechanism and catalytic cycle). After a switch to olefin **2** as our model substrate (lower volatility compared to 4-phenylbut-1-ene), a re-optimization of catalyst, solvent, and hydride source afforded diaziridine **3a** in 99% isolated yield (**Fig. 2**).

Other cobalt catalysts such as Co(dpm)₃ (entry 1), and Co(TPP)CI (entry 2) gave trace amounts of the desired product. Mn(dpm)₃ (entry 3), failed to show any appreciable regioselectivity (*ca.* 1.5:1), however, the products were obtained in 86% isolated yield after only 2 h at 0 °C. Ultimately, this turned out to be an excellent protocol for symmetrical olefins and will be discussed further in Figures 3 and 5. Other counterions for the salen catalyst (e.g. OAc in **cat-2**, Cl in **cat-3**, entries 4 and 5), led to diminished yields as did Carreira's catalyst (**cat-4**, entry 6). While a mixed solvent system (DCE:IPA 4:1) was found to be optimal for most substrates, IPA alone (entry 7) gave **3a** in a slightly lower yield and longer reaction time (30 h). The "IPA only" conditions proved to be useful with tri-substituted alkenes as the substrate scope was further evaluated. Although the olefin was consumed

within 3 h, a longer reaction time was required to achieve completion. At 3 h the isolated yield was 66% (entry 8), whereas prolonging the reaction time to 16 h increased the isolated yield to 93% (entry 9) and to 20 h, 99% (optimized conditions). As mentioned above, di-*t*-butyl peroxide was a crucial additive; in its absence, the yield dropped to 38% (entry 10). When using *t*-butyl peroxide instead, the yield decreased to 36% (entry 11), and the Mukaiyama hydration product was isolated instead as the major product. Other silanes, such as TES and PHMS, gave only trace amounts of **3a** (entries 12 and 13). Higher temperatures were deleterious to the yield (60 °C, 56% yield, entry 14) as was running the reaction with no precautions (open to air and light, 54% yield, entry 15).

Substrate Scope



Figure 3. Scope of the hydroamination - building blocks and pharmaceutical core stuctures. Reactions conducted with alkene (0.2 mmol), 1 (0.3 mmol, 1.5 eq), cat-1 (0.01 mmol, 0.05 eq), *t*-BuOO*t*-Bu (0.2 mmol, 1 eq), PhSiH₃ (0.2 mmol, 1 eq), DCE (800 μ L) and IPA (200 μ L) at 40 °C for 20 h under argon atmosphere with protection from light. Isolated yields are reported. ^aReaction run at 0.1 mmol scale. ^bReaction run at 1 g scale. ^cReaction run with IPA (1 mL) as the only solvent for 30 h. ^dReaction conducted with alkene (0.2 mmol), 1 (0.3 mmol, 1.5 eq), Mn(dpm)₃ (0.01 mmol, 0.05 eq), PhSiH₃ (0.2 mmol, 1 eq), DCE (800 μ L) and IPA (200 μ L) at 0 °C for 2 h under argon atmosphere with protection from light.

With the optimized conditions in hand, the scope of building blocks containing unactivated olefins was broadly evaluated. Monosubstituted olefins (**Fig. 3A**), cyclic or acyclic disubstituted olefins (**Fig. 3B**), cyclic or acyclic trisubstituted olefins (**Fig. 3C**), and *cis*-cyclic olefins (**Fig. 3D**) all performed well, affording the desired diaziridine products as single regioisomers in good to excellent yields. Nitroarenes (**3b**) and classical aryl and aliphatic amine protecting groups were tested (e.g. Boc (**3c**, **3n**, **3o**, **3aa and 5a/b**), Ac (**3d**), Cbz (**3e**) and Ts (**3p**, **3w** and **3x**); all were found to be compatible with the reaction conditions and, importantly, offer the opportunity for downstream synthetic manipulations of the orthogonally protected "diamine" motifs. In a similar vein, a free amine was tolerated (**3f**, 54% yield), and the presence of a nitrile group (**3j**) was also supported. Free hydroxy groups (**3q**, **4a**) were compatible, as were many of their common protecting groups including acetyl (**3r**), MOM (**3s**), THP (**3t**), Cbz (**3u**), Bn (**3v**) and TBS (**4e**). Phenolic (**3a-f**, **3g**) and thiophenolic ethers (**3l**), ketones (**4b**) and esters (**4f** and **4g**) could be incorporated affording the desired products in good to excellent yields.

Various simple hydrocarbons were successfully employed: linear alkenes (**3h**, **3i**), methylenecyclohexane derivatives (**3ab**, **3ac**) phenyl derivatives (**3k**, **3m**, **3y**), an adamantyl derivative (**3z**), methyl cyclohexenes (**4c**, **4d**), norbornene (**5c/d**), and cyclooctene (**5f/g**). Heterocyclic substrates included Boc- or Ts-protected piperidines (**3o**, **3p**, **3w**, **3x**, **3aa**), an azepine (**5a/b**), and an 8-amino quinoline (**3n**). This last example is notable since both the starting material and diaziridine product may act as a ligand with the catalyst, yet **3n** is still obtained in 54% isolated yield. *Cis*-olefins were generally hydroaminated in good yields (**5c**, **5f**), however with azepine derivative **5a**, a lower isolated yield was obtained (44%) due to loss of the Boc protecting group. This appears unique to this example since this phenomenon was not observed with other Boc-protected substrates (e.g. **3o**, **3n**, **3aa**). While the cobalt-catalyzed protocol worked well in these examples, for symmetrical alkenes (**Fig. 3D**) the manganese-catalyzed version proved superior, giving the desired products in shorter reaction times with better yields (**5b**, **5d**, **5g**). To our delight, [1.1.1]propellane engaged well with the alternative manganese conditions, delivering **5e** in 72% yield, which affords a different pathway into amine, hydrazine, and heterocyclic bicyclopentyl building blocks (*22*).

The reaction was conducted on gram-scale where **3h** was isolated in 73% yield. Previously we demonstrated the use of a perfluorinated diazirine reagent (with C_8F_{17}) that allowed for both the initial diazirine reaction and subsequent diaziridine cleavage to be incorporated into fluorous phase workflows (*18*). This eliminates the need for column chromatography and expedites the library synthesis of nitrogen-containing compounds for discovery scientists. Monosubstituted olefins (**3g**), disubstituted olefins (**3x**), and trisubstituted olefins (**4d**) all gave the expected hydroamination products, though **4d** suffered a loss in yield. Many of the olefins displayed in Figure 3 can be readily crafted into pharmaceuticals through known transformations including clobenzorex (from **3k**), labetalol (from **3m**), neramexane (from **3ac**), quinocide (from **3n**) and azelastine (from **5a/b**).

Having evaluated the initial scope and functional group compatibility of the hydroamination we turned our attention to the late-stage functionalization of more complex natural products with an emphasis on terpenoids (Fig. 4). Terpenes and terpenoids, many of which are commercially available, are not only common in the food and fragrance industries, but also serve as effective building blocks in medicinal chemistry and natural product synthesis. While these scaffolds appear sporadically throughout the hydrofunctionalization literature, to date there is no general hydroamination that has been demonstrated to be effective across numerous members of the class. Classical methods reported for C-N bond formation on terpenes include the Ritter reaction (23), mercuration with anilines and azides (24,25), and others. More recent hydrofunctionalization methods include Boger's hydroazidation of citronellol (26), Carreira's hydrohydrazination of camphene (6), Glorius' iminative bisfunctionalization of perillol, limonene, and nootkatone (27), Lin and Xu's independent approaches to the electrochemical diazidation of limonene oxide and nootkatone (28,29), and Engle's aminoarylation of sclareol (30). Monosubstituted olefins that successfully underwent the diazirine-based hydroamination include eugenol (6a) and sclareol (6i). Reaction on the disubstituted isopropenyl group of nookatone (6b), cannabidiol (6c), perillol (6d), limonene oxide (6e), isopulegol (6f), valencene (6g), and limonene (6k) all afforded the desired products in moderate to good yields. Bisabolol (6h), citronellic acid (6l), citronellal (6m), and citronellol (6n) serve as examples of suitable trisubstituted prenyl groups that were able to undergo the hydroamination. Finally, camphene was converted to 6j selectively as the exo-product in 75% yield.

Importantly, each of the above reactions were performed directly on the natural products, without the need for protecting groups. As a result, the functional group tolerance was further illuminated: α , β -unsaturated ketones (**6b**), free phenols (**6a**, **6c**), allylic alcohols (**6d**), aliphatic alcohols (**6f**, **6h**, **6i**, **6n**), epoxides (**6e**), carboxylic acids (**6l**) and aldehydes (**6m**) were all tolerated. Notably, kinetically-driven chemoselectivity was observed whereby

disubstituted olefins reacted significantly faster than trisubstituted olefins. Thus, for cannabidiol (**6c**), perillol (**6d**), valencene (**6g**), bisabolol (**6h**), and limonene (**6k**) the observed products were a result of mono-hydroamination only on the isopropenyl group, leaving the trisubstituted olefins untouched. However, when the reaction time was increased to 30 h in the presence of a twofold amount of diazirine, catalyst, peroxide, and silane, the bis-hydroamination products appeared and slowly became more dominant. Presumably this is a steric effect of the initial binding of the metal and migratory insertion to the olefin. However, in the mono-aminated substrates displayed in Figure 3 and 4, the steric environment around (but not directly attached to) the reactive olefin



Figure 4. Scope of the hydroamination - terpene natural products. Reactions conducted with alkene (0.2 mmol), 1 (0.3 mmol, 1.5 eq), cat-1 (0.01 mmol, 0.05 eq), *t*-BuOO*t*-Bu (0.2 mmol, 1 eq), PhSiH₃ (0.2 mmol, 1 eq), DCE (800 μ L) and IPA (200 μ L) at 40 °C for 20 h under argon atmosphere with protection from light. Isolated yields are reported. ^aReaction run with IPA (1 mL) as only solvent for 30 h.

appears to have significantly less influence as shown in **30**, **6c**, and **6j**.

Synthetic applications

With the substrate scope in hand, we endeavored to highlight the utility of the diaziridine intermediates through their conversion to amines, hydrazines, and various heterocyclic species. As shown in Figure 5, this was achieved with a variety of target-oriented syntheses, diversity-oriented approaches, and late-stage functionalization, all from diazirine reagent 1. Primaquine, a member of the 8-aminoquinoline class of drugs, is used for the prevention and treatment of malaria and is found on the World Health Organization's List of Essential Medicines. Quinocide is the constitutional isomer of primaquine and the major contaminant formed during its synthesis (*31*). Access to quinocide (8) is critical for quality control and impurity assessment of primaquine; however, its availability is somewhat limited and at an exorbitant cost. The treatment of quinoline 7 under the cobalt-catalyzed conditions gave diaziridine **3n** in 54% yield, which was then exposed to TMSCI and LiCl in DMF to reveal the amine with concomitant cleavage of the Boc group to produce quinocide **+HCI (8)** in 80% yield (**Fig. 5A**). This compares favorably to the previous route where the amine was installed via a six-step sequence originating from nitroethane (*32*). Mecamylamine is an antagonist of nicotinic acetylcholine receptors that is used for smoking cessation and hypertension (*33*). Following the selective hydroamination of camphene (**9**) as previously discussed, diaziridine **6j** was converted to the free amine with HI (82% yield) followed by reductive amination with paraformaldehyde to afford mecamylamine•HCI (**10**) in 62% yield. This provides a more tractable

preparatory scale route for discovery, which avoids the use of hydrogen cyanide that is common in the industrial process (**Fig. 5A**). Neramexane is an NMDA antagonist that is being clinically investigated for a number of



Figure 5. Synthetic applications: Target- and diversity-oriented synthesis of pharmaceuticals and related compounds via hydroamination. A. Target-oriented synthesis of mecamylamine, neramexane, and quinocide. B. Diversity-oriented synthesis of four piperidine-based fragments of a muscarinic M1 receptor candidate. C. Target-oriented and late-stage functionalization approaches to a splicing modulator candidate.

indications including Alzheimer's disease and tinnitus (34,35). The reaction of diazirine 1 with 1,1,3,3-

tetramethyl-5-methylenecyclohexane (11) produced diaziridine **3ac** in 59% yield, which was hydrolyzed in the presence of HI to furnish neramexane•HCl (12) in 57% yield (Fig. 5A).

Synthetic medicinal chemistry workflows are best expedited through common intermediates that can be readily diversified for structure-activity relationship (SAR) studies. Azaspirocycle **19** is a muscarinic M1 receptor agonist from Heptares Therapeutics with potential applications in various neurological disorders (*36*). Piperidine building blocks **15**, **16**, and **18** were targeted by the researchers via three distinct routes and three different sets of intermediates using classical chemistry: reductive aminations, $S_N 2$ displacements, and nucleophilic additions. Piperidine **17**, an obvious derivative to be examined for SAR in this subset of fragments, could not be prepared through these routes and thus was not evaluated. In a more streamlined approach, piperidines **13** and **14** were subjected to the hydroamination conditions with diazirine **1** to afford diaziridines **3p** and **3w** in 87% and 67% yield, respectively. Treatment with either pyrazole-forming conditions (*Method A*) or cleavage to the free amine (*Method B*) rapidly delivered all four building blocks (**15-18**) in 55-77% yield (**Fig. 5B**). Taken together with our previously reported diversification reactions, it is easily envisioned how a small number of strategically chosen

intermediates can be multiplied into a large library of medicinally relevant scaffolds covering a significant amount of chemical space.

Phthalazinone **25** is a splicing modulator, developed by Remix Therapeutics, that may be useful in treating disease through targeting RNA (*37*). In the original route, it was prepared from 6-bromophthalazin-1(2*H*)-one over three steps in *ca.* 3% overall yield. The lowest yielding steps are the Mitsunobu reaction to install the azepine, followed by deprotection (**Fig. 5C**, inset). By leveraging the manganese-catalyzed version of our hydroamination, symmetrical cyclic olefin **20** was smoothly converted into diaziridine **5b** in 85% yield. Condensation of **5b** with ester **21** effected a late-stage functionalization to afford splicing modulator **25** in 64% yield. This approach can be used not only to rapidly prepare **25** in high yield (54% yield over two steps from diazirine **1**), but also interrogate the SAR around the azepine. Alternatively, a three-step approach can be conducted that would efficiently enable the late-stage variation of heterocycle **25**. Here, diaziridine **5b** was first condensed with 5-bromo-3-hydroxyisobenzofuran-1(3*H*)-one **22** in 89% yield, followed by Suzuki coupling with **24** to afford the desired product in 77% yield (58% yield over three steps from diazirine **1**) (**Fig. 5C**).

¹⁵N isotopic labeling

The stable isotopic labeling of small molecules is an important analytical tool used across a variety of fields due to its low natural isotopic abundance (e.g. ²H, ¹³C, ¹⁵N), which delivers a high signal to noise ratio when observed in NMR, MS, and MRI. Recent advances in newer techniques such as hyperpolarization further increase the sensitivity (*38*). The synthesis of ¹⁵N-labeled pharmaceuticals and agrochemicals allows for the in vivo study of both the metabolism and environmental fate of candidate molecules (**Fig. 6A**) (*39*). ¹⁵N NMR has



Figure 6: Late-stage ¹⁵**N installation and its potential applications. A.** Applications of ¹⁵N-containing molecules. **B.** Hydroamination with ¹⁵N-diazirine to insert ¹⁵N-atoms into target molecules. **C.** Previous unlabeled decagram synthesis of diazirine **1**. **D.** Redesigned synthesis of ¹⁵N-diazirine (¹⁵*N*-**1**) from ¹⁵N-ammonia as the only nitrogen source. **E.** Synthesis of ¹⁵N-containing drug candidate ¹⁵*N*-**25**.

also been used to study protein packing and the conformation of protein-ligand complexes (40). The incorporation of ¹⁵N monitoring by mass spectrometry enables the quantification of the metabolism of proteins and other biomolecules (e.g. metabolomics) (41). The elucidation of organic and organometallic mechanisms is also enhanced through the monitoring of reaction intermediates and determination of kinetics (42).

Despite the utility of ¹⁵N-containing molecules, their syntheses still predominantly rely on ¹⁵N-ammonium salts (*43*), or a small number of simple building blocks such as ¹⁵N-ammonia (*44*), ¹⁵N-hydroxylamine (*44*), ¹⁵N-hydrazine monohydrate (*45*), and ¹⁵N-urea (*46*). In nearly all cases, this necessitates a time-consuming radical redesign of the synthetic route for the small molecule that needs to be labeled. Several efforts have been made toward the development of an electrophilic ¹⁵N-nitrogen source. Jones reported a ¹⁵N-diazonium reagent that afforded an azo species en route to the preparation of ¹⁵N-labeled deoxynucleosides (*47*). Unkefer developed 1-chloro-1-[¹⁵N]nitrosocyclohexane for the generation of ¹⁵N-labeled amino acids (*48*). While useful for their given targets, both approaches lack generality. Bis-¹⁵N labeled diazirine (**1**) can serve as a more universal ¹⁵N-labeled electrophilic nitrogen source when coupled not only with the hydroamination chemistry described above, but also with the previously reported decarboxylative aminations. The mild reactions conditions allow for the label(s) to be installed at a late-stage, minimizing cost and waste, and the diversification of the diaziridines affords immediate access to myriad ¹⁵N amines, hydrazines, and nitrogen-containing heterocycles. Most importantly, the tedious redesign of synthetic routes would no longer be required when unlabeled diazirine **1** is used for the initial synthesis (**Fig. 6B**).

The current synthesis of diazirine 1 requires using hydroxylamine to install the first nitrogen atom and a large excess of liquid ammonia to install the second. This route is neither practical nor economically feasible for the synthesis of ¹⁵N-labeled diazirine 1 with ca. 100 eq of ¹⁵N-ammonia going to waste (Fig. 6C). While other methods exist for the conversion of ketones to diazirines, none are suitable for α, α, α -trifluoroacetophenone (44, 49). Instead, we endeavored to devise a new route to 15 N-1 that used 15 NH₃ as the lone source of isotopically labeled nitrogen (Fig. 6D). Toward this end, a modified literature procedure was used to generate ¹⁵Nphosphorus-nitrogen ylide **30** (50), which required ca. 7 equivalents of $^{15}NH_3$. This bench stable intermediate was treated with *n*-BuLi and ketone **26** to affect a Wittig-like reaction that furnished the corresponding ¹⁵N-labeled imine. Treatment of the imine with t-BuOCI followed by ¹⁵NH₃ (ca. 5 equiv.) allowed the oxidation/cyclization sequence to proceed and delivered ¹⁵N-diaziridine **32**. Oxidation with NaOCI and catalytic TBAI gave ¹⁵N-1 (> 95% yield by NMR). To demonstrate the effectiveness of the late-stage isotopic incorporation, cyclic olefin 21 was treated with ¹⁵N-1 under the manganese-catalyzed hydroamination conditions, which furnished ¹⁵N-labeled diaziridine 33 in 72% yield. Condensation of ¹⁵N-diaziridine 33 with ester 21 delivered splicing modulator ¹⁵N-25 in 75% yield (Fig. 6E). The simple reagent synthesis, mild late-stage functionalization conditions, and ability to engage ¹⁵N-1 under any present or future diazirine protocol makes this an attractive option for stable isotope incorporation.

Conclusion

In summary, we report the cobalt-catalyzed, regioselective hydroamination of unactivated olefins. The mild reaction conditions afford broad functional group tolerance including amines, alcohols, ketones, aldehydes, acids, and epoxides. Alternatively, a manganese-catalyzed variant is suitable for symmetrical olefins, delivering the products in high yields with short reaction times. Among the more than fifty examples are fourteen terpenoid natural products, all of which were directly aminated without the use of protecting groups. Synthetic applications highlighted the mild conversion of diaziridines to amines, hydrazines, and heterocycles, featuring both target-and diversity-oriented approaches for the concise preparation of neramexane, mecamylamine, quinocide, and two other clinical candidates. A redesigned route toward diazirine **1** enabled the synthesis of a bis ¹⁵N diazirine, which was demonstrated in the two-step, late-stage isotopic labeling of an RNA splicing modulator candidate.

References

- 1. E. Vitaku, D. T. Smith, J. T. Njardarson, Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **57**, 10257-10274 (2014).
- 2. N. Kerru, L. Gummidi, S. Maddila, K. K. Gangu, S. B. Jonnalagadda, A Review on Recent Advances in Nitrogen-Containing Molecules and Their Biological Applications. *Molecules* **25**, 1909 (2020).

- 3. S. Ma, J. F. Hartwig, Progression of Hydroamination Catalyzed by Late Transition-Metal Complexes from Activated to Unactivated Alkenes. *Acc. Chem. Res.* **56**, 1565-1577 (2023).
- 4. S. W. M. Crossley, C. Obradors, R. M. Martinez, R. A. Shenvi, Mn-, Fe-, and Co-Catalyzed Radical Hydrofunctionalizations of Olefins. *Chem. Rev.* **116**, 8912-9000 (2016).
- 5. M. Villa, A. J. von Wangelin, Hydroaminations of Alkenes: A Radical, Revised, and Expanded Version. *Angew. Chem. Int. Ed.* **54**, 11906-11908 (2015).
- 6. J. Waser, B. Gaspar, H. Nambu, E. M. Carreira, Hydrazines and Azides via the Metal-Catalyzed Hydrohydrazination and Hydroazidation of Olefins. *J. Am. Chem. Soc.* **128**, 11693-11712 (2006).
- B. Gaspar, J. Waser, E. M. Carreira, Cobalt-Catalyzed Synthesis of Tertiary Azides from α,α-Disubstituted Olefins under Mild Conditions Using Commercially Available Reagents. *Synthesis* 2007, 3839-3845 (2007).
- 8. P.-W. Sun *et al.*, Cobalt-Catalyzed Intermolecular Hydroamination of Unactivated Alkenes Using NFSI as Nitrogen Source. *Chin. J. Chem.* **40**, 1066-1072 (2022).
- 9. T. Qin *et al.*, Cobalt-Catalyzed Radical Hydroamination of Alkenes with N-Fluorobenzenesulfonimides. *Angew. Chem. Int. Ed.* **60**, 25949-25957 (2021).
- 10. S.-M. Jia, Y.-H. Huang, F. Wang, Aminium-Radical-Mediated Intermolecular Hydroamination of Nonactivated Olefins. *Synlett* **34**, 93-100 (2022).
- 11. S.-M. Jia *et al.*, Hydroamination of Unactivated Alkenes with Aliphatic Azides. *J. Am. Chem. Soc.* **144**, 16316-16324 (2022).
- 12. Y. Zhang *et al.*, Modular Synthesis of Alkylarylazo Compounds via Iron(III)-Catalyzed Olefin Hydroamination. *Org. Lett.* **21**, 2261-2264 (2019).
- 13. J. Waser, E. M. Carreira, Convenient Synthesis of Alkylhydrazides by the Cobalt-Catalyzed Hydrohydrazination Reaction of Olefins and Azodicarboxylates. *J. Am. Chem. Soc.* **126**, 5676-5677 (2004).
- 14. X. Shen *et al.*, Ligand-promoted cobalt-catalyzed radical hydroamination of alkenes. *Nat. Commun.* **11**, 783 (2020).
- 15. J. Zheng, J. Qi, S. Cui, Fe-Catalyzed Olefin Hydroamination with Diazo Compounds for Hydrazone Synthesis. *Org. Lett.* **18**, 128-131 (2016).
- 16. J. Gui et al., Practical olefin hydroamination with nitroarenes. Science 348, 886-891 (2015).
- 17. H. Song, Z. Yang, C.-H. Tung, W. Wang, Iron-Catalyzed Reductive Coupling of Nitroarenes with Olefins: Intermediate of Iron–Nitroso Complex. *ACS Catal.* **10**, 276-281 (2020).
- 18. P. P. Chandrachud, L. Wojtas, J. M. Lopchuk, Decarboxylative Amination: Diazirines as Single and Double Electrophilic Nitrogen Transfer Reagents. *J. Am. Chem. Soc.* **142**, 21743-21750 (2020).
- 19. V. Maharaj, P. P. Chandrachud, W. Che, L. Wojtas, J. M. Lopchuk, Photodecarboxylative Amination of Redox-Active Esters with Diazirines. *Org. Lett.* **23**, 8838-8842 (2021).
- 20. M. W. Halloran, J.-P. Lumb, Recent Applications of Diazirines in Chemical Proteomics. *Chem. Eur. J.* **25**, 4885-4898 (2019).
- 21. T. Tokuyasu, S. Kunikawa, A. Masuyama, M. Nojima, Co(III)–Alkyl Complex- and Co(III)–Alkylperoxo Complex-Catalyzed Triethylsilylperoxidation of Alkenes with Molecular Oxygen and Triethylsilane. *Org. Lett.* **4**, 3595-3598 (2002).
- 22. K. D. Bunker, N. W. Sach, Q. Huang, P. F. Richardson, Scalable Synthesis of 1-Bicyclo[1.1.1]pentylamine via a Hydrohydrazination Reaction. *Org. Lett.* **13**, 4746-4748 (2011).
- 23. M.-E. Chen *et al.*, Recent advances of Ritter reaction and its synthetic applications. *Org. Chem. Front.* **8**, 4623-4664 (2021).
- 24. J. Barluenga *et al.*, Unexpected behavior of limonene in the oxidative aminomercuration reaction with HgO/HBF₄ and aromatic amines: stereospecific synthesis of 1,2-diamines. *J. Org. Chem.* **56**, 2930-2932 (1991).
- 25. R. M. Carman, R. P. C. Derbyshire, Azacineole (1,3,3-Trimethyl-2-azabicyclo[2.2.2]octane). *Aust. J. Chem.* **56**, 319-322 (2003).
- 26. E. K. Leggans, T. J. Barker, K. K. Duncan, D. L. Boger, Iron(III)/NaBH₄-Mediated Additions to Unactivated Alkenes: Synthesis of Novel 20'-Vinblastine Analogues. *Org. Lett.* **14**, 1428-1431 (2012).
- 27. G. Tan *et al.*, Photochemical single-step synthesis of β-amino acid derivatives from alkenes and (hetero)arenes. *Nat. Chem.* **14**, 1174-1184 (2022).

- 28. J. C. Siu, J. B. Parry, S. Lin, Aminoxyl-Catalyzed Electrochemical Diazidation of Alkenes Mediated by a Metastable Charge-Transfer Complex. *J. Am. Chem. Soc.* **141**, 2825-2831 (2019).
- 29. C.-Y. Cai, Y.-T. Zheng, J.-F. Li, H.-C. Xu, Cu-Electrocatalytic Diazidation of Alkenes at ppm Catalyst Loading. *J. Am. Chem. Soc.* **144**, 11980-11985 (2022).
- 30. T. Kang *et al.*, Nickel-Catalyzed 1,2-Carboamination of Alkenyl Alcohols. *J. Am. Chem. Soc.* **143**, 13962-13970 (2021).
- 31. A. A. Elbashir, B. Saad, A. S. M. Ali, M. I. Saleh, H. Y. Aboul-Enein, Determination of quinocide as impurity in primaquine tablets by capillary zone electrophoresis. *Biomed. Chromatogr.* **23**, 464-471 (2009).
- 32. R. C. Elderfield *et al.*, Further Syntheses of Primaquine Analogs1. *J. Am. Chem. Soc.* **77**, 4819-4822 (1955).
- 33. D. Mangan *et al.*, A new synthesis and preliminary evaluation of some analogues of mecamylamine a compound with anti-addiction properties. *Org. Biomol. Chem.* **14**, 10787-10798 (2016).
- 34. A. Jirgensons, V. Kauss, I. Kalvinsh, M. R. Gold, A Practical Synthesis of tert-Alkylamines via the Ritter Reaction with Chloroacetonitrile. *Synthesis* **2000**, 1709-1712 (2000).
- 35. W. Danysz, C. G. Parsons, Neuroprotective potential of ionotropic glutamate receptor antagonists. *Neurotox. Res.* **4**, 119-126 (2002).
- 36. B. G. Albert, C. M. Stuart, P. Mark, T. B. Gerald, Muscarinic Agonists. Patent CA 2993484 C (2023).
- 37. D. Reynolds *et al.*, Systems and methods for determining patient disease load. Patent WO 2021/207553 A1 (2021).
- 38. J. Bae, G. Zhang, H. Park, W. S. Warren, Q. Wang, ¹⁵N-Azides as practical and effective tags for developing long-lived hyperpolarized agents. *Chem. Sci.* **12**, 14309-14315 (2021).
- 39. D. Tsikas *et al.*, S-Nitroso-N-acetyl-L-cysteine ethyl ester (SNACET) and N-acetyl-L-cysteine ethyl ester (NACET)–Cysteine-based drug candidates with unique pharmacological profiles for oral use as NO, H₂S and GSH suppliers and as antioxidants: Results and overview. *J. Pharm. Anal.* **8**, 1-9 (2018).
- 40. S. C. Chiliveri, A. J. Robertson, Y. Shen, D. A. Torchia, A. Bax, Advances in NMR Spectroscopy of Weakly Aligned Biomolecular Systems. *Chem. Rev.* **122**, 9307-9330 (2022).
- 41. A. Lafaye, J. Labarre, J.-C. Tabet, E. Ezan, C. Junot, Liquid Chromatography–Mass Spectrometry and ¹⁵N Metabolic Labeling for Quantitative Metabolic Profiling. *Anal. Chem.* **77**, 2026-2033 (2005).
- M. S. Davies, M. D. Hall, S. J. Berners-Price, T. W. Hambley, [¹H, ¹⁵N] Heteronuclear Single Quantum Coherence NMR Study of the Mechanism of Aquation of Platinum(IV) Ammine Complexes. *Inorg. Chem.* 47, 7673-7680 (2008).
- 43. V. G. Chandrashekhar, W. Baumann, M. Beller, R. V. Jagadeesh, Nickel-catalyzed hydrogenative coupling of nitriles and amines for general amine synthesis. *Science* **376**, 1433-1441 (2022).
- 44. T. Glachet *et al.*, Iodonitrene in Action: Direct Transformation of Amino Acids into Terminal Diazirines and ¹⁵N₂-Diazirines and Their Application as Hyperpolarized Markers. *J. Am. Chem. Soc.* **141**, 13689-13696 (2019).
- 45. J. M. Herbert, Preparation of di-tert-butyl[U-¹⁵N]-azodicarboxylate and [U-¹⁵N]-(s)-piperazic acid. *J. Labelled Comp. Radiopharm.* **41**, 859-862 (1998).
- 46. T. Murase, M. Oka, M. Nampei, A. Miyachi, T. Nakamura, A highly sensitive assay for xanthine oxidoreductase activity using a combination of [¹³C₂,¹⁵N₂]xanthine and liquid chromatography/triple quadrupole mass spectrometry. *J. Labelled Comp. Radiopharm.* **59**, 214-220 (2016).
- 47. B. L. Gaffney, P. P. Kung, R. A. Jones, Nitrogen-15-labeled deoxynucleosides. 2. Synthesis of 7-¹⁵Nlabeled deoxyadenosine, deoxyguanosine, and related deoxynucleosides. *J. Am. Chem. Soc.* **112**, 6748-6749 (1990).
- 48. S. N. Lodwig, L. A. Silks III, C. J. Unkefer, Synthesis of 1-chloro-1-[¹⁵N]nitrosocyclohexane, an electrophilic aminating reagent. *J. Labelled Comp. Radiopharm.* **38**, 161-167 (1996).
- 49. Q. Ibert *et al.*, One-Pot Synthesis of Diazirines and ¹⁵N₂-Diazirines from Ketones, Aldehydes and Derivatives: Development and Mechanistic Insight. *Adv. Synth. Catal.* **363**, 4390-4398 (2021).
- 50. H. J. Cristau, E. Manginot, E. Torreilles, A Convenient Synthesis of Triphenylphosphine Acylimides Using N-Lithiated Triphenylphosphine Imide. *Synthesis* **1991**, 382-384 (1991).

Data availability

All data including experimental procedures, graphical procedures, and compound characterization data are available within the article and its Supporting Information file.

Acknowledgements

We gratefully acknowledge the National Science Foundation (CHE-2301063, J.M.L., discovery and scope of the hydroamination) and the National Institutes of Health (NIGMS R35-GM142577, J.M.L., synthetic applications and ¹⁵N labeling) for support of this research. This work has also been supported in part by the Chemical Biology Core Facility at the H. Lee Moffitt Cancer Center & Research Institute, an NCI designated Comprehensive Cancer Center (P30-CA076292). We thank Harshani Lawrence Moffitt for NMR and HRMS support.

Author contributions

P.P.C. and J.M.L. conceived and designed the initial project. Q.X. and J.M.L. conceived and designed the applications and ¹⁵N labeling. Q.X, P.P.C., and K.T. performed the experimental studies. Q.X, P.P.C. K.T., and J.M.L. analyzed and interpreted experimental data. Q.X., K.T., and J.M.L. wrote the manuscript. All authors reviewed, commented on, and approved the final draft.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to Justin M. Lopchuk (Justin.Lopchuk@Moffitt.org).