

A BN-Benzvalene

Tomoya Ozaki, Sierra K. Bentley, Nina Rybansky, Bo Li, Shih-Yuan Liu*

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States

ABSTRACT: The synthesis and crystallographic characterization of BN-benzvalene, the first light-element hetero-benzvalene, is described. BN-benzvalenes are produced via photoexcitation of C5-aryl-substituted 1,2-azaborines under flow conditions. Mechanistic studies support a boron-specific, two-step photoisomerization pathway involving a BN-Dewar benzene intermediate, which is distinct from the photoisomerization pathway proposed in benzene and phospho- and sila-benzenes for the formation of their respective benzvalene analogues.


Benzene has four valence isomers:¹ benzvalene,² Dewar benzene,³ prismane,⁴ and bicyclopropenyl.⁵ Among them, benzvalene has captured the attention of synthetic chemists due to its unique bonding and reactivity.² The development of a practical and facile synthesis of benzvalene by Katz⁶ has led to the access to additional strained hydrocarbon scaffolds, including bicyclo[1.1.1]pentane^{7,8} and bicyclo[2.1.1]hexane⁹ derivatives that are of current interest as 3D bioisosteres of benzene in medicinal chemistry.¹⁰

In comparison to the carbonaceous benzvalene, main group heteroatom-containing benzvalene derivatives have been less developed. Kobayashi pioneered the first example in 1977 by synthesizing the 1,4-diphospha-benzvalene through the photoisomerization of 1,4-diphospha-benzene.¹¹ Subsequently, Regitz,¹² Ando,¹³ Tokitoh,¹⁴ Sekiguchi,¹⁵ Cummins,¹⁶ and Kyushin¹⁷ et al. independently synthesized additional phosphorus- and silicon-containing heteroatom benzvalenes (Schemes 1A and 1B).¹⁸ Conspicuously, second-row element (B, N, O) hetero-benzvalenes have remained elusive to date.¹⁹ Arguably, the involvement of heavier and larger 3rd-row main group elements (P and Si) reduces the resonance stabilization (due to weak π -bonding) of the corresponding heteroarenes and attenuates the ring strain, resulting in some thermodynamic driving force for the formation of the heavier hetero-benzvalene structures.^{17c, 20}

In this communication, we report the first example of a light-element hetero-benzvalene. Specifically, we describe the synthesis, characterization, and initial reactivity studies of the boron- and nitrogen-containing BN-benzvalene (Scheme 1C). Deuterium labeling analysis is consistent with a distinct mechanism that involves a corresponding Dewar valence isomer species (BN-Dewar benzene) as an intermediate that undergoes a photo-induced 1,2-boron shift to furnish the observed BN-benzvalene.


Our group has had a long-standing interest in the synthetic development and applications of 1,2-azaborines,²¹ which are boron(B)-nitrogen(N) containing isosteres of benzene. In collaboration with the Bettinger group, we discovered that 1,2-azaborine undergoes a clean photoisomerization to BN-Dewar benzene upon UV light irradiation ($h\nu > 280$ nm).²² Subsequently, the BN-Dewar benzene species has been investigated as potential molecular solar thermal

Scheme 1. Main Group Heteroatom-Containing Benzvalenes



fuels,^{23,24} new synthetic building blocks for 1,2-aminoborylated cyclobutane derivatives,²⁵ and as a monomer for ring-opening metathesis polymerization.²⁶ During our investigation of the photochemistry of 1,2-azaborines, we discovered that a C5-aryl-functionalized 1,2-azaborine (with *N*-TBS and *B*-Mes substitution) produced the BN-benzvalene isomer in addition to the BN-Dewar isomer upon photolysis (Scheme 2).

Scheme 2. Initial Discovery




To optimize the formation of BN-benzvalene, we selected C5-*p*-Tol-*N*-TBS-*B*-Mes-1,2-azaborine **1a** as a model substrate using a Vapourtec UV-150 continuous-flow photochemical reactor with adjustable irradiation wavelength (by applying select filters to a mercury lamp). As can be seen from Table 1, we determined that

280 nm UV irradiation with a residence time of 20 minutes in THF (0.10 M) were optimal for the generation of BN-benzvalene.

Under these conditions, we obtained BN-benzvalene **2a** in 88% yield without the formation of BN-Dewar benzene **3a** as a byproduct (Table 1, entry 1). Shortening the residence time (RT) (entry 2) or using a longer wavelength (entry 4) resulted in incomplete conversion, with BN-Dewar benzene **3a** being the major observed product. On the other hand, extending the residence time (entry 3) or using a shorter wavelength (entry 5) resulted in a slightly reduced yield of BN-benzvalene **2a**, likely due to product degradation. Hexane and CH₂Cl₂ are compatible reaction solvents, however, they produced lower yields compared to THF (entries 6 and 7). The reaction is scalable to at least 0.50 mmol (200 mg) scale without a decrease in yield (entry 8).²⁷ Notably, both products **2a** and **3a** are stable toward silica gel column chromatography under an inert atmosphere.

Table 1. Optimization of the Photoisomerization



Entry	Variations from the "standard conditions"	Conv. (%) ^a	2a	3a
1	None	>99	88	0
2	RT = 10 min	76	28	41
3	RT = 50 min	>99	81	0
4	hv > 300 nm	53	6	43
5	hv > 260 nm	>99	76	0
6	hexane, RT = 50 min	>99	77	0
7	CH ₂ Cl ₂	93	56	21
8	0.50 mmol, RT = 50 min	>99	87 (80)	0
9	0.50 mmol, RT = 50 min, hv > 300 nm	62	7	53 (44)

^a Yields were determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. Scale: 0.10 mmol, unless otherwise stated. Yields of isolated product are reported in parentheses. RT: residence time.

We were able to isolate compounds **2a** and **3a** (Table 1, entry 8 and entry 9, respectively), and Figure 1 shows their ¹H NMR spectra. In Dewar isomer **3a**, the two methyl groups on TBS (~0 ppm, orange arrow) are diastereotopic as evidenced by two distinct signals in the ¹H NMR spectrum. Similarly, the benzvalene isomer **2a** also exhibits two signals for the same methyl groups in its ¹H NMR spectrum, albeit with less chemical shift difference. The mesityl *o*-methyl groups (~2.2 ppm, purple arrow) exhibit exchange dynamics (broadened signals) at room temperature for the BN-Dewar benzene **3a** compound, whereas sharp diastereotopic mesityl *o*-methyl signals are observed for BN-benzvalene **2a**. These observations are consistent with a more restricted B–Mes bond rotation for **2a** relative to **3a**. The original 1,2-azaborine arene C–H proton signals H^a, H^b, H^c are upfield shifted in BN-benzvalene **2a** relative to the signals for BN-Dewar **3a** and are consistent with previously reported similar tricyclic structures.^{2a,19a}





Figure 1. ¹H NMR spectra in CDCl₃ of C5-*p*-Tol-BN-Dewar benzene **3a** (top trace) and *p*-Tol-BN-benzvalene **2a** (bottom trace) in CDCl₃.


We next evaluated the scope of C5-aryl functional group under our optimized conditions. A broad range of *p*-substituents are compatible, affording the corresponding products containing alkyl groups (entries **2a–2c**), trifluoromethoxy group (entry **2d**), and halogens (entries **2e–2g**) in good yields. The unsubstituted C5-Ph-1,2-azaborine substrate (entry **2h**), along with C5-3,5-dimethoxy-phenyl- (entry **2i**), and C5-*o*-tolyl- (entry **2j**) derivatives were also successfully converted to the BN-benzvalene products. Moreover, C5-1,2-azaborines containing electronically diverse heterocycles such as thiophene, pyridine, and thiazole (entries **2k–2m**) prove also to be suitable substrates.

Table 2. Reaction Scope at the C5 Position



^a Yields of isolated products are reported as an average of two trials. Scale: 0.10 mmol, unless otherwise stated. ^b 0.50 mmol scale. ^c Hexane was used as a solvent. ^d Yields were determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard; reported yields are the average of two trials. ^e 0.05 M Substrate concentration. RT: residence time.

We then turned our attention to determine the scope with respect to the nitrogen substituent. The presence of a silicon on nitrogen appears to be crucial for product formation. While silicon-based groups, such as TMS, TBS, and SiPh₃ (entries **2a**, **2n**, **2o**) generate the product cleanly, other groups on nitrogen, such as H, Me, Ph, and Ts (entries **2p–2s**) completely shut down the reaction, with unreacted starting material and/or decomposition being observed.

Table 3. Reaction Scope at the N Position

^a Yields were determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. Scale: 0.10 mmol. Reported yields are the average of two runs. ^b Yields of isolated products are reported as an average of two trials. RT: residence time.

BN-benzvalenes (**2a–2n**) are generally non-crystalline oily substances, with the exception being the *N*-SiPh₃-substituted **2o**, which is a white solid. We successfully grew crystals of **2o** (CCDC 2359775) suitable for single crystal X-ray diffraction analysis by slow evaporation from a CH₂Cl₂ solution at –40 °C. We also obtained the structure of the corresponding 1,2-azaborine starting material **1o** (CCDC 2359773) for comparative bonding analysis. Two structural features of **2o** are noteworthy (Figure 2): 1) The N-B bond distance (1.419(3) Å) in **2o** is shorter than that for **1o** (1.446(3) Å), which is consistent with loss of electron delocalization/aromaticity²⁸ after the photoisomerization. 2) In **2o**, the C–C bond distances associated with C6 (C6–C5 = 1.497(3) Å, C6–C3 = 1.494(3) Å) are shorter than the C–C bond distances associated with C4 (C4–C3 = 1.531(3) Å, C4–C5 = 1.546(3) Å). The bond distance differences are consistent with reported crystallographically characterized aminoborylated cyclopropanes.²⁹ Notably, density functional theory (DFT) calculations nicely reproduce the experimentally observed structural asymmetry in **2o** whereas the direct carbonaceous benzvalene analogue exhibits symmetrical σ-bonding according to DFT predictions (See Supporting Information for details).




Figure 2. Crystal Structure of **2o** and **1o** (hydrogen atoms are omitted for clarity). Thermal ellipsoids are drawn at the 50% probability level. Selected bond lengths in Å for **2o** are: N–B = 1.419(3), N–C6 = 1.479(3), B–C4 = 1.575(3), C4–C3 = 1.531(3), C4–C5 = 1.546(3), C6–C3 = 1.494(3), C6–C5 = 1.497(3); Selected bond lengths in Å for **1o** are: N–B = 1.446(3), N–C6 = 1.388(2), B–C3 = 1.519(3), C3–C4 = 1.355(3), C4–C5 = 1.421(3), C6–C5 = 1.355(3).

We also used DFT calculations at the B3LYP/6-311G**//B3LYP/6-31G** level of theory to evaluate the relative energies of


Table 4. Relative Energies of 1,2-Azaborine’s Valence Isomers (calculated at the B3LYP/6-311G**//B3LYP/6-31G** level of theory.)

Isomer	Relative Energy (kcal/mol)
1a	0
3a	46.3
2a	49.5
CC-1a	0
CC-3a	65.3
CC-2a	68.1

Dewar benzene and benzvalene isomers vs. their (hetero)arene structures for both the BN and CC series (Table 4). We determined that benzvalenes **2a** and **CC-2a** are slightly less stable than the Dewar benzene isomers **3a** and **CC-3a** by 3.2 and 2.8 kcal/mol, respectively). BN-benzvalene **2a** is higher in energy by 49.5 kcal/mol compared to 1,2-azaborine **1a** whereas benzvalene **CC-2a** is higher in energy by 68.1 kcal/mol relative to its benzene isomer **CC-1a**. The difference in the relative stability between the BN and CC congeners (BN: 49.5 kcal/mol vs. CC: 68.1 kcal/mol) can be ascribed to the stronger resonance stabilization energy present in **CC-1a** vs. **1a**.³⁰


We have also conducted preliminary reactivity studies of BN-benzvalene **2a** and determined that **2a** reacts with thiophenol to cleanly furnish the ring-opened adduct **4a** (Scheme 3).^{9b,31} Notably, the reaction is sluggish under dark conditions, and irradiation with purple light accelerates the reaction.³² Compound **4a** (CCDC 2359774), the first example of a BN-bicyclo[2.1.1]hexane, is stable towards air and moisture and can be isolated by column chromatography under ambient conditions.

Scheme 3. Synthesis of BN-Bicyclo[2.1.1]hexane



To elucidate the reaction mechanism for the formation of BN-benzvalene, we conducted the experiments as outlined in Scheme 4. The observations during our reaction optimization work (Table 1, entries 2 and 4 vs. entry 1) hinted at BN-Dewar benzene **3a** serving as an intermediate toward the formation of BN-benzvalene **2a**. We were able to isolate pure BN-Dewar benzene **3a** and determine that **3a** cleanly converts to **2a** under the standard reaction conditions (Scheme 4, eq 1). Thus, BN-Dewar benzene **3a** is a chemically and likely kinetically competent intermediate for BN-benzvalene formation. Next, we conducted a deuterium labeling study to identify the bond connectivity over the course of the reaction. We synthesized C6- and C3-deuterated 1,2-azaborine analogues **5a** and **7a**, respectively. These substrates, when subjected to the standard photoisomerization conditions, furnish labeled BN-benzvalene products **6a** and **8a**. Formation of **6a** (Scheme 4, eq 2) is consistent with N–C6 bond remaining connected throughout the reaction. To our surprise, the deuterium in **8a** switched its


Scheme 4. Mechanistic Studies



location to the bridgehead C-H position, indicating that B-C3 bond is cleaved during the formation of the BN-benzvalene product. By shortening the residence time, we were able to monitor the formation of the deuterium-labeled BN-Dewar benzenes **9a** and **10a**. Consistent with the previously reported mechanism for BN-Dewar benzene formation,³³ no N-C6 and B-C3 bond cleavage was observed (eqs 4 and 5). The findings illustrated in Scheme 4 indicate that the B-C3 bond must be cleaved during the transformation from BN-Dewar benzene to BN-benzvalene.

Scheme 5 illustrates our proposed mechanism that is consistent with our deuterium labeling and reaction intermediate analysis. First, the starting 1,2-azaborine undergoes a photoinduced formal 4π disrotatory electrocyclization to generate the BN-Dewar benzene intermediate.³³ Then, the styrene chromophore in the BN-Dewar benzene is excited by UV-light to generate a biradical species. The radical at the C4 position subsequently undergoes a 1,2-boron shift,³⁴ and producing a C3 radical via a C3-B bond cleavage. Finally, radical-radical recombination of the C3 and C5 radicals creates the central C-C bond of BN-benzvalene. The proposed reaction mechanism is consistent with C5-aryl functional group facilitating the generation of BN-benzvalene by providing extra conjugation in the BN-Dewar intermediate for the photoinduced formation of the biradical. The proposed 1,2-boron shift for BN-benzvalene formation is distinct from mechanisms proposed in benzene^{35,36} and phospho¹²- and sila^{14,15a}-benzene analogues to yield their respective benzvalenes.

Scheme 5. Proposed Mechanism for Formation of BN-Benzvalene



In conclusion, we presented the synthesis and crystallographic characterization of the first light-element hetero-benzvalene exemplified as the boron-nitrogen containing BN-benzvalene. Photoexcitation of C5-aryl-substituted 1,2-azaborines under flow conditions furnish BN-benzvalenes with a broad substrate scope, including halogen functional groups and heteroaromatic rings. Mechanistic studies support a distinct stepwise photoisomerization pathway from 1,2-azaborine via a BN-Dewar benzene intermediate involving a radical-induced 1,2-boron shift. BN-benzvalene reacts with thiophenol in the presence of light to generate a BN-bicyclo[2.1.1]hexane. Current efforts are directed toward exploring novel modes of reactivity of BN-benzvalene derivatives.

ASSOCIATED CONTENT

The Supporting Information is available free of charge at <https://pubs.acs.org>. Experimental procedures, compound characterization data, computational and crystallographic information (PDF)

Accession Codes

CCDC 2359773, CCDC 2359774, and CCDC 2359775 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Shih-Yuan Liu – Department of Chemistry, Boston College, Chestnut Hill, Massachusetts 02467-3860, United States; orcid.org/0000-0003-3148-9147; Email: shihyuan.liu@bc.edu

Authors

Tomoya Ozaki – Department of Chemistry, Boston College, Chestnut Hill, Massachusetts 02467-3860, United States
 Sierra K. Bentley – Department of Chemistry, Boston College, Chestnut Hill, Massachusetts 02467-3860, United States; present address: Metrohm USA, Riverview, Florida 33578, United States
 Nina Rybansky – Department of Chemistry, Boston College, Chestnut Hill, Massachusetts 02467-3860, United States
 Bo Li – Department of Chemistry, Boston College, Chestnut Hill, Massachusetts 02467-3860, United States

Notes

The authors declare no competing financial interests. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENT

Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health (NIGMS) under Award Number R01GM136920 and by Boston College IGNITE fund. We also acknowledge the NIH-S10

(award: S10OD026910), the NSF-MRI (award: CHE-2117246) for the support of Boston College's NMR facilities, and the NIH-S10 (award: S10OD030360) for the support of Boston College's X-ray facilities. T.O. thanks the Takenaka Scholarship Foundation for support. T.O. thanks Mr. Yiling Zhou for helpful discussions related to DFT calculations.

DEDICATION

This work is dedicated to Prof. Lawrence T. Scott on the occasion of his 80th birthday.

REFERENCES

- ¹ For an overview, see: (a) Bolesov, I. G. Valence Isomers of Benzene. *Russ. Chem. Rev.* **1968**, *37*, 666–670. (b) Van Tamelen, E. E. Valence-Bond Isomers of Aromatic Systems. *Angew. Chem. Int. Ed.* **1965**, *4*, 738–745. (c) Scott, L. T.; Jones M. Rearrangements and Interconversions of Compounds of the Formula (CH)_n. *Chem. Rev.* **1972**, *72*, 181–202.
- ² (a) Wilzbach, K. E.; Ritscher, J. S.; Kaplan, L. Benzvalene, the Tricyclic Valence Isomer of Benzene. *J. Am. Chem. Soc.* **1967**, *89*, 1031–1032. (b) Christl, M. Benzvalene—Properties and Synthetic Potential. *Angew. Chem. Int. Ed.* **1981**, *20*, 529–546.
- ³ (a) Dewar, J. S. On the Oxidation of Phenyl Alcohol, and a Mechanical Arrangement adapted to illustrate Structure in the Unsaturated Hydrocarbons. *Proc. R. Soc. Edinburgh* **1869**, *6*, 82–86. (b) Van Tamelen, E. E.; Pappas, S. P. Chemistry of Dewar Benzene. 1,2,5-Tri-*t*-Butylbicyclo[2.2.0]Hexa-2,5-Diene. *J. Am. Chem. Soc.* **1962**, *84*, 3789–3791. (c) Van Tamelen, E. E.; Pappas, S. P. Bicyclo [2.2.0]hexa-2,5-diene. *J. Am. Chem. Soc.* **1963**, *85*, 3297–3298. (d) Van Tamelen, E. E.; Pappas, S. P.; Kirk, K. L. Valence bond isomers of aromatic systems. Bicyclo[2.2.0]hexa-2,5-dienes (Dewar benzenes). *J. Am. Chem. Soc.* **1971**, *93*, 6092–6101.
- ⁴ (a) Ladenburg, A. Bemerkungen zur aromatischen Theorie. *Ber. Dtsch. Chem. Ges.* **1869**, *2*, 140–142. (b) Katz, T. J.; Acton, N. Synthesis of prismane. *J. Am. Chem. Soc.* **1973**, *95*, 2738–2739.
- ⁵ (a) Billups, W. E.; Haley, M. M. Bicycloprop-2-enyl (C₆H₆). *Angew. Chem. Int. Ed.* **1989**, *28*, 1711–1712. (b) Boese, R.; Bläser, D.; Gleiter, R.; Pfeifer, K. H.; Billups, W. E.; Haley, M. M. Structure and photoelectron spectrum of 3,3'-bicyclopropenyl. *J. Am. Chem. Soc.* **1993**, *115*, 743–746.
- ⁶ (a) Katz, T. J.; Wang, E. J.; Acton, N. Benzvalene synthesis. *J. Am. Chem. Soc.* **1971**, *93*, 3782–3783. (b) Katz, T. J.; Roth, R. J.; Acton, N.; Carnahan, E. J. Synthesis of Benzvalene. *J. Org. Chem.* **1999**, *64*, 7663–7664.
- ⁷ (a) Christl, M.; Brüntrup, G. Reaktionen am Benzvalen: Reduktion mit Diimin und Ozonolyse. *Chem. Ber.* **1974**, *107*, 3908–3914. (b) Stulgies, B.; Pigg, D. P.; Kaszynski, P.; Kudzin, Z. H. 9,9-Dimethyl-8,10-dioxapentacyclo[5.3.0.0^{2,5}.0^{3,6}.0^{3,6}]decane and naphthotetra-cyclo[5.1.0.0^{1,6}.0^{2,7}]oct-3-ene: new substituted [1.1.1]propellanes as precursors to 1,2,3,4-tetrafunctionalized bicyclo[1.1.1]pentanes. *Tetrahedron* **2005**, *6*, 89–95. (c) Mazal, C.; Škarka, O.; Kaleta, J.; Michl, J. *cis*-endo-Bicyclo[1.1.1]pentane-1,2,3,4-tetracarboxylic Acid and Its Derivatives. *Org. Lett.* **2006**, *8*, 749–752. (d) Kaleta, J.; Michl, J.; Mazal, C. T-Shaped Molecular Building Blocks by Combined Bridgehead and Bridge Substitution on Bicyclo[1.1.1]pentanes. *J. Org. Chem.* **2010**, *75*, 2350–2356.
- ⁸ Anderson, J. M.; Measom, N. D.; Murphy, J. A.; Poole, D. L. Bridge Functionalisation of Bicyclo[1.1.1]pentane Derivatives. *Angew. Chem. Int. Ed.* **2021**, *60*, 24754–24769.
- ⁹ (a) Roth, R. J.; Katz, T. J. Uses of benzvalene in synthesis. Synthesis of tricyclo[2.2.0.0^{2,6}]hexane. *J. Am. Chem. Soc.* **1972**, *94*, 4770–4771. (b) Herbert, R.; Christl, M. Bicyclo[2.1.1]hexane aus Tricyclo[3.1.0.0^{2,6}]hexanen. *Chem. Ber.* **1979**, *112*, 2012–2021. (c) Roth, R. J.; Katz, T. J. Reactions and rearrangements of 5,6-disubstituted bicyclo[2.1.1]hexanes. *J. Org. Chem.* **1980**, *45*, 961–965.
- ¹⁰ (a) Mykhailiuk, P. K. Saturated bioisosteres of benzene: where to go next? *Org. Biomol. Chem.* **2019**, *17*, 2839–2849. (b) Semeno, V. V.; Vasylichenko, V. O.; Fesun, I. M.; Ruzhlyo, L. Y.; Kipriianov, M. O.; Melnykov, K. P.; Skreminskyi, A.; Iminov, R.; Mykhailiuk, P.; Vashchenko, B. V.; Grygorenko, O. O. Bicyclo[m.n.k]alkane Building Blocks as Promising Benzene and Cycloalkane Isosteres: Multigram Synthesis, Physicochemical and Structural Characterization. *Chem. Eur. J.* **2024**, *30*, e202303859.
- ¹¹ (a) Kobayashi, Y.; Fujino, S.; Hamana, H.; Kumadaki, I.; Hanzawa, Y. Trifluoromethylated Diphosphabenzvalene: 1,3,4,6-Tetrakis(trifluoromethyl)-2,5-diphosphatricyclo[3.1.0.0^{2,6}]hexene-3. *J. Am. Chem. Soc.* **1977**, *99*, 8511. (b) Kobayashi, Y.; Fujino, S.; Hamana, H.; Hanzawa, Y.; Morita, S.; Kumadaki, I. Studies on organic fluorine compounds. 36. Synthesis and reactions of trifluoromethylated diphosphabenzvalene. *J. Org. Chem.* **1980**, *45*, 4683–4685.
- ¹² Blatter, K.; Rösch, W.; Vogelbacher, U.-J.; Fink, J.; Regitz, M. Isomerization Reactions in the System Dewar-Phosphinine/Phosphaprismane/Phosphabenzvalene/Phosphinine. *Angew. Chem. Int. Ed.* **1987**, *26*, 85–86.
- ¹³ (a) Ando, W.; Shiba, T.; Hidaka, T.; Morihashi, K.; Kikuchi, O. Syntheses and Characterization of Bis(silacyclopropene) and Disilabenzvalene. *J. Am. Chem. Soc.* **1997**, *119*, 3629–3630. (b) Kabe, Y.; Ohkubo, K.; Ishikawa, H.; Ando, W. 1,4-Disila(Dewar-benzene) and 1,4-Disilabenzene: Valence Isomerization of Bis(alkylsilacyclopropenyl)s. *J. Am. Chem. Soc.* **2000**, *122*, 3775–3776.
- ¹⁴ (a) Wakita, K.; Tokitoh, N.; Okazaki, R.; Takagi, N.; Nagase, S. Crystal Structure of a Stable Silabenzene and Its Photochemical Valence Isomerization into the Corresponding Silabenzvalene. *J. Am. Chem. Soc.* **2000**, *122*, 5648–5649. (b) Sugahara, T.; Sasamori, T.; Tokitoh, N. The formation of a 1,4-disilabenzene and its isomerization into a disilabenzvalene derivative. *Dalton Trans.* **2019**, *48*, 9053–9056.
- ¹⁵ (a) Nakata, N.; Oikawa, T.; Matsumoto, T.; Kabe, Y.; Sekiguchi, A. Silyl-Substituted 1,4-Disila(Dewar benzene): New Synthesis and Unexpected Insertion of CO into the Si–Si Bond To Form a Disilyl Ketone. *Organometallics* **2005**, *24*, 3368–3370. (b) Takanashi, K.; Lee, V. Y.; Ichinohe, M.; Sekiguchi, A. 1,2,5,6-Tetrasilabenzobenzvalene: A Valence Isomer of 1,2,3,4-Tetrasilanaphthalene. *Chem. Lett.* **2007**, *36*, 1158–1159.
- ¹⁶ Piro, N. A.; Cummins, C. C. Tetrachosphabenzene Obtained via a Triphosphacyclobutadiene Intermediate. *Angew. Chem. Int. Ed.* **2009**, *48*, 934–938.
- ¹⁷ (a) Tsurusaki, A.; Iizuka, C.; Otsuka, K.; Kyushin, S. Cyclopentasilane-Fused Hexasilabenzvalene. *J. Am. Chem. Soc.* **2013**, *135*, 16340–16343. (b) Tsurusaki, A.; Kyushin, S. The Radical Anion of Cyclopentasilane-Fused Hexasilabenzvalene. *Chem. Eur. J.* **2016**, *22*, 134–137. (c) For an overview, see: Ota, K.; Kinjo, R. Inorganic Benzene Valence Isomers *Chem. Asian J.* **2020**, *15*, 2558–2574.
- ¹⁸ For examples transition-metal derived rhoda- and iridabenzvalenes, see: (a) Gilbertson, R. D.; Weakley, T. J. R.; Haley, M. M. Synthesis, Characterization, and Isomerization of an Iridabenzvalene. *Chem. Eur. J.* **2000**, *6*, 437–441. (b) Wu, H.-P.; Weakley, T. J. R.; Haley, M. M. Metallabenzenes and Valence Isomers. 5. Synthesis and Structural Characterization of a Rhodabenzvalene: A Rare η²-Cyclopropene/σ-Vinylrhodium(I) Complex. *Organometallics* **2002**, *21*, 4320–4322. (c) Gilbertson, R. D.; Lau, T. L. S.; Lanza, S.; Wu, H.-P.; Weakley, T. J. R.; Haley, M. M. Synthesis, Spectroscopy, and Structure of a Family of Iridabenzenes Generated by the Reaction of Vaska-Type Complexes with a Nucleophilic 3-Vinyl-1-cyclopropene. *Organometallics* **2003**, *22*, 3279–3289. (d) Wu, H.-P.; Weakley, T. J. R.; Haley, M. M. Regioselective Formation of β-Alkyl-α-phenyliridabenzenes via Unsymmetrical 3-Vinylcyclopropenes: Probing Steric and Electronic Influences by Varying the Alkyl Ring Substituent. *Chem. Eur. J.* **2005**, *11*, 1191–1200. (e) Wu, H.-P.; Ess, D. H.; Lanza, S.; Weakley, T. J. R.; Houk, K. N.; Baldrige, K. K.; Haley, M. M.

Rearrangement of Iridabenzvalenes to Iridabenzenes and/or η^5 -Cyclopentadienyliridium(I) Complexes: Experimental and Computational Analysis of the Influence of Silyl Ring Substituents and Phosphine Ligands. *Organometallics* **2007**, *26*, 3957–3968.

¹⁹ For related oxa- or aza-tricyclohexane derivatives, see: (a) Corey, E. J.; Pirkle, W. H. Tricyclo[2^{3,6}.1.1.0]pyran-2-one. *Tetrahedron Lett.* **1967**, *8*, 5255–5256. (b) Burger, U.; Dreier, F. Reactions of nitrogen containing aromatic anions with chlorocarbene. *Tetrahedron* **1983**, *39*, 2065–2071. (c) Fink, J.; Regitz, M. Thermal cycloaddition of carbonyl compounds to a stable cyclobutadiene. *Tetrahedron Lett.* **1984**, *25*, 1711–1714. (d) Fink, J.; Regitz, M. Synthesen mit Cyclobutadienen, 7. Ein Zugang zum 2-Oxabicyclo[2.2.0]hexen/3-Oxatricyclo-[3.1.0.0^{2,6}]hexan-System durch thermische Cycloaddition von Carbonylverbindungen an ein stabiles Cyclobutadien. *Chem. Ber.* **1985**, *118*, 2255–2265. (e) Michels, G.; Hermesdorf, M.; Schneider, J.; Regitz, M. Cycloaddition von *N*-Acyliminen an Cyclobutadiene. *Chem. Ber.* **1988**, *121*, 1775–1783.

²⁰ Gimarc, B. M.; Zhao, M. Strain and resonance energies in main-group homoatomic rings and clusters. *Coord. Chem. Rev.* **1997**, *158*, 385–412.

²¹ For an overview, see: (a) Campbell, P. G.; Marwitz, A. J. V.; Liu, S.-Y. Recent Advances in Azaborine Chemistry. *Angew. Chem. Int. Ed.* **2012**, *51*, 6074–6092. (b) Giustra, Z. X.; Liu, S.-Y. The State of the Art in Azaborine Chemistry: New Synthetic Methods and Applications. *J. Am. Chem. Soc.* **2018**, *140*, 1184–1194. (c) McConnell, C. R.; Liu, S.-Y. Late-stage functionalization of BN-heterocycles. *Chem. Soc. Rev.* **2019**, *48*, 3436–3453.

²² Brough, S. A.; Lamm, A. N.; Liu, S.-Y.; Bettinger, H. F. Photoisomerization of 1,2-Dihydro-1,2-Azaborine: A Matrix Isolation Study. *Angew. Chem. Int. Ed.* **2012**, *51*, 10880–10883.

²³ Edel, K.; Yang, X.; Ishibashi, J. S. A.; Lamm, A. N.; Maichle-Mössmer, C.; Giustra, Z. X.; Liu, S.-Y.; Bettinger, H. F. The Dewar Isomer of 1,2-Dihydro-1,2-azaborinines: Isolation, Fragmentation, and Energy Storage. *Angew. Chem. Int. Ed.* **2018**, *57*, 5296–5300.

²⁴ Richter, R. C.; Biebl, S. M.; Einholz, R.; Walz, J.; Maichle-Mössmer, C.; Ströbele, M.; Bettinger, H. F.; Fleischer, I. Facile Energy Release from Substituted Dewar Isomers of 1,2-Dihydro-1,2-azaborinines Catalyzed by Coinage Metal Lewis Acids. *Angew. Chem. Int. Ed.* **2024**, *63*, e202405818.

²⁵ Giustra, Z. X.; Yang, X.; Chen, M.; Bettinger, H. F.; Liu, S.-Y. Accessing 1,2-Substituted Cyclobutanes through 1,2-Azaborine Photoisomerization. *Angew. Chem. Int. Ed.* **2019**, *58*, 18918–18922.

²⁶ Lin, H.; Yang, X.; Liu, S.-Y.; Jäkle, F. Ring-Opening Metathesis Polymerization of the Dewar Isomer of 1,2-Azaborinine, a B–N Isostere of Benzene. *ACS Macro Lett.* **2024**, *13*, 21–27.

²⁷ Buglioni, L.; Raymenants, F.; Slattery, A.; Zondag, S. D. A.; Noël, T. Technological Innovations in Photochemistry for Organic Synthesis: Flow Chemistry, High-Throughput Experimentation, Scale-up, and Photoelectrochemistry. *Chem. Rev.* **2022**, *122*, 2752–2906.

²⁸ Abbey, E. R.; Zakharov, L. N.; Liu, S.-Y. Crystal Clear Structural Evidence for Electron Delocalization in 1,2-Dihydro-1,2-azaborines. *J. Am. Chem. Soc.* **2008**, *130*, 7250–7252.

²⁹ (a) Shi, Y.; Yang, Y.; Xu, S. Iridium-Catalyzed Enantioselective C(sp³)-H Borylation of Aminocyclopropanes. *Angew. Chem. Int. Ed.* **2022**, *61*, e202201463. (b) Hohn, E.; Pietruszka, J.; Solduga, G. Synthesis of enantiomerically pure cyclopropyl trifluoroborates. *Synlett* **2006**, *10*, 1531–1534.

³⁰ Burford, R. J.; Li, B.; Vasiliu, M.; Dixon, D. A.; Liu, S.-Y. Diels–Alder Reactions of 1,2-Azaborines. *Angew. Chem. Int. Ed.* **2015**, *54*, 7823–7827.

³¹ Szeimien, G.; Schloer, A.; Philipp, F.; Dietz, P.; Mickler, W. Thioadditionen an Bicyclo[1.1.0]butan-Systeme. *Chem. Ber.* **1978**, *111*, 1922–1937.

³² (a) Griesbaum, K. Problems and Possibilities of the Free-Radical Addition of Thiols to Unsaturated Compounds. *Angew. Chem. Int. Ed.* **1970**, *9*, 273–287. (b) Bär, R. M.; Kirschner, S.; Nieger, M.; Bräse, S. Alkyl and Aryl Thiol Addition to [1.1.1]Propellane: Scope and Limitations of a Fast Conjugation Reaction. *Chem. Eur. J.* **2018**, *24*, 1373–1382.

³³ (a) Arpa, E. M.; Stafström, S.; Durbeej, B. Photochemical formation of the elusive Dewar isomers of aromatic systems: why are substituted azaborines different? *Phys. Chem. Chem. Phys.* **2024**, *26*, 11295–11305. (b) Jeong, S.; Park, E.; Kim, J.; Kim, K. H. Ultrafast photoisomerization mechanism of azaborine revealed by nonadiabatic molecular dynamics simulations. *Phys. Chem. Chem. Phys.* **2023**, *25*, 17230–17237.

³⁴ For similar radical 1,2-boron shift, see: (a) Batey, R. A.; Smil, D. V. The First Boron-Tethered Radical Cyclizations and Intramolecular Homolytic Substitutions at Boron. *Angew. Chem. Int. Ed.* **1999**, *38*, 1798–1800. (b) Kaiser, D.; Noble, A.; Fasano, V.; Aggarwal, V. K. 1,2-Boron Shifts of β -Boryl Radicals Generated from Bis-boronic Esters Using Photoredox Catalysis. *J. Am. Chem. Soc.* **2019**, *141*, 14104–14109. (c) Jana, K.; Bhunia, A.; Studer, A. Radical 1,3-difunctionalization of allylboronic esters with concomitant 1,2-boron shift. *Chem* **2020**, *6*, 512–522. (d) Guo, Y.; Wang, X.; Li, C.; Su, J.; Xu, J.; Song, Q. Decarboxylation of β -boryl NHPI esters enables radical 1,2-boron shift for the assembly of versatile organoborons. *Nat. Commun.* **2023**, *14*, 5693.

³⁵ Photoexcitation of Dewar benzene results in the formation of benzene, see: Harman, P. J.; Kent, J. E.; O'Dwyer, M. F.; Griffith, D. W. T. Photochemistry of benzene isomers. 2. Benzvalene and Dewar benzene. *J. Phys. Chem.* **1981**, *85*, 2731–2733.

³⁶ For a computational analysis of the photochemistry of benzene, see: Pieri, E.; Lahana, D.; Chang, A. M.; Aldaz, C. R.; Thompson, K. C.; Martinez, T. J. The non-adiabatic nanoreactor: towards the automated discovery of photochemistry. *Chem. Sci.* **2021**, *12*, 7294–7307.