

Stereoselective Synthesis of Mirogabalin via 1,4-Selective Addition of Lithioacetonitrile to Alkylidene Malonate

Hidenori Ochiai,* Taiki Mihara, Miwa Sasagawa, Akira Nishiyama

Pharma & Supplemental Nutrition Solutions Vehicle, Pharma Business Division, Kaneka Corporation, 1-8, Miyamae-cho, Takasago-cho, Takasago, Hyogo 676-8688, Japan

KEYWORDS. Mirogabalin, lithioacetonitrile, 1,4-addition, alkylidene malonate, Hofmann rearrangement

ABSTRACT: The synthesis of mirogabalin was studied for industrial production, and an alternative to the Daiichi–Sankyo’s method was established. The developed synthesis involves the introduction of a two-carbon unit with the stereoselective 1,4-selective addition of lithioacetonitrile to alkylidene malonate and one-carbon degradation by Hofmann rearrangement. The precursor for the Hofmann rearrangement was readily prepared from the 1,4-adduct via a one-pot reaction involving decarboxylation, hydrolysis, and hydration.

INTRODUCTION

Mirogabalin is a class of gabapentinoids originally developed by Daiichi–Sankyo Co., Ltd. These gabapentinoids function as an $\alpha 2\delta$ ligand and are derivatives of the inhibitory neurotransmitter γ -aminobutyric acid (GABA).¹ Since its approval in Japan in 2019, mirogabalin has been marketed as a drug for the treatment of neuropathic pain under the trade name Tarlige. Other well-known gabapentinoids are pregabalin and gabapentin, which are increasingly used worldwide (Figure 1).




Figure 1. Structures of gabapentinoids.


From the structural viewpoint, this compound has a highly strained bicyclo[3.2.0]heptane skeleton and three contiguous stereogenic centers despite its small molecular size, making it unique and extremely difficult to synthesize. Therefore, a sophisticated design strategy is necessary for synthesizing mirogabalin to ensure a robust commercial supply.

The original synthetic route for mirogabalin developed by Daiichi–Sankyo is shown in Scheme 1.¹ The optically active bicyclo ketone (–)-**2** was readily derived by separating its racemic form (*rac*)-**2** by various optical resolution methods (i.e., enzymatic,² salt crystallization,³ and asymmetric aldol methods⁴), and the requisite (*rac*)-**2** was readily synthesized via Claisen rearrangement⁵ or Aza-Claisen rearrangement.⁶ (–)-**2** was transformed to diethyl cyclobutylidene malonate **3** through Knoevenagel condensation. Reportedly, the undesired precipitation of insoluble Ti complexes was suppressed by

conducting the reaction in cyclopentyl methyl ether (CPME).¹ Furthermore, reports have stated that the use of TiCl₃(*Oi*-Pr), generated in situ by mixing TiCl₄ and Ti(*Oi*-Pr)₄ in a 3:1 ratio, could adjust the Lewis acidity and suppress the decomposition of **2**, while furnishing **3** in good yield. 1,4-Addition of a cyano group from the convex face of **3** yielded the 1,4-adduct **4** with perfect stereoselectivity. Aqueous KOH was slowly added to an ethanolic solution of **4**, while heating to induce decarboxylation of the ethyl malonate unit to afford **5**. Further addition of KOH solution completed the hydrolysis to furnish **6** in a one-pot manner. The resulting **6** was solidified with benzylamine via salt formation to obtain **7**. Subsequently, after the dissolution of salt **7** with LiOH, the nitrile moiety of the resulting **7** (free form) was reduced by hydrogenation under high-pressure conditions using sponge cobalt to obtain the corresponding amine moiety. After neutralizing the γ -amino acid with malonic acid, **1** (free form) was isolated as crystals. Finally, salt crystallization of **1** (free form) with benzenesulfonic acid (besylic acid) afforded mirogabalin besylate **1** in good yield with good purity.

Given the interesting and unique structure of **1**, we have also gained our interest and embarked on its process development, aiming for a robust commercial supply of **1**. Prior to the study, issues to be addressed based on Daiichi–Sankyo’s original synthetic route were identified by considering scale-up. The generation of toxic HCN gas during the introduction of the cyano group was a concern, where the amount of generated gas needed to be controlled. Controlling the residual heavy metals during reduction with pyrophoric sponge cobalt was also necessary.⁷ Furthermore, pressure-resistant equipment is needed for nitrile reduction to achieve efficient hydrogenation.⁸ All of these requirements arise from the use of inorganic NaCN as a nucleophile for constructing the asymmetric quaternary center.


Scheme 1. Original synthetic route (Daiichi-Sankyo, Co., Ltd.)



RESULTS AND DISCUSSION

Based on the above background, we envisioned that these matters of concern, including controlling HCN gas, the use of a pyrophoric metal, controlling the residual metal, and facility limitations, could be avoided in a single stroke by changing the strategy to introduce a two-carbon unit bearing a carbamoyl group to obtain **8**, followed by one-carbon degradation (Scheme 2, top), instead of introducing a cyano group followed by the reduction of the nitrile (Scheme 2, bottom).

Scheme 2. Working hypothesis of this study



For the one-carbon degradation reaction, the Hofmann rearrangement is one of the most inexpensive and commercially effective methods. There are many similar examples of the derivation from 4-carbamoylbutyric acids to γ -amino acids using this reaction; therefore, this transformation is considered promising.⁹


Based on this working hypothesis, an alternative to the original Daiichi-Sankyo's synthesis was established, which is explained in detail below. The synthetic route is summarized in Scheme S1 (Supporting Information).

The Knoevenagel condensation product **3** was synthesized with (*rac*)-**2** according to a method published by Daiichi-Sankyo¹ and a patent description (Scheme S2).⁸

First, various malonate-type derivatives were investigated as soft-carbon nucleophiles for the construction of quaternary centers via 1,4-addition (Scheme 3). Ethyl cyanoacetate afforded a complex mixture; however, **9a** was not obtained. When Meldrum's acid was used to generate **9b**, the reaction did not proceed and the starting material was completely recovered. When diethyl malonate was

used, the target product **9c** was obtained in moderate yield; however, a high yield could not be achieved despite modification of the reaction conditions. 1,4-Addition of soft nucleophiles to **3** did not proceed efficiently because of steric hindrance near the newly formed quaternary center and undesired competition from the reverse reaction of the malonate as a good leaving group.

Scheme 3. 1,4-Addition of soft nucleophiles



Therefore, a monoester derivative was considered as an alternative nucleophile to malonate, aiming for relaxation of the steric hindrance and suppression of reverse reactions, although a stronger base is required for deprotonation. Incidentally, to carry out the Hofmann rearrangement in the later stage of synthesis, a carbamoyl moiety is required. Herein, acetonitrile was considered the best nucleophile among the monoester derivatives, providing easy access to the carbamoyl moiety in a few steps because it already has a nitrogen atom in its chemical structure. Importantly, because the anion species generated from acetonitrile is a relatively hard nucleophile, the competitive 1,2-addition, rather than 1,4-addition, was a concern,¹⁰ and the reactivity of this species has not been well investigated.


The results of the optimization study are shown in Tables 1 and S1–6. Upon treatment with strong bases such as lithium diisopropylamide (LDA), lithium tetramethylpiperidide (LiTMP), and lithium hexamethyldisilazide (LHMDS) under cryogenic conditions at $-78\text{ }^\circ\text{C}$, the 1,4-addition proceeded smoothly from the convex face of **3** to afford the desired **10** in high yields (entries 1–3; Table S1), in which the competitive 1,2-addition was not observed, intriguingly, the same reaction was achieved using *n*-BuLi instead of lithium amide (entry 4); to a suspension of *n*-BuLi (1.5 M in hexane, 1.5 equiv.) and MeCN (1.5 equiv.) in THF (10 V), **3** (1 equiv.) was added at $-78\text{ }^\circ\text{C}$, followed by stirring at $25\text{ }^\circ\text{C}$

for 1 h to obtain the desired **10** in 97.2% yield (assayed by HPLC). This result suggests that lithioacetonitrile was generated efficiently without the undesired nucleophilic addition¹¹ to acetonitrile or without dimerization¹² under these conditions.¹³ Interestingly, when NaH was used, the desired product was not obtained (entry 5). In this case, **3** was recovered, and the active species were deactivated. To the best of our knowledge, the 1,4-selective addition of lithioacetonitrile to β,β -dialkyl-substituted alkylidene malonates, as in this case, is not well known. At this stage, it is difficult to conclude whether the reaction is specific to mirogabalin¹⁴, or whether it can be generalized.

From the viewpoint of material cost, the reaction conditions in entry 4 are highly attractive and were considered provisionally optimal. Further investigations of the reaction conditions were conducted. The effect of solvents were investigated. (entries 6–10; Table S2). Interestingly, significant differences were observed in ether-based solvents. When cyclopentyl methyl ether (CPME) or 4-methyltetrahydropyran (MTHP) was used, the reaction proceeded well and furnished the target product in good yields (entries 6 and 7); however, when methyl *tert*-butyl ether (MTBE) was used, a significant decrease in the yield was observed (entry 8). Low-polarity solvents such as toluene and *n*-hexane significantly reduced the yield and gave complex mixtures (entries 9 and 10). The effect of reducing the amount of base was also examined (entries 11 and 12; Table S3). The reduction of *n*-BuLi to 1.2 equiv. reduced the yield to 89.1% (entry 11), whereas upon further reduction to 1.0 equiv, the reaction did not proceed to completion and the yield was reduced to 76.0% (entry 12). The concentrations of the reactants were

also studied (entries 13 and 14; Table S4). When the solvent volume was reduced to 5 V, the product yield remained unchanged (entry 13). When the solvent volume was further reduced to 3 V, the product yield decreased slightly (entry 14). The reaction temperature was also examined (entries 15–17; Table S5). The reaction proceeded smoothly even at -78 °C, suggesting that the 1,4-addition reaction proceeds rapidly (entry 15). Increasing the temperature to -40 °C gave the desired product in relatively good yield (entry 16), whereas the reaction at -20 °C did not give the desired product (entry 17). In this case, the ester moiety was damaged. Because there is an example in which lithioacetonitrile was added at approximately room temperature,^{12a,15} we were encouraged to re-examine the protocol at a temperature at which commercial production is possible (entries 18–21; Table S6). After examining the procedure for reagent addition, it was found that adding *n*-BuLi to a solution of **3** and MeCN in THF at -20 °C gave the desired product in 75.4% yield (entry 18). The amounts of *n*-BuLi and MeCN were increased to improve the yield; however, contrary to expectations, the reaction yield decreased (entry 19). Because the ratio of MeCN to *n*-BuLi seemed important, the amount of MeCN was further increased to 16 equiv, which yielded the desired product in 92.9% yield; this suggested that the deprotonation of MeCN was promoted (entry 20). The product was obtained even at 0 °C, albeit in moderate yield (entry 21). Notably, almost no diastereomers were observed and the diastereoselectivity was extremely high in every case, which is similar to the effect of adding NaCN to **3**, as reported by Daiichi-Sankyo.¹

Table 1. Optimization of 1,4-addition



entry	Base (X equiv.)	Solvent (Y V)	temp. (°C)	Conv. (%) ^a	3 (%) ^b	10 (%) ^b	<i>epi-10</i> (%) ^b
1 ^c	LDA (1.5)	THF (10 V)	-78 to 25	99.8	0.2	95.3	1.8
2 ^d	LiTMP (1.5)	THF (10 V)	-78 to 25	99.7	0.2	95.9	1.9
3	LHMDS (1.5)	THF (10 V)	-78 to 25	82.8	7.7	80.0	0.0
4	<i>n</i>-BuLi (1.5)	THF (10 V)	-78 to 25	98.6	0.6	97.2	1.0
5	NaH (1.5)	THF (10 V)	-78 to 25	0.0	54.6	0.0	0.0
6	<i>n</i> -BuLi (1.5)	CPME (10 V)	-78 to 25	97.0	2.2	72.1	0.1
7	<i>n</i> -BuLi (1.5)	MTHP (10 V)	-78 to 25	97.9	1.9	89.5	0.6
8	<i>n</i> -BuLi (1.5)	MTBE (10 V)	-78 to 25	62.4	14.4	24.2	0.5
9	<i>n</i> -BuLi (1.5)	toluene (10 V)	-78 to 25	72.1	5.8	15.0	0.0
10	<i>n</i> -BuLi (1.5)	hexane (10 V)	-78 to 25	18.4	11.1	2.5	0.0
11	<i>n</i> -BuLi (1.2)	THF (10 V)	-78 to 25	99.3	0.3	89.1	0.9
12	<i>n</i> -BuLi (1.0)	THF (10 V)	-78 to 25	77.4	10.2	76.0	0.5
13	<i>n</i> -BuLi (1.5)	THF (5 V)	-78 to 25	99.0	0.5	97.1	0.5
14	<i>n</i> -BuLi (1.5)	THF (3 V)	-78 to 25	98.9	0.5	91.3	0.4
15	<i>n</i> -BuLi (1.5)	THF (10 V)	-78	99.5	0.2	95.6	0.6
16	<i>n</i> -BuLi (1.5)	THF (10 V)	-40 to 25	98.9	0.4	89.3	1.0
17	<i>n</i> -BuLi (1.5)	THF (10 V)	-20 to 25	100	0.0	0.4	0.2
18 ^e	<i>n</i> -BuLi (1.5)	THF (10 V)	-20 to 25	87.0	5.5	75.4	1.2
19 ^{e,f,h}	<i>n</i> -BuLi (2.0)	THF (10 V)	-20 to 25	91.4	6.5	67.8	1.4
20^{e,g,h}	<i>n</i>-BuLi (2.0)	THF (10 V)	-20 to 25	97.3	2.6	92.9	1.0
21 ^{e,f}	<i>n</i> -BuLi (2.0)	THF (10 V)	0 to 25	82.4	7.5	73.3	1.5

^aDetermined by HPLC [(area¹⁰ + area^{*epi-10*})/(area³ + area¹⁰ + area^{*epi-10*})]

^bQuantified using external standard method

^cLDA (1.5 equiv) was prepared *in situ* from diisopropylamine (1.5 equiv) and *n*-BuLi (1.5 equiv) at -78 °C for 30 min.

^dLiTMP (1.5 equiv) was prepared *in situ* from 2,2,6,6-tetramethylpiperidine (1.5 equiv) and *n*-BuLi (1.5 equiv) at -78 °C for 30 min

^e*n*-BuLi was added to the solution containing **3** and MeCN in THF

^fMeCN (2.0 equiv) was used instead of MeCN (1.5 equiv). ^gMeCN (16.0 equiv.) was used instead of MeCN (1.5 equiv)

^hReaction time was 30 min

Encouraged by the successful 1,4-selective addition of lithioacetonitrile to **3** with good yield and stereoselectivity, which was achieved using only inexpensive *n*-BuLi and MeCN, we focused

on the decarboxylation and hydrolysis of **10** to **13** (Tables 2 and S7).

Table 2. Optimization of one-pot process (decarboxylation and hydrolysis)

entry	Base (X equiv.)	Base solution in dropping funnel	EtOH total (Y V)	10 (%) ^a	11 (%) ^a	12 (%) ^a	13 (%) ^a	14 (%) ^a	12+13 (%)
1	KOH (1.5)	45% aq. solution	9	0.0	2.1	20.1	69.4	11.5	89.5
2 ^b	KOH (1.5)	EtOH (4 V)	9	0.0	0.0	16.4	74.3	9.2	90.7
3 ^c	LiOH (1.5)	insoluble	9	42.4	3.2	52.4	0.0	0.0	52.4
4 ^b	NaOH (1.5)	EtOH (4 V)	9	7.2	0.0	16.6	73.6	2.9	90.2
5 ^b	CsOH·H ₂ O (1.5)	EtOH (4 V)	9	0.0	0.0	0.0	84.2	15.3	84.2
6 ^c	KOH (1.5)	EtOH (4 V)	20	0.0	0.0	5.8	88.9	8.0	94.7
7 ^d	KOH (1.2)	EtOH (4 V)	20	0.0	2.3	49.9	49.2	3.8	99.1

^aQuantified using external standard method

^bBase in EtOH (4 V) was added to a solution of **10** in EtOH (5 V). Total volume of EtOH was 9 V

^cLiOH was added to a solution of **10** in EtOH (9 V)

^dKOH in EtOH (4 V) was added to a solution of **10** in EtOH (16 V) Total volume of EtOH was 20 V

Referring to the report by Daiichi-Sankyo,¹ a 45% aqueous KOH solution was added to a solution of **10** in EtOH (9 V) over 4 h under reflux conditions (entry 1). Hydrolysis and decarboxylation occurred simultaneously, yielding a mixture of monoester **12** (20.1%) and the target carboxylic acid **13** (69.4%). In this case, 11.5% of dicarboxylic acid **14** was generated as a byproduct, but this compound was not decarboxylated under the reaction conditions and was not converted to **12**.¹⁶ Compounding the issue, **14** was difficult to remove by liquid separation or crystallization, which led to lower purity and lower yields of the products, including **12** and **13**. Therefore, the generation of **14** was minimized by strictly controlling the reaction conditions.

To suppress the generation of the undesired byproduct dicarboxylic acid **14**, the amount of H₂O was first reduced (entry 2); a KOH solution in dehydrated EtOH (4 V, 5.6 wt% solution) was used instead of the original 45% KOH aqueous solution. This solution was then added to a solution of **10** in EtOH (5 V) over 4 h (the total EtOH volume was 9 V). In this case, the production of **14** was slightly suppressed.


Bases were then examined (entries 2–5). When less basic LiOH or NaOH was used, the generation of byproduct **14**, formed by the hydrolysis of **11**, was suppressed. The hydrolysis of **10** to **11** was also suppressed, and the net reaction was dramatically delayed (entries 3 and 4). When excess base was added to complete the reaction, a non-negligible amount of **14** was formed (data not shown). CsOH·H₂O as a stronger base was found to increase the formation of **14** (entry 5).

Therefore, to slow the intermolecular conversion of **11** to **14** and relatively accelerate the intramolecular conversion of **11** to **12**, the reaction mixture was diluted by increasing the amount of total EtOH from 9 to 20 V, which slightly improved the reaction yield,

and the formation of **14** was reduced as expected (entry 6). From this point, we further reduced the amount of base to 1.2 equiv, which successfully minimized the generation of **14** to 3.8%, while allowing **10** to be completely consumed (entry 7). The resulting mixture of **12** and **13** at 25 °C was fully converted to carboxylic acid **13** by further addition of KOH to the solution.

Subsequently, the hydration reactions of nitriles targeting amide **8** were studied separately.


Scheme 4. Hydration of nitrile on **13**



Several reaction conditions have been reported for nitrile hydration.¹⁷ Under acidic conditions, the olefin moiety of **13** was damaged prior to hydration, whereas under basic conditions, the resulting carbamoyl group underwent overhydrolysis, yielding a non-negligible amount of symmetric dicarboxylic acid. Undesired overhy-


hydrolysis could not be suppressed despite several efforts. Gratifyingly, the hydration of nitriles could be induced in high yields under mild conditions with the Radziszewskii reaction,¹⁸ in which aqueous hydrogen peroxide was added under basic conditions. Hydrolysis of ester **12**, followed by hydration of the resulting **13**, proceeded smoothly to furnish **8** in good yield in a one-pot manner (Schemes 4 and S7).

Scheme 5. One-pot transformation




With **8** in hand, which was readily obtained from **10** via a one-pot transformation, the Hoffman rearrangement was subsequently investigated (Schemes 6 and S9). One of the least expensive reagents, $\text{NaClO} \cdot 5\text{H}_2\text{O}$, was used as the oxidant. The reaction proceeded smoothly under the standard reaction conditions. After completion of the reaction, hydrochloric acid was added to adjust the pH to 6. Neutralization and crystallization proceeded, and γ -amino acids were precipitated from the aqueous solution and isolated as a solid in high yield. Through this solidification process, both the purity and content of the resulting solid were improved compared to those of crude **8**. Notably, the isolation of **1** (free form) as a solid is required only once, which simplifies the process and reduces the operating time.

Scheme 6. Hofmann rearrangement



The remaining step was the salt crystallization of **1** (free form) (Schemes 7 and S10). As in Daiichi-Sankyo's patent,⁸ besylic acid was added to a suspension of **1** (free form) in anisole to obtain target product **1** with good purity.

Scheme 7. Salt formation with besylic acid




The transformations, including decarboxylation, hydrolysis, and hydration, were also conducted and examined in a one-pot process (Schemes 5 and S8). After the decarboxylation, the solvent was exchanged from EtOH to water to facilitate liquid separation. Subsequently, hydrolysis was conducted in one pot, yielding **13** in 98.4% assayed yield. Aqueous H_2O_2 was added to this basic aqueous solution to afford **8** in 92.8% assay yield. After workup, a residue containing **8** was obtained in 84.4% yield as an oil, which was used for the next step without further purification.

DISCUSSION OF 1,4-SELECTIVE ADDITION

The key to the developed synthesis of mirogabalin was the successful 1,4-selective addition of lithioacetonitrile to alkylidene malonates. As mentioned above, lithioacetonitrile seems to be a relatively hard nucleophile, and 1,2-additions are expected to be preferential; however, 1,4-selective addition occurred even under kinetic reaction conditions. To ascertain whether this phenomenon is specific to mirogabalin or can be generalized, various electrophiles were explored to investigate the selectivity and discuss its origin (Tables 3 and S8).

Table 3. Investigation of selectivity



entry	15	16 (%) ^a	17 (%) ^a	Others
1	15a	79	ND	
2	15b	ND	87	
3	15c	13	54	olefin isomer: 14%
4	15d	ND	93	
5	15e	73	ND	

^aIsolated yield.

^b“ND” stands for “not detected”.

When the reaction conditions (Table 1, entry 4) were employed for cinnamic acid ester **15a**, only the 1,2-adduct **16a** was obtained; the 1,4-adduct **17a** was not observed (entry 1). On the other hand, quite interestingly, when applied to benzylidene malonate **15b**, only the 1,4-adduct **17b** was obtained in 87% yield, suggesting that the activation of the double bond by two ester groups accelerates the 1,4-addition reaction (entry 2). A competition experiment between **15a** and **15b** also supported this observation.¹⁹


When the reaction conditions were employed for isopropylidene malonate **15c** bearing the β,β -dialkylsubstituted olefin, 1,2-addition and 1,4-addition competed, with isomerization of the olefin occurring in addition to these reactions (entry 3; Table S8). With **15d**, which is ethylidene malonate with less steric hindrance around the reaction site, only the 1,4-adduct **17d** was obtained in high yield (entry 4). With **15e** bearing a methyl group at the α -position, only the 1,2-addition product **16e** was observed in spite of the steric hindrance around the reaction site, suggesting that the second ester behaves as an activator of the double bond and does not play the role of a bulky substituent (entry 5).

Based on the above, the 1,4-selective addition of **3** proceeds with high efficiency, mainly owing to the activation of the olefins by the

two ester groups. The fact that olefin isomerization does not occur because of the ring strain of the cyclobutene ring and that the ring strain is partly released by the transformation from sp^2 to sp^3 -hybridization also contributes to the high efficiency.


Other than malonate addition to unsaturated nitriles **21** (Scheme 8, path b), this 1,4-selective addition of lithioacetonitrile to alkylidene malonate is expected to provide a cheaper alternative for accessing cyanoethylmalonate derivatives **19**, which are important intermediates for synthesizing γ -amino acid derivatives (Scheme 8, path a).

Scheme 8. Alternative retrosynthesis




Scheme 9. Summary of the developed synthetic route

Our synthetic route



Daiichi-Sankyo's original route



CONCLUSION

An alternative to Daiichi-Sankyo's method was established for the commercial production of mirogabalin (Scheme 9). The developed synthesis involves the introduction of a two-carbon unit, followed by one-carbon degradation. The method features the introduction of a two-carbon unit by the stereo- and 1,4-selective addition of lithioacetonitrile to alkylidene malonate, which can be generated from inexpensive *n*-BuLi and acetonitrile. This novel transformation potentially broadens the scope of retrosynthesis of GABA derivatives. The one-pot transformation, including decarboxylation, hydrolysis, and hydration, enables easy access to the precursor of the Hofmann rearrangement.

EXPERIMENTAL SECTION

General Information

All reactions were performed under a nitrogen atmosphere unless otherwise indicated. All manipulations of air- and/or moisture-sensitive compounds were performed using standard Schlenk techniques. ¹H NMR (500 MHz) and ¹³C NMR (125 MHz) spectra were obtained from measurements at ambient temperature on a JEOL

500SS spectrometer (JNM-ECA500). Chloroform-*d*₁ (CDCl₃) containing 0.03% tetramethylsilane (TMS) (99.8%D, KANTO Chemical Co., Inc.) was used as a solvent for NMR measurements at room temperature. Chemical shifts (δ) are given in parts per million (ppm) downfield from the signal of TMS in CDCl₃ (δ 0.00 ppm for ¹H NMR) or CDCl₃ (δ 77.0 ppm for ¹³C NMR) as an internal standard, and as an external standard with coupling constants (*J*) in hertz (Hz). Abbreviations s, d, t, q, sep, m, and br signify singlet, doublet, triplet, quartet, septet, multiplet, and broad, respectively.

HPLC Method

High performance liquid chromatography (HPLC) was performed by LC-2050C Plus with 5C18-AR-II column (250 \times 4.6 mm I.D., 5.0 μ m). The column oven was set at 40 $^{\circ}$ C. The UV detector was set at 210 nm. Mobile phases A (0.1% phosphoric acid aq.) and B (acetonitrile) were utilized at a flow rate of 1.4 mL/min. Mobile phase B was increased linearly from 50% to 70% over 3.2 min, held at 70% for 6 min, increased linearly from 70% to 100% over 4.8 min, and held at 100% for 4.8 min.

Retention times: **2** (5.19 min), **3** (9.56 min), **10** (9.20 min), *epi*-**10** (8.97 min), **12** (7.58 min), **13** (4.32 min), **14** (3.26 min), **8** (2.91 min), **1** (1.58 min), BsOH (2.39 min).

Preparation for **3** (Knoevenagel Condensation)

To a stirred solution of $\text{Ti}(\text{O}i\text{-Pr})_4$ (3.50 mL, 11.8 mmol, 0.40 equiv.(gross basis)) in CPME (32.0 mL, 8.0 V (gross basis)) was added TiCl_4 (3.52 mL, 32.3 mmol, 1.1 equiv.(gross basis)) dropwise at 20 °C. After stirring for 30 min at 10 °C, diethyl malonate (4.93 mL, 32.3 mmol, 1.1 equiv.(gross basis)) was added at the same temperature. After stirring at 5 °C for 15 min, **2** (gross: 4.00 g, net: 3.96 g, content: 99.0 wt%, purity: 98.8 area%, 29.4 mmol, 1 equiv.(basis)) was added at the same temperature. After stirring at 30 °C for 3 h [conversion: 93.7%, yield: 95.8% (HPLC)], to the reaction mixture was added H_2O (20 mL, 5.0 V (gross basis)) at 5 °C. The organic phase was separated and washed by 3.8% HCl (8 mL, 2.0 V (gross basis)) and 3.0% NaHCO_3 (8 mL, 2.0 V (gross basis)). The organic phase was concentrated under reduced pressure and dried in vacuo at 25 °C for 18 h to afford **3** (gross: 8.07 g, net: 7.59 g, content: 92.4 wt%, purity: 94.5 area%, 27.3 mmol, 93.8% yield) as reddish oil.

Preparation for **10** (1,4-addition of lithioacetonitrile)

To a solution of *n*-BuLi (1.6 M in hexane, 11.2 mL, 18.0 mmol, 1.5 equiv.(net basis)) in THF (36 mL, 10 V (gross basis)) was added MeCN (0.937 mL, 18.0 mmol, 1.5 equiv.(net basis)) at -78 °C. After stirring for 30 min, **3** (gross: 3.60 g, net: 3.33 g, content: 92.4 wt%, purity: 94.5 area%, 12.0 mmol, 1 equiv.(basis)) was added dropwise at the same temperature. After stirring at 25 °C for 1 h [conversion: 100%, yield: 102.6% (HPLC)], to the reaction mixture was added sat. NH_4Cl aq. (18 mL, 5 V (gross basis)) at the same temperature. The organic phase was separated. The aqueous layer was extracted with CH_2Cl_2 (18 mL, 5 V (gross basis)) twice. The combined organic phases were concentrated under reduced pressure and dried in vacuo at 25 °C for 18 h to afford **10** (gross: 4.17 g, net: 3.57 g, content: 85.6 wt%, purity: 89.7 area%, 1.47 mmol, 91.1% yield.) as reddish oil.

Preparation for **8** (one-pot transformation)

To a solution of **10** (gross: 4.05 g, net: 3.47 g, content: 85.6 wt%, purity: 89.7 area%, 10.9 mmol, 1 equiv.) in EtOH (55.5 mL, 16 V (net basis)) was added KOH (5.6 wt% solution, 730 mg, 13.0 mmol, 1.2 equiv.) in EtOH (15.6 mL, 4 V (net basis)) at reflux temperature over 4 h. After stirring for 72 h [yields: **10** (0.0%), **12** (39.9%), **13** (62.7%) (HPLC)], the reaction mixture was cooled to 25 °C. To the mixture was added H_2O (13.9 mL, 4 V (net basis)) at the same temperature. The reaction mixture was concentrated under reduced pressure (30 °C, 50 mmHg) to afford an aqueous solution (total weight: 24.4 g). To the mixture was added aq. KOH (20% solution, 1.22 g, 21.7 mmol, 2.0 equiv.) at the same temperature. After stirring at the same temperature for 3 h [yields: **12** (0.0%), **13** (98.4%)], the reaction mixture was cooled to 0 °C. To the mixture was added aq. H_2O_2 (30% solution, 2.58 g, 76.0 mmol, 7.0 equiv.) at the same temperature. After stirring at 60 °C for 3 h [yields: **13** (1.0%), **8** (92.8%) (HPLC)], the reaction mixture was cooled to 25 °C. To this mixture was added toluene (20.3 mL, 5 V (gross basis)) at the same temperature. The phases were separated. To the aqueous phase was added HCl aq. (17.5% solution, 2.35 mL) at room temperature. The aqueous solution was extracted by EtOAc (40.5 mL, 10 V (gross basis)) three times. The combined organic phases were concentrated under reduced pressure and dried in vacuo at 25 °C for 18 h to afford **8** (gross: 3.20 g, net: 2.17 g, content: 67.5 wt%, purity: 81.7 area%, 12.5 mmol, 84.4% yield) as a viscous yellow oil.

Preparation for **1** (free form) (Hofmann rearrangement)

To a stirred suspension of **8** (gross: 1.00 g, net: 675 mg, content: 67.5 wt%, purity: 81.7 area%, 2.85 mmol, 1 equiv.(basis)) in H_2O (3 mL, 3 V (gross basis)) were added NaOH aq. (20% in H_2O , 8.53 mmol, 3.0 equiv. (net basis)) and $\text{NaClO} \cdot 5\text{H}_2\text{O}$ (561 mg, 3.41 mmol, 1.2 equiv. (net basis)) at 0 °C. After stirring for 1 h at the same temperature, the reaction mixture was warmed up to 25 °C. After stirring for 3 h at the same temperature [(conversion: 100%, yield: 92.4% (HPLC)], the reaction mixture was cooled to 0 °C. To

the mixture was added HCl (17.5 wt% in H_2O , 1.78 g) until reaching pH6 to afford precipitation. Then NaCl (1.04 g, 17.8 mmol) was added to the reaction mixture. After stirring at the same temperature for 1 h, the resulting slurry solution was filtered off. The resulting wet crystal was washed with iced water (2 mL, 2 V (gross basis)) twice, and dried in vacuo at 40 °C for 20 h to afford **1** (free form) as a white crystal (gross: 483 mg, net: 465 mg, content: 96.2 wt%, purity: 88.8 area%, 2.22 mmol, 78.4% yield).

Preparation for **1** (Salt formation)

To a suspension of **1** (free form) (gross: 100 mg, net: 96.2 mg, content: 96.2 wt%, purity: 88.8 area%, 0.460 mmol, 1 equiv.(basis)) in anisole (4.0 mL, 40 V (gross basis)) was added a solution of besylic acid (79.5 mg, 0.483 mmol, 1.05 equiv.(net basis)) in anisole (0.40 mL, 4.0 V (gross basis)) at 25 °C. The reaction mixture was cooled to 0 °C. After stirring at the same temperature for 1 h, the resulting suspension was filtered off. The wet crystal was washed with acetone (0.50 mL, 5.0 V (gross basis)) once and dried in vacuo at 40 °C for 20 h to afford **1** as a white solid (gross: 111 mg, net: 113 mg, content: 101.7 wt%, purity: 99.1 area%, 0.303 mmol, 65.9% yield).

General Procedure for 1,4-addition of lithioacetonitrile

To a solution of *n*-BuLi (1.5 M in *n*-hexane, 5.00 mL, 7.50 mmol, 1.5 equiv.) in THF (10 mL, 0.5 M) was added MeCN (0.392 mL, 7.50 mmol, 1.5 equiv.) dropwise at -78 °C. After stirring at the same temperature for 30 min, **15** (5.00 mmol, 1 equiv.(basis)) was added dropwise at the same temperature. The reaction mixture was warmed up to 25 °C. After stirring at the same temperature for 1 h, 1 M HCl (10 mL) was added to the mixture. The mixture was extracted by EtOAc (10 mL) three times. The combined organic phase was dried over Na_2SO_4 , and filtered. The resulting organic phase was concentrated under the reduced pressure to afford a residue. The residue was purified by silica-gel column chromatography to afford the products.

Characterization

(*1R,5S*)-3-Ethylbicyclo[3.2.0]hept-3-en-6-one (**2**). Colorless oil; ^1H NMR (CDCl_3) δ 5.24–5.22 (m, 1H), 4.21–4.18 (m, 1H), 3.24–3.17 (m, 1H), 2.85–2.76 (m, 3H), 2.35–2.29 (m, 1H), 2.14 (q, J = 7.5 Hz, 2H), 1.07 (t, J = 7.5 Hz, 3H); ^{13}C NMR (CDCl_3) δ 209.1 (1C), 150.1 (1C), 117.1 (1CH), 73.3 (1CH), 53.1 (1CH₂), 42.7 (1CH₂), 26.1 (1CH), 24.2 (1CH₂), 12.2 (1CH₃).

Diethyl 2-((*1R,5S*)-3-ethylbicyclo[3.2.0]hept-3-en-6-ylidene)malonate (**3**). Colorless oil; ^1H NMR (CDCl_3) δ 5.37–5.35 (m, 1H), 4.29–4.23 (m, 1H), 4.26 (qd, J = 7.2, 1.4 Hz, 2H), 4.21 (qd, J = 7.2, 0.9 Hz, 2H), 3.32 (ddd, J = 18.9, 8.6, 3.7 Hz, 1H), 2.95–2.89 (m, 1H), 2.72 (ddd, J = 18.9, 5.8, 3.7 Hz, 1H), 2.66 (dd, J = 16.8, 8.2 Hz, 1H), 2.22 (dd, J = 16.8, 3.4 Hz, 1H), 2.11 (q, J = 7.5 Hz, 2H), 1.33 (t, J = 7.2 Hz, 3H), 1.28 (t, J = 7.2 Hz, 3H), 1.07 (t, J = 7.5 Hz, 3H); ^{13}C NMR (CDCl_3) δ 176.1 (1C), 164.5 (1C), 164.2 (1C), 148.8 (1C), 121.0 (1CH), 118.2 (1C), 60.6 (1CH₂), 60.6 (1CH₂), 58.3 (1CH), 42.7 (1CH₂), 39.7 (1CH₂), 32.7 (1CH), 24.3 (1CH₂), 14.22 (1CH₃), 14.20 (1CH₃), 12.2 (1CH₃).

1,3-Diethyl 2-[(*1R,5S,6R*)-(6-cyanomethyl)-3-ethylbicyclo[3.2.0]hept-3-en-6-yl]propanedioate (**10**). Colorless oil; ^1H NMR (CDCl_3) δ 5.28–5.26 (m, 1H), 4.22 (qd, J = 7.2, 3.4 Hz, 2H), 4.28 (q, J = 7.2 Hz, 2H), 3.71 (s, 1H), 3.30 (br s, 1H), 3.08–3.02 (m, 1H), 3.01 (d, J = 2.9 Hz, 2H), 2.54 (dd, J = 16.6, 8.0 Hz, 1H), 2.31 (ddd, J = 13.0, 8.9, 2.7 Hz, 1H), 2.19–2.12 (m, 2H), 2.07 (d, J = 16.6 Hz, 1H), 1.70 (dd, J = 13.0, 7.5 Hz, 1H), 1.29 (t, J = 7.0 Hz, 3H), 1.26 (t, J = 7.0 Hz, 3H), 1.09 (t, J = 7.5 Hz, 3H); ^{13}C NMR (CDCl_3) δ 168.0 (1C), 167.8 (1C), 151.8 (1C), 121.0 (1CH), 118.9 (1C), 61.52 (1CH₂), 61.47 (1CH₂), 55.7 (1CH), 55.1 (1CH), 43.8 (1C), 42.6 (1CH₂), 37.9 (1CH₂), 31.4 (1CH), 24.5 (1CH₂), 24.4 (1CH₂), 14.01 (1CH₃), 13.99 (1CH₃), 12.4 (1CH₃).

1,3-Diethyl 2-[(1R,5S,6S)-(6-cyanomethyl)-3-ethylbicyclo[3.2.0]hept-3-en-6-yl]propanedioate (3:2 mixture of diastereomixture, **epi-10** is a minor product) (**epi-10**). Colorless oil; ¹H NMR (CDCl₃) (minor) δ 5.39–5.38 (m, 1H), 4.27–4.21 (m, 4H), 3.80 (s, 1H), 3.35 (br s, 1H), 2.80 (s, 2H), 2.76–2.71 (m, 1H), 2.55–2.48 (m, 2H), 2.19–2.12 (m, 3H), 1.66 (dd, *J* = 13.2, 6.8 Hz, 1H), 1.32–1.27 (m, 6H), 1.10 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CDCl₃) (minor) δ 167.9 (1C), 166.6 (1C), 152.1 (1C), 120.5 (1CH), 118.4 (1C), 61.7 (1CH₂), 61.6 (1CH₂), 57.4 (1CH), 53.4 (1CH), 43.7 (1C), 41.7 (1CH₂), 36.6 (1CH₂), 30.5 (1CH), 24.4 (1CH₂), 22.5 (1CH₂), 14.01 (1CH₃), 13.99 (1CH₃), 12.3 (1CH₃).

1-Ethyl 2-[(1R,5S,6S)-(6-cyanomethyl)-3-ethylbicyclo[3.2.0]hept-3-en-6-yl]propanedioate (3:2 mixture of diastereomixture) (**11**). Colorless oil; ¹H NMR (CDCl₃) δ 7.85 (br s, 1H), 5.34 [s, (2/5)1H], 5.27 [s, (3/5)1H], 4.29–4.18 (m, 2H), 3.78 [s, (2/5)1H], 3.77 [s, (3/5)1H], 3.31 (br s, 1H), 3.09–3.01 (m, 1H), 3.01 [s, (2/5)2H], 3.00 [s, (3/5)2H], 2.54 (dd, *J* = 16.0, 7.9 Hz, 1H), 2.36–2.29 (m, 1H), 2.19–2.13 (m, 2H), 2.07 (d, *J* = 16.0 Hz, 1H), 1.74 [dd, *J* = 12.5, 7.5 Hz, (3/5)1H], 1.70 [dd, *J* = 12.5, 7.5 Hz, (2/5)1H], 1.31 [t, *J* = 7.5 Hz, (3/5)3H], 1.28 [t, *J* = 7.5 Hz, (2/5)3H], 1.09 [t, *J* = 7.5 Hz, (2/5)3H], 1.08 [t, *J* = 7.5 Hz, (3/5)3H]; ¹³C NMR (CDCl₃) δ 173.3 [(3/5)1C], 173.0 [(2/5)1C], 167.9 [(2/5)1C], 167.7 [(3/5)1C], 152.3 [(3/5)1C], 152.2 [(2/5)1C], 121.1 [(2/5)1CH], 120.9 [(3/5)1CH], 118.8 (1C), 62.01 [(2/5)1CH₂], 61.96 [(3/5)1CH₂], 55.6 [(3/5)1CH], 55.4 [(2/5)1CH], 55.3 [(2/5)1CH], 55.1 [(3/5)1CH], 43.8 (1C), 42.7 (1CH₂), 38.1 [(3/5)1CH₂], 37.9 [(2/5)1CH₂], 31.6 [(2/5)1CH], 31.5 [(3/5)1CH], 24.6 (1CH₂), 24.54 [(3/5)1CH₂], 24.52 [(2/5)1CH₂], 14.10 [(2/5)1CH₃], 14.07 [(3/5)1CH₃], 12.52 [(2/5)1CH₃], 12.47 [(3/5)1CH₃].

Ethyl (1R,5S,6S)-6-(cyanomethyl)-3-ethylbicyclo[3.2.0]hept-3-ene-6-acetate (**12**). Colorless oil; ¹H NMR (CDCl₃) δ 5.29–5.27 (m, 1H), 4.12 (qd, *J* = 7.2, 2.3 Hz, 2H), 3.13 (br s, 1H), 2.92–2.86 (m, 1H), 2.82 (s, 2H), 2.51 (d, *J* = 2.3 Hz, 2H), 2.50 (dd, *J* = 16.4, 8.7 Hz), 2.17–2.11 (m, 3H), 2.04 (d, *J* = 16.4 Hz, 1H), 1.57 (dd, *J* = 13.0, 7.5 Hz, 1H), 1.26 (t, *J* = 7.2 Hz, 3H), 1.08 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CDCl₃) δ 171.4 (1C), 150.7 (1C), 121.2 (1CH), 118.7 (1C), 60.4 (1CH₂), 54.4 (1CH), 42.2 (1CH₂), 41.6 (1C), 38.8 (1CH₂), 37.9 (1CH₂), 31.0 (1CH), 27.3 (1CH₂), 24.4 (1CH₂), 14.2 (1CH₃), 12.4 (1CH₃).

(1R,5S,6S)-6-(Cyanomethyl)-3-ethylbicyclo[3.2.0]hept-3-ene-6-acetic acid (**13**). White solid; ¹H NMR (CD₃OD-*d*₄) δ 5.36–5.35 (m, 1H), 3.11 (br s, 1H), 2.95–2.87 (m, 1H), 2.91 (s, 2H), 2.51 (dd, *J* = 16.5, 8.0 Hz, 1H), 2.45 (d, *J* = 4.5 Hz, 2H), 2.17–2.11 (m, 1H), 2.17 (q, *J* = 7.5 Hz, 2H), 2.07 (d, *J* = 16.5 Hz, 1H), 1.57 (dd, *J* = 12.6, 7.5 Hz, 1H), 1.11 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CD₃OD-*d*₄) δ 174.8 (1C), 151.6 (1C), 122.7 (1CH), 120.0 (1C), 55.8 (1CH), 43.0 (1CH₂), 42.6 (1C), 39.8 (1CH₂), 38.9 (1CH₂), 32.2 (1CH), 27.5 (1CH₂), 25.3 (1CH₂), 12.8 (1CH₃).

2-[(1R,5S,6R)-6-(Cyanomethyl)-3-ethylbicyclo[3.2.0]hept-3-en-6-yl]propanedioic acid (**14**). White solid; ¹H NMR (CDCl₃) δ 5.38–5.36 (m, 1H), 3.64 (s, 1H), 3.28 (br s, 1H), 3.10 (dd, *J* = 17.0, 0.6 Hz, 1H), 3.04 (d, *J* = 17.0 Hz, 1H), 3.06–3.00 (m, 1H), 2.54 (dd, *J* = 16.6, 8.0 Hz, 1H), 2.31 (ddd, *J* = 12.9, 9.0, 2.3 Hz, 1H), 2.18 (q, *J* = 7.5 Hz, 2H), 2.11 (d, *J* = 16.6 Hz, 1H), 1.73 (dd, *J* = 12.9, 7.5 Hz, 1H), 1.12 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CDCl₃) δ 171.5 (1C), 171.3 (1C), 152.7 (1C), 122.6 (1CH), 120.3 (1C), 57.0 (1CH), 56.5 (1CH), 44.6 (d, *J* = 10.2 Hz, 1C), 43.4 (1CH₂), 39.1 (d, *J* = 6.0 Hz, 1CH₂), 32.7 (1CH), 25.3 (1CH₂), 25.0 (1CH₂), 12.8 (1CH₃).

(1R,5S,6S)-6-(2-Amino-2-oxoethyl)-3-ethylbicyclo[3.2.0]hept-3-ene-6-acetic acid (**8**). White solid; ¹H NMR (CD₃OD-*d*₄) δ 5.39–5.38 (m, 1H), 3.16 (br s, 1H), 2.87–2.81 (m, 1H), 2.67 (d, *J* = 14.0 Hz, 1H), 2.63 (d, *J* = 14.0 Hz, 1H), 2.51 (s, 2H), 2.48 (dd, *J* = 16.3, 7.7 Hz, 1H), 2.21 (ddd, *J* = 12.3, 8.8, 2.6 Hz, 1H), 2.15 (q, *J* = 7.5 Hz, 2H), 2.03 (d, *J* = 16.3 Hz, 1H), 1.49 (dd, *J* = 12.3, 7.5 Hz, 1H), 1.10 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CD₃OD-*d*₄) δ 177.4 (1C), 176.2

(1C), 150.7 (1C), 123.6 (1CH), 56.5 (1CH), 44.1 (1CH₂), 43.1 (1CH₂+1C), 40.3 (1CH₂), 39.5 (1CH₂), 32.4 (1CH), 25.4 (1CH₂), 12.9 (1CH₃).

(1R,5S,6S)-6-(Aminomethyl)-3-ethylbicyclo[3.2.0]hept-3-ene-6-acetic acid (**1** (free form)). White solid; ¹H NMR (CD₃OD-*d*₄) δ 5.37–5.35 (m, 1H), 3.17 (d, *J* = 13.2 Hz, 1H), 3.13 (d, *J* = 13.2 Hz, 1H), 3.09 (br s, 1H), 2.87–2.81 (m, 1H), 2.50–2.45 (m, 1H), 2.49 (d, *J* = 16.0 Hz, 1H), 2.45 (d, *J* = 16.0 Hz, 1H), 2.14 (q, *J* = 7.5 Hz, 2H), 2.07–2.02 (m, 2H), 1.46 (dd, *J* = 12.3, 7.5 Hz, 1H), 1.09 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CD₃OD-*d*₄) δ 180.3 (1C), 151.0 (1C), 122.8 (1CH), 53.7 (1CH), 48.5 (1CH₂), 46.1 (1CH₂), 43.3 (1C), 43.0 (1CH₂), 37.8 (1CH₂), 32.3 (1CH), 25.4 (1CH₂), 12.8 (1CH₃).

Mirogabalin besylate (**1**). White solid; ¹H NMR (CD₃OD-*d*₄) δ 7.85–7.82 (m, 2H), 7.46–7.41 (m, 3H), 5.32 (s, 1H), 3.34 (d, *J* = 13.2 Hz, 1H), 3.29 (d, *J* = 13.2 Hz, 1H), 3.13 (br s, 1H), 2.91–2.84 (m, 1H), 2.52 (dd, *J* = 16.0, 6.9 Hz, 1H), 2.51 (s, 2H), 2.20–2.13 (m, 3H), 2.09 (d, *J* = 16.0 Hz, 1H), 1.50 (dd, *J* = 12.6, 7.5 Hz, 1H), 1.11 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (CD₃OD-*d*₄) δ 175.5 (1C), 152.1 (1C), 146.3 (1C), 131.3 (1CH), 129.3 (2CH), 126.9 (2CH), 122.0 (1CH), 53.7 (1CH), 47.7 (1CH₂), 43.2 (1CH₂), 42.9 (1C), 38.5 (1CH₂), 37.3 (1CH₂), 32.4 (1CH), 25.3 (1CH₂), 12.8 (1CH₃).

(4E)-3-Oxo-5-phenyl-4-pentenitrile (**16a**). Off-white solid; ¹H NMR (CDCl₃) δ 7.69 (d, *J* = 16.0 Hz, 1H), 7.61–7.58 (m, 2H), 7.49–7.42 (m, 3H), 6.88 (d, *J* = 16.0 Hz, 1H), 3.73 (s, 2H); ¹³C NMR (CDCl₃) δ 186.3 (1C), 146.6 (1CH), 133.3 (1C), 131.7 (1CH), 129.2 (2CH), 128.8 (2CH), 122.3 (1CH), 114.0 (1C), 30.8 (1CH₂).

1,3-Diethyl 2-(2-cyano-1-phenylethyl)propanedioate (**17b**). Colorless oil; ¹H NMR (CDCl₃) δ 7.35–7.27 (m, 5H), 4.25 (q, *J* = 7.2 Hz, 2H), 3.99–3.93 (m, 2H), 3.86 (d, *J* = 10.3 Hz, 1H), 3.73 (ddd, *J* = 10.3, 8.2, 4.6 Hz, 1H), 2.92 (dd, *J* = 16.9, 8.2 Hz, 1H), 2.86 (dd, *J* = 16.9, 4.6 Hz, 1H), 1.29 (t, *J* = 7.2 Hz, 3H), 0.99 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (CDCl₃) δ 167.6 (1C), 166.9 (1C), 137.9 (1C), 128.9 (2CH), 128.2 (1CH), 127.7 (2CH), 117.6 (1C), 62.1 (1CH₂), 61.7 (1CH₂), 55.9 (1CH), 41.2 (1CH), 22.8 (1CH₂), 14.0 (1CH₃), 13.7 (1CH₃).

Ethyl 3-methyl-2-(2-cyanoacetyl)-2-butenolate (**16c**). Colorless oil; ¹H NMR (CDCl₃) δ 4.29 (q, *J* = 7.2 Hz, 2H), 3.68 (s, 2H), 2.22 (s, 3H), 2.02 (s, 3H), 1.34 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (CDCl₃) δ 189.8 (1C), 164.3 (1C), 160.0 (1C), 129.0 (1C), 113.6 (1C), 61.4 (1CH₂), 32.6 (1CH₂), 24.4 (1CH₃), 23.5 (1CH₃), 14.1 (1CH₃).

1,3-Diethyl 2-(2-cyano-1,1-dimethylethyl)propanedioate (**17c**). Colorless oil; ¹H NMR (CDCl₃) δ 4.21 (q, *J* = 7.2 Hz, 4H), 3.41 (s, 1H), 2.81 (s, 2H), 1.28 (t, *J* = 7.2 Hz, 6H), 1.27 (s, 6H); ¹³C NMR (CDCl₃) δ 167.5 (2C), 118.0 (1C), 61.5 (2CH₂), 58.9 (1CH), 35.0 (1C), 27.9 (1CH₂), 25.5 (2CH₃), 14.0 (2CH₃).

1,3-Diethyl 2-(1-methylethenyl)propanedioate (**18c**). Colorless oil; ¹H NMR (CDCl₃) δ 5.09–5.08 (m, 1H), 4.99 (br s, 1H), 4.27–4.18 (m, 4H), 4.09 (s, 1H), 1.90–1.89 (m, 3H), 1.29 (t, *J* = 7.2 Hz, 6H); ¹³C NMR (CDCl₃) δ 167.9 (2C), 137.6 (1C), 117.1 (1CH₂), 61.5 (2CH₂), 59.4 (1CH), 20.9 (1CH₃), 14.0 (2CH₃).

1,3-Diethyl 2-(2-cyano-1-methylethyl)propanedioate (**17d**). Colorless oil; ¹H NMR (CDCl₃) δ 4.26–4.19 (m, 4H), 3.39 (d, *J* = 7.5 Hz, 1H), 2.66–2.53 (m, 3H), 1.288 (t, *J* = 7.2 Hz, 3H), 1.285 (t, *J* = 7.2 Hz, 3H), 1.20 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (CDCl₃) δ 167.71 (1C), 167.68 (1C), 118.0 (1C), 61.7 (2CH₂), 55.5 (1CH), 30.2 (1CH), 22.2 (1CH₂), 17.3 (1CH₃), 14.00 (1CH₃), 13.98 (1CH₃).

(4E)-4-Methyl-3-oxo-4-hexenenitrile (**16e**). ¹H NMR (CDCl₃) δ 6.74 (qq, *J* = 6.9, 1.4 Hz, 1H), 3.80 (s, 2H), 1.94 (dq, *J* = 6.9, 1.4 Hz, 3H), 1.83 (q, *J* = 1.4 Hz, 3H); ¹³C NMR (CDCl₃) δ 188.1 (1C), 141.4 (1CH), 137.0 (1C), 114.4 (1C), 28.2 (1CH₂), 15.1 (1CH₃), 11.0 (1CH₃).

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the Publications website.

General remarks, HPLC conditions, Summary of synthetic route, Knoevenagel condensation, Stereoselective 1,4-addition of lithioacetonitrile, Synthesis of diastereomer for *epi*-**10**, Optimization of decarboxylation, One-pot transformation including decarboxylation, hydrolysis, hydration, Hofmann rearrangement, Salt formation, Investigation of the selectivity of lithioacetonitrile, Competition experiments, Characterization of compounds, References for supporting information, ¹H and ¹³C NMR spectra of compounds (PDF)

AUTHOR INFORMATION

Corresponding Author

Hidegori Ochiai - Pharma & Supplemental Nutrition Solutions Vehicle, Pharma Business Division, Kaneka Corporation, 1-8, Miyamae-cho, Takasago-cho, Takasago, Hyogo 676-8688, Japan; orcid.org/0000-0003-1544-9663;

Email: Hidegori.Ochiai@kaneka.co.jp

Authors

Taiki Mihara - Pharma & Supplemental Nutrition Solutions Vehicle, Pharma Business Division, Kaneka Corporation, 1-8, Miyamae-cho, Takasago-cho, Takasago, Hyogo 676-8688, Japan; Email: Taiki.Mihara@kaneka.co.jp

Miwa Sasagawa - Pharma & Supplemental Nutrition Solutions Vehicle, Pharma Business Division, Kaneka Corporation, 1-8, Miyamae-cho, Takasago-cho, Takasago, Hyogo 676-8688, Japan; Email: Miwa.Sasagawa@kaneka.co.jp

Akira Nishiyama - Pharma & Supplemental Nutrition Solutions Vehicle, Pharma Business Division, Kaneka Corporation, 1-8, Miyamae-cho, Takasago-cho, Takasago, Hyogo 676-8688, Japan; orcid.org/0000-0002-3846-0411; Email: Akira.Nishiyama@kaneka.co.jp

Author Contributions

H.O. directed this project and designed the experiments. H.O., T.M., and M.S. conducted experiments. H.O. drafted and edited manuscript and supporting information. A.N. supervised. All authors have read and given approval to the final version of the manuscript.

Funding Sources

There are no funders to report for this submission.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

The authors express our special thanks to Mr. Takuro Yamakawa, Mr. Shunichi Kubota, and Ms. Yuka Morishige for their technical support. The authors are also grateful to members of API team in KANEKA for material supports and helpful discussions.

ABBREVIATIONS

LDA, Lithium diisopropylamide;
LiTMP, lithium tetramethylpiperidide;
LHMDS, Lithium hexamethyldisilazide;

CPME, Cyclopentyl methyl ether;
MTHP, 4-Methyltetrahydropyran;
MTBE, Methyl *tert*-butyl ether.

REFERENCES

- (1) Development of Efficient Manufacturing Method for Mirogabalin Utilizing Chiral Organocatalyst. Ukai, K.; Kaneda, T.; Kitawaki, T.; Wakayama, M.; Nakamura, Y. *Journal of Synthetic Organic Chemistry, Japan* **2024**, *82*, 328–335. <https://doi.org/10.5059/yukigoseikyokaiishi.82.328>
- (2) Nakamura, Y.; Furuya, Y.; Shimada, K.; Onishi, Y. Optical resolution method for bicyclic compound using enzyme. WO 2013154066 A1, 2013.
- (3) Nakamura, Y.; Kitawaki, T. Optical resolution method for bicyclic compound using hydrosulfonic acid. WO 2013089189 A1, 2013.
- (4) Nakamura, Y.; Kitawaki, T.; Kaneda, T. Optical resolution method for bicyclic compound using asymmetric catalyst. WO 2013089188 A1, 2013.
- (5) Nakamura, Y.; Kimura, K. Method for producing bicyclic compound via iminium salt. WO 2012169474 A1, 2012.
- (6) Nakamura, Y. Method for producing bicyclic compound via Claisen rearrangement. WO 2012169475 A1, 2012.
- (7) (a) Aller, B. V. Raney cobalt hydrogenation catalysts. I. the preparation of the catalyst. *J. appl. Chem.* **1957**, *7*, 130–134. <https://doi.org/10.1002/jctb.5010070307> (b) Aller, B. V. Raney cobalt hydrogenation catalysts. II. The physical and chemical properties of the catalyst. *J. appl. Chem.* **1958**, *8*, 163–167. <https://doi.org/10.1002/jctb.5010080304> (c) Aller, B. V. Raney cobalt hydrogenation catalysts. III. Applications and promoter effects. *J. appl. Chem.* **1958**, *8*, 492–495. <https://doi.org/10.1002/jctb.5010080805>
- (8) Nakamura, Y.; Ukai, K.; Kitawaki, T.; Nakajima, T.; Kitagawa, Y.; Furuya, Y.; Imai, M.; Numagami, E.; Wakayama, M.; Saito, A. Method for producing optically active bicyclic γ -amino acid derivative. WO 2015005298 A1, 2015.
- (9) For pregabalin, see: Gade, S. R.; Harikeerthi, N. M.; Amirisetty, R. T.; Chitta, V. W.; Muvva, V. Process for preparing pregabalin. WO 2008137512 A2, 2008. For baclofen, see: (a) Heinrich, K.; Johann, W. F.; Max, W. Gamma-amino-beta-(para-halophenyl)-butyric acids and their esters. US3471548A, 1969. (b) Heinrich, K.; Johann, W. F.; Max, W. Beta-(para-halo-phenyl)-glutaric acid imides. US3634428A, 1972.
- (10) For example, it is known that 1,2-addition occurs for methyl cinnamate with lithioacetonitrile, see; Mamuye, A. D.; Castoldi, L.; Azzena, U.; Holzer, W.; Pace, V. Chemoselective efficient synthesis of functionalized β -oxonitriles through cyanomethylation of Weinreb amides. *Org. Biomol. Chem.* **2015**, *13*, 1969–1973. <https://doi.org/10.1039/C4OB02398F>
- (11) Sumrell, G. Investigation of the reaction of organolithium reagents with nitriles having α -hydrogen. *J. Org. Chem.* **1954**, *19*, 817–819. <https://doi.org/10.1021/jo01370a020>
- (12) Reynolds, G. A.; Humphlett, W. J.; Swamer, F. W.; Hauser, C. R. Condensations of nitriles having α -hydrogen to form β -iminonitriles, cyclic trimers, and substituted 2-hydroxy-4-aminopyridines. *J. Org. Chem.* **1951**, *16*, 165–172. <https://doi.org/10.1021/jo01142a001>
- (13) For generation of lithioacetonitrile with *n*-BuLi and MeCN, see: (a) Kaiser, E. M.; Hauser, C. R. Ionization of an α -hydrogen of acetonitrile by *n*-butyllithium and alkali amides. Condensations with ketones and aldehydes to form β -hydroxynitriles. *J. Org. Chem.* **1968**, *33*, 3402–3404. <https://doi.org/10.1021/jo01273a009> (b) Crouse, D. N.; Seebach, D. Erzeugung und Reaktionen von *a*-Li-N,N-Dimethyl-acetamid und -propionamid und von *a*-Li-Acetonitril. *Chem. Ber.* **1968**, *101*, 3113–3121. *Chem. Ber.* **1968**, *101*, 3113–3121.

- (14) For example of addition to β -substituted methylenemalonate, see; Stanetty, P.; Kremslehner, M.; Voellenkle, H. Addition of Lithiated Acetonitrile to β -Substituted Methylenemalonates. Revision and Correction. *Monatshfte fuer Chemie* **1998**, *129*, 291–296. <https://doi.org/10.1007/pl000000087>
- (15) Selected examples of reactions with lithioacetonitrile at higher temperature, see (a) Bonomo, L.; Solari, E.; Scopelliti, R.; Floriani, C.; Re, N. The Porphyrinogen–Porphodimethene Relationship Leading to Novel Synthetic Methodologies Focused on the Modification and Functionalization of the Porphyrinogen and Porphodimethene Skeletons. *J. Am. Chem. Soc.* **2000**, *122*, 5312–5326. <https://doi.org/10.1021/ja000253s>. (b) Eagon, S.; Ball-Jones, N.; Haddenham, D.; Saavedra, J.; DeLieto, C.; Buckman, M.; Singaram, B. Enantioselective reduction of α -substituted ketones mediated by the boronate ester TarB-NO₂. *Tetrahedron Lett.* **2010**, *51*, 6418–6421. <https://doi.org/10.1016/j.tetlet.2010.09.146>
- (16) A similar phenomenon where dicarboxylic acid do not participate in decarboxylation, was reported by Daiichi-Sankyo, see, ref 1.
- (17) Recent examples of nitrile hydrations, see; (a) Kalimani, F. M.; Khorshidi, A. Ag-embedded manganese oxide octahedral molecular sieve (Ag-OMS-2) nano-rods as efficient heterogeneous catalysts for hydration of nitriles to amides in aqueous solution. *RSC Advances* **2023**, *13*, 6909–6918. <https://doi.org/10.1039/d3ra00292f>. (b) Wang, K.; Jiang, P.; Yang, M.; Ma, P.; Qin, J.; Huang, X.; Ma, L.; Li, R. Metal-free nitrogen -doped carbon nanosheets: a catalyst for the direct synthesis of imines under mild conditions. *Green Chem.* **2019**, *21*, 2448–2461. <https://doi.org/10.1039/c9gc00908f>. (c) Vyas, K. M.; Mandal, P.; Singh, R.; Mobin, S. M.; Mukhopadhyay, S. Arene-ruthenium(II)-phosphine complexes: Green catalysts for hydration of nitriles under mild conditions. *Inorg. Chem. Commun.* **2020**, *112*, 107698. <https://doi.org/10.1016/j.inoche.2019.107698>. (d) Gonzalez-Fernandez, R.; Crochet, P.; Cadierno, V.; Menendez, M. I.; Lopez, R. Phosphinous Acid-Assisted Hydration of Nitriles: Understanding the Controversial Reactivity of Osmium and Ruthenium Catalysts. *Chem. Eur. J.* **2017**, *23*, 15210–15221. <https://doi.org/10.1002/chem.201703481>. (e) Tomas-Mendivil, E.; Cadierno, V.; Menendez, M. I.; Lopez, R. Unmasking the Action of Phosphinous Acid Ligands in Nitrile Hydration Reactions Catalyzed by Arene-Ruthenium(II) Complexes. *Chem. Eur. J.* **2015**, *21*, 16874–16886. <https://doi.org/10.1002/chem.201503076> (f) Woo, H.; Lee, K.; Park, S.; Park, K. H. Magnetically separable and recyclable Fe₃O₄-supported Ag nanocatalysts for reduction of nitro compounds and selective hydration of nitriles to amides in water. *Molecules* **2014**, *19*, 699–712. <https://doi.org/10.3390/molecules19010699>. (g) Battilocchio, C.; Hawkins, J. M.; Ley, S. V. Mild and Selective Heterogeneous Catalytic Hydration of Nitriles to Amides by Flowing through Manganese Dioxide. *Org. Lett.* **2014**, *16*, 1060–1063. <https://doi.org/10.1021/ol403591c>. (h) Garcia-Alvarez, R.; Zablocka, M.; Crochet, P.; Duhayon, C.; Majoral, J.-P.; Cadierno, V. Thiazolyl-phosphine hydrochloride salts: effective auxiliary ligands for ruthenium-catalyzed nitrile hydration reactions and related amide bond forming processes in water. *Green Chem.* **2013**, *15*, 2447–2456. <https://doi.org/10.1039/c3gc41201f>.
- (18) (a) Radziszewski, B. Ueber die Oxidation mittelst Wasserstoffperoxyd. *Ber. Dtsch. Chem. Ges.* **1885**, *18*, 355–356. <https://doi.org/10.1002/cber.18850180171>. (b) Panzarasa, G. Just add luminol to turn the spotlight on Radziszewski amidation. *ACS Omega* **2018**, *3*, 13179–13182. <https://doi.org/10.1021/acsomega.8b02208>. (c) Casella, G.; Carlotto, S.; Ribeiro, A. P. C.; Martins, L. M. D. R. S. The reaction of acetonitrile with hydrogen peroxide in alkaline medium: a DFT mechanistic study of green production of amides. *Phys. Chem. Chem. Phys.* **2023**, *25*, 22775–22781. <https://doi.org/10.1039/d3cp02024j>.
- (19) The competition experiment between **15a** and **15b**, showed that 1,4-adduct was obtained in preference to 1,2-adduct (for detail, see Supporting Information S9).

